



HAL
open science

A new sulfonylated flavonoid and other bioactive compounds isolated from the aerial parts of *Cotula anthemoides* L.

Wafa Tadrent, Abdulmagid Alabdul Magid, Ahmed Kabouche, Dominique Harakat, Laurence Voutquenne-Nazabadioko, Zahia Kabouche

► To cite this version:

Wafa Tadrent, Abdulmagid Alabdul Magid, Ahmed Kabouche, Dominique Harakat, Laurence Voutquenne-Nazabadioko, et al.. A new sulfonylated flavonoid and other bioactive compounds isolated from the aerial parts of *Cotula anthemoides* L.. *Natural Product Research*, 2016, 31 (12), pp.1437-1445. 10.1080/14786419.2016.1261342 . hal-01834071

HAL Id: hal-01834071

<https://hal.univ-reims.fr/hal-01834071v1>

Submitted on 5 Nov 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

A new sulfonylated flavonoid and other bioactive compounds isolated from the aerial parts of *Cotula anthemoides* L.

Wafa Tadrent^a, Abdulmagid Alabdul Magid^{b*}, Ahmed Kabouche^a, Dominique Harakat^c, Laurence Voutquenne-Nazabadioko^b and Zahia Kabouche^{a*}

^aUniversité des frères Mentouri Constantine, Département de chimie, Laboratoire d'Obtention des Substances Thérapeutiques (L.O.S.T), Campus Chaabet-Ersas, 25000 Constantine, Algeria.

^bGroupe Isolement et Structure, Institut de Chimie Moléculaire de Reims (ICMR), CNRS UMR 7312, UFR de Pharmacie, BP 1039, 51687 Reims, France.

^cService Commun d'Analyses, Institut de Chimie Moléculaire de Reims (ICMR), CNRS UMR 7312, Bât. 18 B.P.1039, 51687 Reims, France.

*Corresponding author: Prof. Zahia KABOUCHE; E-mail: zkabouche@yahoo.com; Tel/fax: 213-31811100

Abstract

A new sulfonyl flavonol glucoside, quercetin 6'-[propanoic acid-(2-acetoxy-3-sulfonyl)]-3'-O-β-D-glucopyranoside (**1**) was isolated from the aerial parts of *Cotula anthemoides* L. in addition to fifteen known compounds (**2-16**). The structure elucidation of these compounds was based on analyses of spectroscopic data including 1D-, 2D-NMR and HR-ESI-MS techniques and by comparing their NMR data with those reported in the literature. These compounds were evaluated for their DPPH radical scavenging and tyrosinase inhibitory activity. Compound **6** showed a high DPPH radical scavenging with EC₅₀ value of 9.1 ± 0.4. Compound **11**, **9** and **1** exhibited a mild tyrosinase inhibitory activity with IC₅₀ values of 85 ± 0.8, 95 ± 1.5 and 100 ± 0.5, respectively.

Keywords: *Cotula anthemoides*, Asteraceae, antioxidant activity, tyrosinase inhibitory activity.

1. Introduction

The genus *Cotula* (Asteraceae) consists of approximately 80 species, among them, three are distributed in Algeria (Ozenda, 1958). Plants of this genus are used in traditional medicine for their anti-inflammatory, analgesic, antiseptic properties (Jana et al., 1992) and as aromatic and digestive substance in tea (Bellakhdar, 1997). In addition, it has been reported as antipyretic (Larhsini et al., 2002), bacteriostatic (Jana et al., 1992), antiprotozoal (Markouk et al., 1999) and anticorrosive (Benmenine et al., 2011). Several flavonoids (Ahmed et al., 1987; Mahran et al., 1976), alkaloids (Mahjoub et al., 2012), sesquiterpene lactones (Jakupovitch et al., 1988; Metwalli et al., 1986) and coumarins (Greger et al., 1985) were isolated from this genus. *Cotula anthemoides* L commonly known as 'Babunaj' in Algeria is used for colic and as a remedy for head and chest colds (Abhay and Tripathi, 2011). Previous phytochemical investigation on *C. anthemoides* led to the isolation of coumarin derivatives: 6-methoxy-2-oxo-2H-chromene-8-carboxylic acid methyl ester, α-pinene, ursolic acid and gibberellic acid (Showkat, 2012), 5-hydroxy-6,3',4'-trimethoxyisoflavone-7-O-β-D-glucopyranosyl(1→6)-α-L-rhamnopyranoside and 5,7-dihydroxy-6,8-dimethyl-3-

methoxyflavone-7-O-β-D-galactopyranosyl-(1→2)-α-L-rhamnopyranoside (Yadava and Barsainya, 1998). The present work describes the isolation and structural identification of sixteen compounds from *C. anthemoides* that grows wild in Algerian Sahara; fourteen are reported here for the first time from the aerial parts of *Cotula anthemoides* L. In addition, the radical scavenging ability (DPPH assay) and the tyrosinase inhibitory activity of all compounds were investigated for the first time, from this plant.

2. Results and discussion

The *n*-BuOH extract of the aerial parts of *C. anthemoides* was separated by combined chromatographic methods to obtain a new sulfonyl flavonol glucoside (**1**) (Fig. 1), in addition to fifteen known compounds (**2-16**) which were elucidated as luteolin-7-O-β-D-glucopyranoside (**2**) (Furkan et al., 2012), vicenin 2 (**3**) (Yinrong and Yeap foo, 2000), luteolin-7-O-β-D-glucuronopyranoside (**4**) (Tsukasa et al., 2011), apigenin-7-O-β-D-glucuronopyranoside (**5**) (Xiao et al., 2006), 6-hydroxyluteolin-7-O-β-D-glucopyranoside (**6**) (Yinrong and Yeap foo, 2000), isoetin-5'-O-β-D-glucopyranoside (**7**) (Alberto et al., 1988), tachioside (**8**) (Shogo et al.,

1987), pseudolaroside A (**9**) (Peng et al., 2006), junipediol A 8-O- β -D-glucopyranoside (**10**) (Gilles et al., 1997; Arunee et al., 2014), vanillic acid-4-O- β -D-glucopyranoside (**11**) (Akiyo et al., 1995), 2-hydroxy-3-phenylpropionamide (**12**) (Zhi-gang et al., 2012), protocatechuic acid (**13**) (Hui et al., 1998), uridine (**14**) (Yalçin et al., 2003), tuberonic acid β -D-glucopyranoside (**15**) (Baoliang et al., 1993) and rosmarinic acid (Gohari et al., 2009) (Fig.S1). Their structural assignments were made by HR-ESI-MS, 1D-, and 2D-NMR analyses and by comparison with spectral data from the literature values. To the best of our knowledge, compounds **3-16** were isolated for the first time from the genus *Cotula*.

Compound **1** was isolated as a yellow amorphous powder. The positive HR-ESI-MS showed a molecular ion peak m/z 681.0744 $[M+Na]^+$ (calcd for $C_{26}H_{26}O_{18}NaS$, 681.0744) enabling to determine the molecular formula $C_{26}H_{26}O_{18}S$. The UV spectrum revealed absorption bands at 206, 260 and 350 nm. The 1H and ^{13}C NMR spectra of **1** showed characteristic signals indicating the presence of one sugar unit and one flavonol moiety (Ozbek et al., 2015). Signals arising from one anomeric proton at δ_H 5.07 (1H, d, $J = 7.4$ Hz) and other protons of sugar moiety in the range of (δ_H 3.5-4.0 ppm) showed the presence of d-glucose with β -configuration on the basis of the large coupling constants between H-1'' to H-5'' (Table 1) (Alabdul Magid et al., 2008). Complete assignments of osidic protons and carbons were determined by the analysis of COSY, HSQC and HMBC experiments. The relative configuration of β -D-glucopyranosyl moiety was further confirmed by analysis of the ROESY correlations from the α -axial protons H-1/H-3 and H-1/H-5. The A-ring of the flavonol was represented by two meta-coupled resonances at δ_H 6.26 (1H, brs) and δ_H 6.38 (1H, brs) assigned to H-6 and H-8, respectively. The 1H NMR data (see experimental part) exhibited two singlet signals at δ_H 7.43 (1H, s) and δ_H 7.85 (1H, s) assignable to H-5' and H-2', respectively, which were correlated in the HSQC spectrum with their aromatic carbons at δ_C 119.1 and δ_C 113.9 ppm, respectively. In addition, one methyl proton appeared at δ_H 1.90 (3H, s) together with a set of three aliphatic protons δ_H 3.45 (1H, dd, $J = 14.1, 6.8$ Hz), 3.56 (1H, m) and 5.42 (1H, dd, $J = 6.5, 4.1$ Hz). The 1H and ^{13}C NMR signals of compound **1** were completely assigned by a combination of HSQC, HMBC and 1H - 1H COSY experiments. The 1H - 1H COSY correlation between H₂-3''' (δ_H 3.45, 3.56) and H-2''' (δ_H 5.42) and the cross-peaks in the HMBC spectrum (Fig. S17) between H₂-3'''/C-2''' (δ_C 67.1) and C-1''' (δ_C 171.5), H-2'''/C-1''' (δ_C 57.2), and C-4''' (δ_C 171.3) and H₃ (δ_H 1.90)/C-4''', indicated the presence of propanoic acid-2-acetoxy-3-sulfonyl moiety. The chemical shift of C-3''' (δ_C 57.2) in addition to the molecular formula indicated a sulfonyl group. The site of linkage of the glucopyranosyl and sulfonyl groups in **1** was determined by cross-peaks observed in the HMBC experiment between H-1'' of glucose (δ_H 5.07) and C-3' (δ_C 148.9) of quercetin, and

between H₂-3''' (CH_2SO_2) of sulfonyl group and C-6' (δ_C 133.6) of quercetin. Thus, the structure of compound **1** was characterised as quercetin 6'-[propanoic acid-(2-acetoxy-3-sulfonyl)]-3'-O- β -D-glucopyranoside.

Several biological activities of extracts and essential oils of *Cotula* were reported however, the biological potential of isolated pure compounds has not been investigated. Thus, the isolated compounds **1-16** were tested for their DPPH radical scavenging and anti-tyrosinase activities.




Figure 1: Chemical structure of compound **1** isolated from *C. anthemoides* L.

In order to assess the antioxidative properties of **1-16**, the DPPH radical scavenging activity of these compounds was measured. According to the results (Table 2), compounds **1, 2, 4-6, 8,** and **16** have a slightly radical scavenging potential (EC_{50} ranging from 9.1 to 29.9 μM) compared with ascorbic acid, used as positive control (EC_{50} 7.4 μM). The di-OH substitution at 3' and 4' in the B ring is particularly important to the antiradical activity of flavonoids (Arora et al.1998). The trend is consistent with the less active flavonoid **5** compared with **4**. The free 3-OH was also found to be important for the antioxidant activity of rosmarinic acid **16** (3-OH) comparing to tachioside **8** (3-OCH₃) (Sawai and Moon, 2000). Comparison of the DPPH radical scavenging activity of **6** with **1, 2** and **4** showed that the presence of the hydroxyl group at C-6 increased the activity. Compounds **2** and **4** shared a common aglycone which is 3' and 4'-di-OH substituted in the B ring (luteolin skeleton), the only difference was in the nature of the group linked at C-7 which was β -D-glucopyranose (**2**) and β -D-glucuronopyranose (**4**), suggesting that the glucuronic acid may contribute slightly to the increase of the DPPH radical scavenging activity. Thus, we can conclude that the hindrance effect due to the presence of a hydroxy group at C-6, a β -D-glucose or β -D-glucuronic acid linked at C-7 in a luteolin skeleton is determinant in the scavenging of the DPPH free radical.

The tyrosinase inhibitory activity of isolated pure compounds **1-16** was performed using L-DOPA as substrate and kojic acid, a well-known strong tyrosinase inhibitor, as a positive control (Demirkiran et al. 2013;

Kim et al., 2001; Kim and Uyama, 2005; Xie et al., 2003). The results (Table 2) showed that compounds **1**, **9** and **11** exhibited a mild tyrosinase inhibition. The most active compound was **11** (IC₅₀ 85 μM) followed by **9** (IC₅₀ 95 μM) and **1** (IC₅₀ 100 μM).

Table 1. Antiradical potential and anti-tyrosinase activity of compounds **1-16** isolated from the aerial parts of *C. anthemoides* L.

Compounds	DPPH radical scavenging activity	Tyrosinase
	EC ₅₀ ± S.D. (μM) ^a	inhibitory activity IC ₅₀ ± S.D. (mM) ^a
1	22.8 ± 0.5	100 ± 0.5
2	29.9 ± 1.3	- ^b
3	- ^b	- ^b
4	25.1 ± 0.7	- ^b
5	29.5 ± 0.1	- ^b
6	9.1 ± 0.4	- ^b
7	- ^b	- ^b
8	27.6 ± 1.1	- ^b
9	- ^b	95 ± 1.5
10	- ^b	- ^b
11	- ^b	85 ± 0.8
12	- ^b	- ^b
13	- ^b	- ^b
14	- ^b	- ^b
15	- ^b	- ^b
16	12.7 ± 0.3	- ^b
Ascorbic acid ^c	7.4 ± 0.05	-
Kojic Acid ^c		6.4 ± 0.04

^a Values are presented as mean ± S.D. (n = 3).

^b 50% inhibition not achieved at the concentration of 100 μg/ml.

^c used as a positive control.

In summary, one new sulfonyl flavonol glucopyranoside, along with fifteen known compounds, were isolated from the aerial parts of *C. anthemoides*. The DPPH radical scavenging activity assay showed a high activity for compound **6** compared with ascorbic acid used as a positive control. Compounds **1**, **9** and **11** exhibited a mild *in vitro* mushroom tyrosinase inhibition compared with kojic acid, used as a positive control.

3. Experimental

3.1. General experimental procedures

NMR spectra were carried in MeOH-*d*₄ and DMSO on Bruker Avance DRX III 500 instruments. HR-ESI-MS experiments were performed using a Micromass Q-TOF micro instrument. Biological assays were read on a Fluostar Omega microplate reader (BMG labtech), the amounts for DPPH assay were 100 μL (95 μL DPPH solution and 5 μL sample) and for tyrosinase inhibitory activity are 300 μL (100 μL of sample, 100 μL of mushroom tyrosinase solution and 100 μL of L-dopa). 96-well polystyrene microtiter clear plates were used. Thin layer chromatography (TLC) was performed on pre-coated silica-gel 60 F₂₅₄ Merck and compounds were observed under UV light at 254 and 365 nm or visualized by spraying the dried plates with 50% H₂SO₄, followed by heating. CC was carried out on Kieselgel 60 (63-200 mesh) or LiChroprep RP-18 (40-63 mm) Merck. HPLC was performed on a Dionex apparatus equipped with an ASI-100 autosampler, an Ultimate 3000 pump, a diode

array detector UVD 340S and Chromeleon software. RP-18 column (Phenomenex 250x15 mm, Luna 5μ) was used for semi-preparative HPLC with binary gradient eluent (H₂O (pH 2.4 with trifluoroacetic acid (TFA)); MeCN) and a flow rate of 4 mL/min; the chromatogram was monitored at 205, 210, 254, and 360 nm. Absorbance (Ab) values in the DPPH free radical scavenging and antityrosinase assay were read on a Fluostar omega microplate reader (BMG labtech).

3.2. Plant material

The aerial parts of *C. anthemoides* L. were collected in March 2012 from Tinguentourine, Ain Amenas, Eastern Algerian Sahara. The plant has been authenticated by Prof. Gérard De Belair (Annaba University, Algeria) and a voucher specimen (LOST Ca03/12) was deposited at the herbarium of the Laboratory of Therapeutic Substances (LOST), Faculty of Sciences, Université des Frères Mentouri-Constantine-Algeria.

3.3. Extraction and isolation

The powdered dry aerial parts of *C. anthemoides* (1400 g) were macerated at room temperature in methanol 80% (3x7.5 L, 24h). The extract was concentrated under low pressure, diluted with H₂O (560 mL), then successively extracted with petroleum ether (PE) (3x300 mL), chloroform (3x300 mL), ethyl acetate (3x300 mL), and *n*-butanol (3x300 mL). After evaporation of the solvents, 0.4g of PE, 1.5 g of CHCl₃, 0.55 g of EtOAc and 21.7 g of *n*-BuOH extracts were obtained. The *n*-BuOH extract was subjected to VLC over polyamid SC6 (9 cm x 5 cm) eluted with toluene-methanol with increasing polarity to give 12 fractions (F1-F12 respectively). Fractions F6-7 were combined (1.4 g) and subjected to HPLC preparative using (10-25% MeCN in 60 min) to yield 19 fractions. Frs [4-6] (35.5 mg) were purified by semi-prep. HPLC (5-15% MeCN, in 20 min) affording compounds **14** (R_t 9.9 min, 4 mg) and **8** (R_t 13.8 min, 3 mg). Frs [7-10] (36 mg) were purified by semi-prep. HPLC (5-15% MeCN, in 20 min) yielding compounds **9** (R_t 16.5 min, 7 mg) and **10** (R_t 21.5 min, 4 mg). Frs [11-12] (26.7mg) were purified by semi-prep. HPLC (10-25% MeCN, in 20 min) leading to compound **11** (R_t 6.3 min, 3.2 mg). Frs [26-27] (51 mg) were purified by semi-prep. HPLC (15-35% MeCN, in 20 min) to afford compounds **12** (R_t 8.5 min, 1.6 mg) and **15** (R_t 16.5 min, 4.7 mg). Frs [36-44] (42 mg) were purified by semi-prep. HPLC (10-40% MeCN, in 25 min) yielding compounds **16** (R_t 25.9 min, 4.2 mg). The combined fractions F9-11 (827 mg) were subjected to a preparative HPLC (10-40% MeCN in 60 min) to obtain 35 fractions. Frs [7-13] (50.3 mg) were purified by semi-prep. HPLC (15-35% MeCN, in 20 min) yielding compounds **1** (R_t 13.1 min, 7 mg), **4** (R_t 14.2 min, 5 mg) and **16** (R_t 18.2 min, 1.4 mg). Frs [14-28] (46.2 mg) were purified by semi-prep. HPLC (15-35% MeCN, in 20 min) leading to compounds **3** (R_t 8.0 min, 4.0 mg), **2** (R_t 13.7 min, 3.5 mg) and **5** (R_t 15.1 min, 4.1 mg). Compound **6** (17.1min, 9.2 mg) was purified from

Frs [28-29] by semi-prep. HPLC (15-35% MeCN, in 20 min). Frs [30-32] were purified by semi-prep. HPLC (15-35% MeCN, in 30 min) affording compounds **13** (R_f 6.3 min, 1 mg) and **7** (R_f 26.4 min, 11.2 mg).

3.3.1. Quercetin 6'-[propanoic acid-(2-acetoxy-3-sulfonyl)]-3'-O- β -D-glucopyranoside

Yellow amorphous powder; $[\alpha]_D^{20} + 26.8$ (c 0.25, MeOH). UV (MeOH) λ_{max} (ab.): 206 (2.1), 260 (1.3), 350 (0.55). 1H NMR (500 MHz, CD_3OD): δ_H 6.26 (brs, H-6), 6.38 (brs, H-8), 7.85 (s, H-2'), 7.43 (s, H-5'), 5.07 (d, $J = 7.4$ Hz, H-1"), 3.59 (dd, $J = 8.9, 7.4$ Hz, H-2"), 3.56 (t, $J = 8.6$ Hz, H-3"), 3.52 (t, $J = 8.6$ Hz, H-4"), 3.60 (m, H-5"), 3.81 (dd, $J = 12.2, 4.8$ Hz, H-6"a), 4.00 (dd, $J = 12.2, 2.5$ Hz, H-6"b), 5.42 (dd, $J = 6.5, 4.1$, H- 2"), 3.45 (dd, $J = 14.1, 6.8$, H- 3"a), 3.56 (m, H- 3"b), 1.90 (s, H- 5"). ^{13}C NMR (125 MHz, CD_3OD): δ 145.2 (C-2), 138.9 (C-3), 177.6 (C-4), 162.9 (C-5), 99.8 (C-6), 166.1 (C-7), 94.7 (C-8), 158.4 (C-9), 105.1 (C-10), 125.4 (C-1'), 113.9 (C-2'), 148.9 (C-3'), 151.1 (C-4'), 119.1 (C-5'), 133.6 (C-6'), 103.0 (C-1"), 74.7 (C-2"), 77.6 (C-3"), 71.0 (C-4"), 78.3 (C-5"), 62.2 (C-6"), 57.2 (C-1"), 67.1 (C-2"), 171.5 (C-3"), 171.3 (C-4"), 20.2 (C-5"). HR-ESI-MS $[M+Na]^+$ m/z 681.0744 (calcd for $C_{26}H_{26}O_{18}NaS$, 681.0744).

3.3.2. Acid hydrolysis

A part of the *n*-BuOH extract (300 mg) was refluxed (90 °C) with 10 mL of 2M TFA for 3 h. After extraction with ethyl acetate (3 x 10 mL), the aqueous layer was evaporated to furnish the monosaccharide residue (100 mg). Two sugars were identified as glucose and glucuronic acid by comparison with authentic samples on TLC in MeCOEt:*iso*-PrOH:Me₂CO:H₂O (20:10:7:6). The monosaccharide residue (100 mg) was subjected to a preparative TLC using the same solvent. The optical rotation of each purified sugar was measured and compared with authentic samples to afford D-glucose and D-glucuronic acid.

3.4. Biological activities

3.4.1. DPPH free radical scavenging assay

The scavenging activity of isolated compounds against DPPH was investigated by spectrophotometric methodology, as previously described (Gossan et al.,

2015). Briefly, 5 μ L of either the standard or sample solutions (dissolved in DMSO) was mixed with 95 μ L of DPPH solution (158 μ M, dissolved in absolute EtOH). After mixing gently and incubating for 30 min at 37°C, the optical density was measured at λ 515 nm. The percentage of absorbance inhibition at λ 515 nm was calculated using the following equation: % inhibition $[(Ab_{control} - Ab_{sample})/Ab_{control}] \times 100$. DPPH solution in EtOH was used as a control. The curve of the % scavenging activity against the concentration of sample was prepared by the MSEXcel based program to obtain the EC₅₀. All the tests were conducted in triplicate. The experimental data were expressed as mean \pm standard deviation.

3.4.2. Tyrosinase enzyme assay

Tyrosinase activity inhibition was determined by spectrophotometric methodology, as previously described (Ngoc et al., 2009). 100 μ L of each test compound solution (dissolved in 10% DMSO) was added to 96-well plate then mixed with 100 μ L of mushroom tyrosinase solution (135 U/ml mushroom tyrosinase in phosphate buffer solution (PBS, pH 6.8)). After pre-incubation at 25 °C for 10 min, 100 μ L of L-dopa (0.5 mM, PBS pH 6.8) were added into 96-well plate. The reaction mixture was incubated for another 5 min at 25 °C. The amount of dopachrome in the mixture was determined by the measurement of the absorbance of each well at 475 nm. The inhibitory percentage of tyrosinase was calculated according to the following equation: % inhibition = $\{[(A - B) - (C - D)] / (A - B)\} \times 100$. A: Ab at 475 nm without test substance; B: Ab at 475 nm without test substance and tyrosinase; C: Ab at 475 nm with test substance; D: Ab at 475 nm with test substance, but without tyrosinase. Kojic acid was used as positive control agent. All the tests were conducted in triplicate and IC₅₀ was determined by interpolation of concentration % inhibition curve obtained by MSEXcel based program. The experimental data were expressed as mean \pm standard deviation.

Acknowledgements

The authors are grateful to DGRSDT and ATRSS (MESRS, Algeria), to university Frères Mentouri-Constantine and to Groupe Isolement et Structure of the Institut de Chimie Moléculaire de Reims (ICMR), France for financial support.

Supporting Information

HR-ESI-MS, 1H and ^{13}C NMR, COSY, HMBC, HSQC and ROESY spectra for compound **1**.

References

- Abhay KP, Tripathi NN. 2011. Aromatic plants of Gorakhpur division: Their antimycotic properties and medicinal value. *Int J Pharm Sci Rev Res.* 7 (2): 142-147.
- Ahmed AA, EL-Sayed N, EL-Negoumi S, Mabry JT. 1987. Flavonoids of *Cotula cinerea*. *J Nat Prod.* 50 (3): 519-520.
- Akiyo S, Maksut C, Takashi M. 1995. Hydroxyl benzoic acids from *Boreava orientalis*. *Phytochemistry.* 40 (1): 257-261.

- Alabdul Magid A, Voutquenne-Nazabadioko L, Bontemps G, Litaudon M, Lavaud C. 2008. Tyrosinase inhibitors and sesquiterpene diglycosides from *Guioa villosa*. *Planta Med.* 74: 55–60.
- Alberto Marco J, Barbera O, Rodriguez S, Domingo C, Adell J. 1988. Flavonoids and other phenolics from *Artemisia hispanica*. *Phytochemistry.* 27 (10): 3155–3159.
- Arora A, Nair MG, Strasburg GM. 1998. Structure–activity relationships for antioxidant activities of a series of flavonoids in a liposomal system. *Free Radic Biol Med.* 24: 1355–1363.
- Arunee S, Kaoru U, Hiroshi N, Pharkphoom P. 2014. Identification of a new angiotensin-converting enzyme (ACE) inhibitor from Thai edible plants. *Food Chem.* 165: 92–97.
- Baoliang C, Motoyuki N, Junei K, Toshihiro N. 1993. Chemical constituents of *Astragali semen*. *Chem Pharm Bull.* 41 (1): 178–182.
- Bellakhdar J. 1997. La Pharmacopée marocaine traditionnelle. Vol 1, Ibis press. Paris.
- Benmenine AK, Ouahrani MR, Gherraf N, Sekirifa ML, Baameur L. 2011. Potentiodynamic investigation of the anticorrosive action of *Cotula cinerea* extracts on mild steel X 52 in 20 % H₂SO₄ Solution. *J Chem Pharm Res.* 3 (3) : 666–669.
- Chang TS. 2009. An updated review of tyrosinase inhibitors. *Int J Mol Sci.* 10: 2440–2475.
- Demirkiran O, Sabudak T, Ozturk M, Topcu G. 2013. Antioxidant and tyrosinase inhibitory activities of flavonoids from *Trifolium nigrescens* subsp *petrisavi*. *J Agr Food Chem.* 61: 12598–12603.
- Furkan O, Özlem B, Derya Y, Tuğba B, Zuhul G, Medine G. 2012. Isolation of some luteolin derivatives from *Mentha longifolia* (L.) Hudson sbsps *longifolia* and determination of their genotoxic potencies. *Food Chem.* 135: 764–769.
- Gilles C, Daovy PA, Albert JC, Joseph V, Noel P. 1997. Three phenylpropanoids from *Juniperus phænicea*. *Phytochemistry.* 64 (6): 1169–1173.
- Gohari AR, Saeidnia S, Shahverdi AR, Yassa N, Malmir M, Mollazade K, Naghinejad AR. 2009. Phytochemistry and antimicrobial compounds of *Hymenocrater calycinus*. *Eur Asia J Bio Sci.* 3: 64–68.
- Gossan DPA, Alabdul Magid A, Yao-Kouassi PA, Coffy AA, Harakat D, Voutquenne-Nazabadioko L. 2015. New acylated flavonol glycosides from the aerial parts of *Gouania longipetala*. *Phytochem Lett.* 11: 306–310.
- Greger H, Otmar H. 1985. Sesquiterpene coumarin ethers and polyacetylenes from *Brocchia cinerea*. *Phytochemistry.* 24 (1): 85–88.
- Hui-Li Z, Akito N, Harumi O, Hajim M, Jinsaku S. 1998. Sesquiterpene glycosides from Cotton oil cake. *Phytochemistry.* 48 (4): 665–668.
- Jana M, Lazrek HB, Markouk M. 1992. Effets bactériostatiques des extraits flavoniques de *Cotula cinerea*. *El Bayrounia Rev Mar Pharm.* 8 (2): 89–96.
- Jakupovic J, Aad MA, Eid F, Bohlmann F, Mungay MG. 1988. Further glaucolides and other sesquiterpene lactones from *Brocchia cinerea*. *Phytochemistry.* 27 (7): 2219–2224.
- Kim DS, Kim SY, Moon SJ, Chung JH, Kim KH, Cho KH. 2001. Ceramide inhibits cell proliferation through Akt/PKB inactivation and decreases melanin synthesis in Mel-Ab cells. *Pigm Cell Res.* 14: 110 – 115.
- Kim D, Park J, Kim J, Han C, Yoon J, Kim N. 2006. Flavonoids as mushroom tyrosinase inhibitors: A fluorescence quenching study. *J Agric Food Chem.* 54: 935–941.
- Kim JY, Uyama H. 2005. Tyrosinase inhibitors from natural and synthetic sources: Structure, inhibition mechanism and perspective for the future. *Cell Mol Life Sci.* 62: 1707–1723.
- Larhsini M, Markouk M, Jouhari JT, Bekkouche K, Lazrek HB, Jana M. 2002. The antipyretic activity of some Moroccan medicinal plants. *Phytother Res.* 16 (1): S97–98.
- Mahjoub MA, Ammar S, Majouli K, Mighri Z. 2012. Two new alkaloids and a new polyphenolic compound from *Cotula coronopifolia*. *Chem Nat Compd.* 47 (6): 955–958.
- Mahrán GH, Salah Ahmed M, Ansari SA. 1976. A study of the flavonoid content of *Cotula cinerea*. *Bull Fac Pharm Cairo Univ.* 14 (1): 237–246.
- Markouk M, Lazrek HB, Jana M. 1999. Analgesic effects from *Cotula cinerea*. *Phytother Res.* 13: 229–230.
- Matsuda H, Higashino M, Chen W, Tosa H, Iinuma M, Kubo M. 1995. Studies of cuticle drugs from natural sources. III. Inhibitory effect of *Myrica rubra* on melanin biosynthesis. *Biol Pharm Bull.* 18: 1148–1150.
- Metwally MA, EL-Dahmi S, Jakupovic J, Bohlmann F, Dawidar AM, Metwally SA. 1985. Glaucolide-like sesquiterpene lactones from *Cotula cinerea*. *Phytochemistry.* 25 (1): 255–257.
- Ngoc TM, Lee I, Ha DT, Kim HJ, Min BS, Bae KH. 2009. Tyrosinase-inhibitory constituents from the twigs of *Cinnamomum cassia*. *J Nat Prod.* 72: 1205–1208.
- Ozenda P. 1958. Flore du Sahara septentrional et central, CNRS, Paris.
- Özbek H, Güvenalpa Z, Kuruüzüm-Uzb A, Kazaz C, Ömür Demirezerb L. 2016. β -Hydroxydihydrochalcone and flavonoid glycosides along with triterpene saponin and sesquiterpene from the herbs of *Pimpinella rhodantha* Boiss. *Nat Prod Res.* 30 (7): 750–754.
- Peng L, Hongzhu G, Yin T, Qiao W, Dean G. 2006. Benzoic acid allopyranosides from the bark of *Pseudolarix kaempferi*. *Phytochemistry.* 67: 1395–1398.

- Sawai Y, Moon JH. 2000. NMR analytical approach to clarify the molecular mechanisms of the antioxidant and radical-scavenging activities of antioxidants in tea using 1, 1-diphenyl-2-picrylhydrazyl. *J Agric Food Chem.* 48: 6247–6253.
- Shogo I, Manami S, Hiroshi K, Hideaki O, Kazuo Y. 1987. Aromatic glycosides from *Berchemia racemosa*. *Phytochemistry.* 26 (10): 2811-2814.
- Showkat R. 2012. Chemoprofiling of medicinal plants and value addition of their principle constituents through synthetic modifications. Kashmir, Srinagar.
- Tsukasa I, Hiroaki S, Junichi K. 2011. Flavonoids from the leaves of *Vitex rotundifolia* (Verbenaceae) and their qualitative and quantitative comparison between Coastal and Inland populations. *Bull Natl Mus Nat Sci.* 37 (2): 87–94.
- Xiao JB, Ren FL, Xu M. 2006. Flavones from *Marchantia convoluta*: Isolation of apigenin-7-O- β -D-glucuronide and 5-hydroxyl-7-methoxyl-2-methylchromone. *J Pharm All Sci.* 3 (1): 310-313.
- Xie LP, Chen QX, Huang H, Wang HZ, Zhang RQ. 2003. Inhibitory effects of some flavonoids on the activity of mushroom tyrosinase. *Biochemistry.* 68: 487–491.
- Yadava RN, Barsainya D. 1998. A novel flavone glycoside from the seeds of *Cotula anthemoides*. *Fitoterapia.* 69 (5): 437-438.
- Yalçın FN, Ersöz T, Akbay P, Çaliş İ. 2003. Phenolic, megastigmane, nucleotide, acetophenon and monoterpene glycosides from *Phlomis samia* and *P. carica*. *Turk J Chem.* 27: 703-711.
- Yinrong L, Yeap foo L. 2000. Flavonoids and phenolic glucosides from *Salvia officinalis*. *Phytochemistry.* 55: 263-267.
- Zhi-gang W, Hai-yan X, Qiong M, Ye C, Jian-nan M, Chao-mei M. 2012. Isolation, identification and quantification of unsaturated fatty acids, amides, phenolic compounds and glycoalkaloids from *Potatoes peel*. *Food Chem.* 135: 2425-2429.