



HAL
open science

Three new iridolactone derivatives from the whole plant of *Brillantaisia owariensis* P. Beauv

Perrin Lanversin Foning Tebou, David Ngnokam, Dominique Harakat,
Laurence Voutquenne-Nazabadioko

► To cite this version:

Perrin Lanversin Foning Tebou, David Ngnokam, Dominique Harakat, Laurence Voutquenne-Nazabadioko. Three new iridolactone derivatives from the whole plant of *Brillantaisia owariensis* P. Beauv. *Phytochemistry Letters*, Elsevier, 2018, 25, pp.171-174. 10.1016/j.phytol.2018.04.015 . hal-01996239

HAL Id: hal-01996239

<https://hal.univ-reims.fr/hal-01996239>

Submitted on 8 Nov 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Three new iridolactone derivatives from the whole plant of *Brillantaisia owariensis* P. Beauv

Perrin Lanversin FONING TEBOU^a, David NGNOKAM^{a*}, Dominique HAKAKAT^b and

Laurence VOUTQUENNE-NAZABADIOKO^c

^aFaculty of Science, Department of Chemistry, University of Dschang, P.O. Box 67. Dschang
Cameroon

^bService Commun d'Analyses, Institut de Chimie Moléculaire de Reims (ICMR), CNRS
UMR 7312, Bat. 18 B.P.1039, 51687 Reims Cedex2, France

^cGroupe Isolement et Structure, Institut de Chimie Moléculaire de Reims (ICMR), CNRS
UMR 7312, Bat. 18 B.P.1039, 51687 Reims Cedex2, France

*Corresponding author. E-mail: dngnokam@yahoo.fr/dngnokam@univ-dschang.org

Corresponding authors:

* Professor David Ngnokam, Laboratory of Environmental and Applied Chemistry, Department of Chemistry, Faculty of Science, University of Dschang, P.O. Box 67, Dschang, Cameroon, Tel: +237 696 710 992. E-mail: dngnokam@yahoo.fr/dngnokam@univ-dschang.org

Abstract

Owariensisone B-D, three new lactone iridoids, identified as lactone of α -(3-acetoxymethyl-2-hydromethylcyclopent-3-en-yl) ethanoic acid (**1**), lactone of α -(4- β -acetoxy-3- β -acetoxymethyl-2-hydromethyl-3-hydroxycyclopentyl) ethanoic acid (**2**) and lactone of α -(4- β -acetoxy-3- α -acetoxymethyl-2-hydromethyl-3-hydroxycyclopentyl) ethanoic acid (**3**) were isolated from the *n*-hexane extract of the whole plant of *Brillantaisia owariensis*. Their structures were established by interpretation of their spectral data, mainly HR-TOFESIMS, 1D-NMR (^1H , ^{13}C) and 2D-NMR (^1H - ^1H COSY, HSQC, HMBC and ROESY) and by comparison with the literature data.

Keywords: *Brillantaisia owariensis*, Acanthaceae, Lactone iridoids, Owariensisone.

Introduction

The genus *Brillantaisia* belongs to the family Acanthaceae and is found growing throughout tropical Africa and Madagascar and occupies a prominent position in traditional medicine (Ngbolua et al., 2013). Eleven species are known in Cameroon, and among these, *Brillantaisia owariensis* P. Beauv (Synonym *Brillantaisia patula* T. Anderson), a large erect shrub with a purple-blue flowers (Heine, 1963). It is found growing in Nigeria, Togo, West Cameroon and across Uganda and Angola. The leaves are used for the treatment of anaemia (Ngbolua et al., 2013), rheumatism, menstrual pain, stomach ache and for their antiplasmodial and analgesic potentials (Asai et al., 2012; Makambila-Koubemba et al., 2011; Mbatchi et al., 2006). Previous study indicated that the alcoholic extract has antibacterial and antioxidant activities (Aluko et al., 2014; Faparusi et al., 2012,) but the phytochemical investigation were limited. Our previous contribution reported the isolation of one new lactone iridoid in the methanol extract with flavonoid and glycosides (Foning Tebou et al., 2016). In an extension of our studies, the *n*-hexane soluble part of the methanol extract of the whole plant of *B. owariensis* was examined in greater detail and three novel lactone iridoids (**1-3**) were isolated in minor concentration.

2. Results and discussion

Purification of the *n*-hexane soluble fraction of the crude MeOH extract afforded three new compounds, owariensisone B, C and D (**1-3**) (Fig. 1).

Compound **1** was obtained as yellowish gum. Its molecular formula was determined as $\text{C}_{11}\text{H}_{14}\text{O}_5$ on the basis of its HR-TOFESIMS spectrum, exhibiting a pseudo-molecular ion

peak at m/z 249.0734 $[M+Na]^+$ (calcd. for $C_{11}H_{14}O_5Na$. 249.0739) with five degrees of unsaturation. Its 1H -NMR spectrum (Table 1) shows one olefinic proton at δ_H 5.83 (H-7, brqt, $J = 1.6$ Hz), two pairs of *gem*-coupled methylene protons at δ_H 2.60 and 2.65 (H-6 α and H-6 β), and at δ_H 2.80 and 2.83 (H-4 α and H-4 β , each d, $J = 14.6$ Hz), two pairs of oxymethylene protons at δ_H 4.59 and 4.72 (H-10 α and H-10 β , each d, $J = 13.2$ Hz) and δ_H 4.26 and 4.58 (H-1 α and H-1 β), and one methine proton at δ_H 2.99 (H-9, dd, $J = 6.5, 4.9$ Hz) attributed to the boschnialactone type skeleton (Sakan et al., 1967; Sisido et al., 1968; Callant et al., 1983; Tanaka et al., 1993; Hilgraf et al., 2012). The signal observed at δ_H 2.09 (3H, H-2', s) indicated the presence of an acetyl group in this compound. This was supported by the ^{13}C -NMR spectrum exhibiting eleven carbon signals including two ester carbonyls at δ_C 174.4 (C-3) and 172.4 (C-1'), an acetoxymethyl carbon at δ_C 20.7 (C-2'), two ethylenic carbons at δ_C 136.7 (C-8) and 131.1 (C-7), four methylene carbons at δ_C 68.4 (C-1), 62.5 (C-10), 48.9 (C-6) and 44.4 (C-4), one methine carbon at δ_C 55.4 (C-9) and one quaternary carbon at δ_C 78.9 (C-5) bearing an hydroxy group. These NMR data are very closed to owariensisone except for a supplementary acetyl group in C-10 position (Foning Tebou et al., 2016). The $^1J_{C-H}$ correlation in the HSQC spectrum allowed us to attribute to each carbon the corresponding proton. Thus, the two methylene protons at δ_H 2.60 and 2.65 (H-6) were linked to carbon C-6 and those at δ_H 2.80 and 2.83 (H-4) were correlated to carbone C-4, and the two pairs oxymethylene protons group at δ_H 4.26 and 4.58 (H-1) and 4.59 and 4.72 (H-10) were linked to carbon C-1 and C-10, respectively. In the COSY spectrum, protons at δ_H 2.60 and 2.65 (H-6) were correlated with proton at δ_H 5.83 (H-7). Mains correlations were also observed between the protons at δ_H 4.26 and 4.58 (H-1) and proton at δ_H 2.99 (H-9).

The two *gem*-methylene systems H-1 and H-4 were allocated to the δ -lactone group by the HMBC cross peaks with the carbonyl carbon at δ_C 174.4 thus confirming its C-3 position. From the HMBC spectrum, correlations observed between the methylene protons H-10 and carbons C-7, C-8 and C-9 suggested that this oxymethylene was located at C-8. The carbonyl of the methyl ester at 172.4 (C-1') was assigned to C-10 by long-range correlation between protons at δ_H 2.09 (H₃-2') and carbon at δ_C 62.5 (C-10).

The β -orientation of hydrogen at C-9 and hydroxyl group at C-5 was supported by the biosynthetic pathway to iridomyrmercin, isoiridomyrmercin and owariensisone, respectively (Lunn, 1961; Foning Tebou et al., 2016), as in usual iridoids. This is confirmed by the coupling constants of H-9 from 6.5 and 4.9 Hz with both protons H-1, characteristic of *pseudo*-axial-axial and *pseudo*-axial-equatorial system as in owariensisone. On the basis of afore mentioned information, the structure of compound **1** was elucidated as 10-*O*-acetyl

owariensisone or lactone of α -(3-acetoxymethyl-2-hydroxymethylcyclopent-3-en-yl) ethanoic acid named owariensisone **B** (Fig. 1).

Compound **2** was obtained as yellowish gum. Its molecular formula was determined as $C_{13}H_{18}O_8$ on the basis of its pseudo-molecular ion peak at m/z 325.0905 $[M+Na]^+$ (calcd. for $C_{13}H_{18}O_8Na$ 325.0899) in the HR-TOFESIMS spectrum, with five degrees of unsaturation. The 1H -NMR spectrum (Table 1) shows signals for two acetyl groups [δ_H 2.09 (3H, H-2', s), 2.07 (3H, H-2'', s); δ_C 172.5 (C-1'), 171.8 (C-1''), 20.7 (C-2'), 21.0 (C-2'')] and signals of iridoid lactone: two pairs of methylene protons at δ_H 2.00 and 2.43 (H-6 α , dd, J = 14.9, 1.8 Hz and H-6 β , dd, J = 14.9, 5.0 Hz) and δ_H 2.73 and 2.85 (H-4 α , dd, J = 14.8, 0.8 Hz and H-4 β , d, J = 14.8 Hz), two pairs of oxymethylene protons at δ_H 4.49 and 4.53 (H-1 α , dd, J = 11.9, 7.7 Hz and H-1 β , dd, J = 11.9, 10.0 Hz) and δ_H 4.12 and 4.38 (H-10 α , d, J = 11.6 Hz and H-10 β , d, J = 11.6 Hz), one methine proton at δ_H 2.57 (H-9, dd, J = 10.0, 7.7 Hz) and one oxymethine proton at δ_H 5.07 (H-7, dd, J = 5.0, 1.8 Hz). This was supported by the ^{13}C -NMR spectrum exhibiting signals of an ester carbonyl at δ_C 174.6 (C-3), four methylene carbons at δ_C 66.6 (C-1), 67.4 (C-10), 45.3 (C-6) and 44.9 (C-4), two methine carbons at δ_C 80.9 (C-7) and 51.9 (C-9) and two quaternary carbons at δ_C 82.1 (C-8) and 80.1 (C-5) bearing each an hydroxy group (Table 1). Their attributions were assigned by analysis of COSY and $^1J_{C-H}$ HSQC spectra. In the COSY spectrum, proton at δ_H 5.07 (H-7) was correlated with protons at δ_H 2.00 and 2.43 (H-6). Another correlation was also observed between the protons at δ_H 4.49 and 4.53 (H-1) and proton at δ_H 2.57 (H-9). As compared to compound **1**, the ethylenic group in $\Delta_{7,8}$ was replaced by two oxycarbonates at C-7 and C-8 as in patriscabrol (Kouna et al., 1994). Analysis of HMBC spectrum showed correlations between the protons at δ_H 4.12 and 4.38 (H-10) and carbons at δ_C 82.1 (C-8), 80.9 (C-7) and 67.4 (C-9) suggesting that this oxymethylene was located at C-8 as in **1**. The carbonyl of the methyl ester at 172.5 (C-1') was assigned to C-10 by long-range correlation with protons H-10, while the one at 171.8 (C-1'') was linked to C-7 by the long-range correlation with proton at δ_H 5.07 (H-7). The position of the lactone at δ_C 174.6 was determined as in **1**. The β -orientation of H-9 was deduced from the coupling constant of 10.0 and 7.7 Hz with protons H-1 α -axial and H-1 β -equatorial, respectively. The α -equatorial orientation of H-7 was deduced from the coupling constant of 5.0 and 1.8 Hz with H-6 α -axial and H-6 β -equatorial, respectively as observed for similar protons in jioglutoside (Morota et al., 1989) and patriscabrol (Kouna et al., 1994). Thus, the hydroxyl at C-7 is β -oriented. As observed in jioglutoside the δ -lactone ring has a boat conformation (1B4) (Morota et al., 1989), as observed by the ROESY correlations between H-1 and H-4 α -axial. Correlations observed in the ROESY spectrum between H-9 and H-10 β , indicated a β -

orientation for C-10 and an α -orientation of hydroxyl at C-8 position (Fig. 2). On the basis of afore mentioned information, the structure of compound **2** was elucidated as lactone of α -(4- β -acetoxy-3- β -acetoxymethyl-2-hydroxymethyl-3-hydroxycyclopentyl) ethanoic acid named owariensisone **C** (Fig.1).

Compound **3** was obtained as yellowish gum. It has the same molecular formula $C_{13}H_{18}O_8$ as compound **2**, and differs from the latter only by the orientation of C-8 oxymethylene. Its 1H -NMR spectrum (Table 1) was so closed to that of **2** and its ^{13}C -NMR spectrum is almost superimposable on that of **2**, except for the signal of oxymethylene C-10 δ_C 65.0 (Δ -2.4 ppm) and methine C-9 δ_C 54.0 (Δ +2.1 ppm). From the analysis of HMBC spectrum, the oxymethylene, the carbonyl of the methyl ester and the position of the lactone have been solved as in **2**. The absence of correlation between H-10 and H-9 in the ROESY spectrum indicated the α orientation of C-10 (Fig.2). On the basis of afore mentioned information, the structure of compound **3** was elucidated as lactone of α -(4- β -acetoxy-3- α -acetoxymethyl-2-hydroxymethyl-3-hydroxycyclopentyl) ethanoic acid named owariensisone **D** (Fig.1).




Fig.1. Structures of compounds **1-3** isolated from the *n*-hexane soluble extract of *Brillantaisia owariensis*.




Fig.2. Noesy effects in compounds **2-3**

Table 1: ^1H -NMR (600 MHz) and ^{13}C -NMR (150 MHz) data of compounds **1-3** in CD_3OD .

N°	1		2		3	
	^{13}C	^1H (J in Hz)	^{13}C	^1H (J in Hz)	^{13}C	^1H (J in Hz)
1	68.4	4.26 (1H, dd, 11.9, 6.5) 4.58 (1H, dd, 11.9, 4.9)	66.6	4.49 (1H, dd, 11.9, 7.7) 4.53 (1H, dd, 11.9, 10.0)	66.9	4.21 (1H, t, 11.2) 4.57 (1H, dd, 11.6, 6.2)
3	174.4		174.6		174.5	
4	44.4	2.80 (1H, d, 14.6) 2.83 (1H, d, 14.6)	44.9	2.73 (1H, dd, 14.8, 0.8) 2.85 (1H, d, 14.8)	44.7	2.57 (1H, d, 15.1) 2.78 (1H, d, 15.1)
5	78.9		80.1		80.4	
6	48.9	2.60 (1H, dq, 18.1, 1.0) 2.65 (1H, dm, 18.1)	45.3	2.00 (1H, dd, 14.9, 1.8) 2.43 (1H, dd, 14.9, 5.0)	46.1	1.91 (1H, dd, 14.5, 1.6) 2.46 (1H, dd, 14.5, 5.0)
7	131.1	5.83 (1H, brqt, 1.6)	80.9	5.07 (1H, dd, 5.0, 1.8)	80.8	5.17 (1H, dt, 5.0, 1.6)
8	136.7		82.1		82.1	
9	55.4	2.99 (1H, dd, 6.5, 4.9)	51.9	2.57 (1H, dd, 10.0, 7.7)	54.0	2.45 (1H, dd, 11.1, 6.2)
10	62.5	4.59 (1H, d, 13.2) 4.72 (1H, d, 13.2)	67.4	4.12 (1H, d, 11.6) 4.38 (1H, d, 11.6)	65.0	3.91 (1H, d, 11.6) 4.18 (1H, d, 11.6)
11	172.4		172.5		172.4	
12	20.7	2.09 (3H, s)	20.7	2.09 (3H, s)	20.7	2.09 (3H, s)
13			171.8		171.2	
14			21.0	2.07 (3H, s)	21.0	2.07 (3H, s)

3. Experimental

3.1. General

IR spectra were recorded with a Shimadzu FT-IR-8400S (Shimadzu, France) spectrophotometer. ^1H and ^{13}C -NMR spectra were recorded on a Bruker Avance III 600 spectrometer equipped with a cryoprobe (^1H at 600 MHz and ^{13}C at 150 MHz). 2D NMR experiments were recorded by means of standard Bruker microprograms (Xwin-NMR version 2.1 software TopSpin 3.2). Chemical shifts (δ) are reported in parts per million (ppm) using the residual solvent signals as secondary reference relatively to TMS ($\delta = 0$), while the coupling constants (J values) are given in Hertz (Hz). TOF-ESIMS and HR-TOFESIMS spectra were recorded using a Micromass Q-TOF micro instrument (Manchester, UK) equipped with an electrospray source. The samples were introduced by direct infusion in a solution of MeOH at a rate of $5 \mu\text{L min}^{-1}$. The optical rotations were measured on a Bellingham & Stanley ADP 220 polarimeter (Bellingham + Stanley Ltd., United Kingdom). Column chromatography was run on Merck silica gel 60 (70-230 mesh) and gel permeation on Sephadex LH-20 while TLC was carried out on silica gel GF₂₅₄ pre-coated plates with detection accomplished by spraying with 50% H_2SO_4 followed by heating at 100°C , or by visual inspection under UV lamp at 254 and 365 nm.

3.1. Plant material

The whole plant of *B. owariensis* was collected at Tsinkop, Menoua Division, West Region of Cameroon, in October 2013. Authentication was done by Mr Victor Nana, a botanist of the

Cameroon National Herbarium, Yaoundé, where the voucher specimen (N° 34376/HNC) has been deposited.

3.2.Extraction and isolation

The air-dried and plant material (4 Kg) was powdered and extracted at room temperature with methanol (3 x 15L, 72h). The solvent was evaporated under reduced pressure, leaving an extract (110 g). Part of this extract (100 g) was extracted with *n*-hexane yielding **30** g fraction after evaporation to dryness. Part of *n*-hexane-soluble fraction (**25** g) was subjected to silica gel column chromatography using Hex-EtOAc (9:1 to 1:0) gradient graduated elution. Many fractions of 250 mL each were collected and combined on the basis of their TLC profiles to give 3 fractions: A, B and C. Fraction B (5.1 g) was purified on silica gel column chromatography using an isochratic eluent system Hex-EtOAc (1:1). Fractions of 10 mL were collected and combined on the basis of TLC profiles to give 3 sub-fractions B₁, B₂ and B₃. Sub-fraction B₂ (102 mg) was subjected to silica gel column chromatography using Hex-EtOAc (3:2) as eluent to give compound **1** (8 mg) and a mixture of **2** and **3**. This mixture was combined to sub-fractions B₁ and B₃ and purified by chromatography on silica gel column chromatography to yield compounds **3** (12 mg) and **2** (9mg).

New compounds

Owariensisone B: Yellowish gum; ¹H and ¹³C-NMR data, see Table 1; $[\alpha]_D^{21} -5.2$ (MeOH *c* 0.06); IR (NaCl) ν_{\max} (cm⁻¹) 3350-3300, 1070, 1040 (OH), 1643 (C=O), 1605, 1580, 1520 (C=C aromatic), 1650 (C-O); HR-TOFESIMS (positive ion mode) *m/z*: 249.0734 [M+Na]⁺ (calcd. for C₁₁H₁₄O₅Na 249.0739).

Owariensisone C: Yellowish gum; ¹H and ¹³C-NMR data, see Table 1; $[\alpha]_D^{21} -20$ (MeOH *c* 0.05); IR (NaCl) ν_{\max} (cm⁻¹) 3350-3300, 1070, 1040 (OH), 1643 (C=O), 1605, 1580, 1520 (C=C aromatic), 1650 (C-O); HR-TOFESIMS (positive ion mode) *m/z*: 325.0905 [M+Na]⁺ (calcd. for C₁₃H₁₈O₈Na 325.0899).

Owariensisone D: Yellowish gum; ¹H and ¹³C-NMR data, see Table 1; $[\alpha]_D^{21} -87.5$ (MeOH *c* 0.02); IR (NaCl) ν_{\max} (cm⁻¹) 3350-3300, 1070, 1040 (OH), 1643 (C=O), 1605, 1580, 1520 (C=C aromatic), 1650 (C-O); HR-TOFESIMS (positive ion mode) *m/z*: 325.0892 [M+Na]⁺ (calcd. for C₁₃H₁₈O₈Na 325.0899).

Acknowledgments

The authors are grateful to the University of Dschang for financing some consumables used in this work, to the “Service Commun d’Analyses” and “Groupe Isolement et Structure”, to the “Institut de Chimie Moléculaire de Reims” for the spectroscopic and spectrometric analysis

on the ESIMS et NMR equipment of the PIANet Platform. The EU-programme FEDER to the PIANet CPER project is gratefully acknowledged.

References

- Aluko**, B.T., Alli Smith, Y. R., Omoyeni O. A. 2014. Phytochemical analysis and antioxidant activities of ethanolic leaf extract of *Brillantaisia patula*. *World Journal of Pharmaceutical Research* **3**, 4914-4924.
- Asai**, T., Hirayama, Y., Fujimoto, Y., 2012. Epi- α -bisabolol 6-deoxy- β -D-gulopyranoside from the glandular trichome exudate of *Brillantaisia owariensis*. *Phytochemistry Letters* **5**, 376-378.
- Callant**, P., Storme, P., Van der Eycken, E., Vandewalle, M., 1983. Iridoïds: stereospecific synthesis of functionalized cyclopentanoid intermediates *via* bicycle [2.2.1] heptanones. *Tetrahedron Letters* **24**, 5797-5800.
- Faparusi**, F., Bello-Akinosho, M. M., Oyede, R. T., Adewole, A., Bankole, P. O., Ali, F. F., 2012. Phytochemical Screening and Antibacterial Activity of *Brillantaisia patula* Leaf. *Research Journal of Phytochemistry* **6**, 9-16.
- Foning** Tebou, P. L., Mabou, F. D., Ngnokam, D., Harakat, D., Voutquenne-Nazabadioko, L., 2016. Owariensisine: a new iridolactone from the whole plant of *Brillantaisia owariensis* P. Beauv (Acanthaceae). *Natural Product Research* **134**, 1478-1485.
- Heine**, H., 1963. *Flora of West tropical Africa*, Vol. 2. *Acanthaceae*, 2nd edn (ed. By J. Hutchinson and J. M. Dalziel), pp. 391-425. Hepper, F. N. Millbank, London.
- Hilgraf**, R., Zimmermann, M., Lehmann, L., Tröger, A., Francke, W., 2012. Stereoselective synthesis of *trans*-fused iridoïd lactones and their identification in the parasitoid wasp *Alloxy stavictrix*, Part II: Iridomyrmecins. *Beilstein Journal of Organic Chemistry* **8**, 1256-1264.
- Kouno**, I., Yasuda, I., Mizoshiri, H., Tanaka, T., Marubayashi, N., Yang, D. M., 1994. Two new iridolactones and their glycosides from the roots of *Patrinia scabra*. *Phytochemistry* **37**, 467-472.
- Lunn**, W. H., 1961. Part I: The structure of catalposide. Part II: Carbonium ion formation at the C-10 position in 2,5-*seco*-A-bisnorcholestane structures. A thesis submitted to the Faculty of graduate studies and research in partial fulfillment of the requirements for the degree of Doctor of Philosophy. McGill University, Montreal P.Q. Canada. PP. 143.
- Makambila-Koubemba**, M. C., Mbatchi, B., Ardid, D., Gelot, A., Henrion, C., Janisson, R., Abena, A. A., Banzouzi, J. T., 2011. Pharmacological studies of ten medicinal plants used for analgesic purposes in Congo Brazzaville. *International Journal of Pharmacology* **7**, 608-615.

- Mbatchi**, S. F., Mbatchi, B., Banzouzi, J. T., Bansimba, T., Nsonde-Ntandou, G. F., Ouamba, J. M., Berry, A., Benoit-Vical, F., 2006. *In vitro* antiplasmodial activity of 18 plants used in Congo Brazzaville traditional medicine. *Journal of Ethnopharmacology* **104**, 168-174.
- Ngbolua**, K. N., Benamambote, B. M., Muanda, D. M., Ekutsu, E., Tshibangu, D. S. T., Gbolo, B. Z., Muanyishay, C. L., Basosila, N. B., Bongo, G. N., Robijaona, B., 2013. Ethnobotanical survey and Ecological Study of some Medicinal Plants species traditionally used in the District of Bas-Fleuve (Bas-Congo Province, Democratic Republic of Congo). *Research Journal of Chemistry* **1**, 1-10.
- Morota**, T., Nishimura, H., Sasaki, H., Chin, M., Sugama, K., Katsuhara, T., Mitsunashi, H., 1989. Five cyclopentanoid monoterpenes from *Rehmannia glutinosa*. *Phytochemistry* **28**, 2385-2391.
- Sakan**, T., Murai, F., Hayashi, Y., Honda, Y., Shono, T., Nakajima, M., Kato, M., 1967. Structure and stereochemistry of boschniakine, boschnialactone and boschnialinic acid, an oxidation product of boschnialactone. *Tetrahedron* **3**, 4635-4652.
- Sisido**, K., Inomata, K., Kageyema, T., Utimoto, K., 1968. Conformations of iridolactones and the stereochemistry in the syntheses. *Journal of Organic Chemistry* **33**, 3149-3155.
- Tanaka**, D., Yoshino, T., Kouno, I., Miyashita, M., Irie, H., 1993. Synthesis of monoterpene lactones, (+)-boschnialactone and (+)-isoiridomyrmecin, starting from L-(+)-arabinose. *Tetrahedron* **49**, 10253-10262.