



HAL
open science

A new cardenolide and other compounds from *Salsola tetragona*

Hamida Ghoraba, Abdeikrim Khettaf, Merien Lehbili, Ahmed Kabouche,
Abdulmagid Alabdul Magid, Dominique Harakat, Laurence
Voutquenne-Nazabadioko, Zahia Kabouche

► **To cite this version:**

Hamida Ghoraba, Abdeikrim Khettaf, Merien Lehbili, Ahmed Kabouche, Abdulmagid Alabdul Magid, et al.. A new cardenolide and other compounds from *Salsola tetragona*. *Natural Product Communications*, SAGE Publications, 2017, 12 (1), pp.3-5. 10.1177/1934578X1701200102. hal-01996632

HAL Id: hal-01996632

<https://hal.univ-reims.fr/hal-01996632>

Submitted on 28 Oct 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

A New Cardenolide and Other Compounds from *Salsola tetragona*Hamida Ghorab^a, Abdelkrim Khettaf^a, Meriem Lehbili^a, Ahmed Kabouche^a, Abdulmagid Alabdul Magid^b, Dominique Harakat^c, Laurence Voutquenne-Nazabadioko^{b*} and Zahia Kabouche^{a*}.^aUniversité des frères Mentouri-Constantine, Laboratoire d'Obtention de Substances Thérapeutiques (LOST), Campus Chaabet Ersas, 25000 Constantine, Algeria^bGroupe Isolement et Structure, Institut de Chimie Moléculaire de Reims (ICMR), CNRS UMR 7312, UFR de Pharmacie, BP 1039, 51687 Reims, France^cService Commun d'Analyses, ICMR-UMR CNRS 7312, Bât. 18 B.P.1039, 51687 Reims Cedex 2, France

zahiakabouche@gmail.com

Received: August 3rd, 2016; Accepted: November 10th, 2016

One new cardenolide, 3-*O*-β-D-allopyranosylcoroglaucigenin (salsotetragonin) (**1**), was isolated from the aerial parts of *Salsola tetragona* Delile with four known cardenolides (**2-5**), two known flavonoids (**6-7**), three known phenolic compounds (**8-10**) and two known fatty acids (**11-12**). Their structures were identified by spectroscopic analyses and by comparison of their spectral data with those reported in the literature. Compounds **1-5**, **7**, **10** and **12** were isolated from the genus *Salsola* for the first time. This is the first report on cardenolides identified in the Amaranthaceae family.

Keywords: *Salsola tetragona* Delile, Amaranthaceae, Cardenolides, Phenolic glycosides.

The genus *Salsola* (Amaranthaceae, ex. Chenopodiaceae) consists of over 174 species found in the arid regions of Asia, Europe and Africa [1]. *Salsola* species possess antihypertensive, anti-inflammatory, anticancer, antioxidant, anti-Alzheimer, antidepressant and antimicrobial activities [2-5]. Previous phytochemical investigations of this genus reported the isolation of flavonoids and other phenolic compounds [6], alkaloids [7], fatty acids [8] and triterpene glycosides [9]. In continuation of our works on plants from Algerian Septentrional Sahara [10-11], we report here the isolation and structural characterization of one new cardenolide, 3-*O*-β-D-allopyranosylcoroglaucigenin, named salsotetragonin (**1**), in addition to eleven known compounds from aerial parts of *S. tetragona* (Figure S1).

The hydromethanolic extract of the aerial parts of *S. tetragona* was partitioned successively with light petroleum, CH₂Cl₂, EtOAc and *n*-butanol to give four extracts. Purification of the *n*-butanol extract through repeated silica gel columns, flash chromatography and semi-prep HPLC led to the isolation and identification of a new cardenolide (**1**) and seven known compounds (**3**, **4**, **6-10**). The dichloromethane extract was fractionated by silica gel chromatography to give four known compounds (**2**, **5**, **11-12**). These compounds were identified by extensive spectroscopic methods including 1D-(¹H and ¹³C) and 2D-NMR (COSY, HSQC, HMBC and NOESY) experiments, as well as HR-ESI-MS analysis and by comparison of their spectral data with the literature. The eleven known compounds were identified as four cardenolides: uzarigenin (**2**) [12], desglucouzarin (**3**) [13], 12-dehydroxyghalakinoside (**4**), and calactin (**5**) [14] (Figure 1), two flavonoids: kaempferol-3-*O*-β-D-glucopyranoside (**6**) [15] and quercetin-3-*O*-β-D-glucopyranosyl-(1→6)-glucopyranoside (**7**) [16], three phenolic compounds: vanillic acid (**8**) [17], canthoside C (**9**), and canthoside D (**10**) [18] and two fatty acids: oleic acid (**11**) [19], and 2,3-dihydroxypropylpalmitate (**12**) [20] (Figure S1).

Compound **1** was isolated as a white amorphous powder. Its positive HR-ESI-MS showed a pseudomolecular ion peak at *m/z* 575.2829 [M+Na]⁺, corresponding to the molecular formula C₂₉H₄₄O₁₀. The




Figure 1: Structures of compounds 1-5.

¹H NMR spectrum of compound **1** (Table 1) showed characteristic signals of a butenolactone ring at δ_H 4.95 (1H, dd, *J* = 18.4, 1.6 Hz, H-21a), 5.07 (1H, dd, *J* = 18.4, 1.4 Hz, H-21b) and 5.93 (1H, s, H-22), one oxymethylene at δ_H 3.74 (1H, d, *J* = 11.7 Hz, H-19a) and δ_H 3.88 (1H, d, *J* = 11.7 Hz, H-19b), one oxymethine at δ_H 3.78 (m, H-3), and one methyl signal at δ_H 0.95 (3H, s, H-18). Its ¹³C NMR spectrum (Table 1) showed twenty-nine carbon signals including six of the sugar part (Figure 1).

The ¹H NMR and ¹³C NMR data of **1** were very similar to those of 3-β-D-glucopyranosylcoroglaucigenin [21]. The COSY and HSQC spectra showed one anomeric proton of a sugar moiety at δ_H 4.78 (d, *J* = 7.9 Hz, H-1') and δ_C 99.8 with five signals of an hexose unit [H-2' [δ 3.28 (dd, *J* = 7.9, 3.0 Hz)], H-3' [δ 4.06 (t, *J* = 3.0 Hz)], H-4' [δ 3.48 (dd, *J* = 9.5, 3.0 Hz)], H-5' [δ 3.69 (m)], and H₂-6' [δ 3.66 (dd, *J* = 11.3, 5.7 Hz) and 3.85 (dd, *J* = 11.3, 1.8 Hz)]; the sugar was identified as an allopyranoside [22]. The large coupling constant (*J* = 7.9 Hz) of the anomeric proton at δ_H 4.78 indicated that the allose had a β-configuration. The linkage of the allose at C-3 of the aglycone was established from the HMBC correlation between allo-H-1' (δ_H 4.78) and C-3 (δ_C 78.9) (Figure 2). The stereochemistry of **1** was confirmed by the NOE effect (Figure 2) between H-5/H-9 and H-3, which clearly established the α-orientation of H-5. The correlation of H₃-18 to H-21 and H-22 in the

Noesy spectrum indicated the β orientation of the γ lactone at C-17. The chemical shifts of the carbons of the C and D rings of **1** are in agreement with those of coroglaucigenin and madagascarrenilid (A) recorded in deuterated pyridine and in CD_3OD [23,24], indicating a 14- β hydroxy orientation. Thus compound **1** was identified as 3-O- β -D-allopyranosylcoroglaucigenin, which we named salsotetragonin.




Figure 2: ^1H - ^1H COSY, HMBC and NOESY correlations of **1**.

Table 1: ^1H and ^{13}C NMR data for compound **1** (in CD_3OD ; ^{13}C : 150 MHz; ^1H : 600 MHz).

Position	1	
	δ_{H} (m, J in Hz)	δ_{C}
1	0.81 (td, 13.6, 3.5), 2.34 (dt, 13.3, 3.5)	32.8
2	1.57 (dm, 14.9), 1.91 (m)	30.9
3	3.78 (m)	78.9
4	1.44 (m), 1.82 (dd, 11.9, 3.4)	35.9
5	1.24 (m)	45.9
6	1.24 (m), 1.34 (m)	29.6
7	2.10 (dd, 12.3, 3.1), 1.15 (m)	28.8
8	1.82 (td, 11.9, 3.4)	43.1
9	1.03 (td, 13.5, 4.5)	51.5
10	-	40.6
11	1.65 (m)	24.1
12	1.41 (td, 13.6, 4.3), 1.53 (dt, 13.5, 3.0)	41.5
13	-	51.2
14	-	86.3
15	1.73 (dd, 11.6, 8.5), 2.14 (m)	33.5
16	2.16 (m), 1.90 (m)	28.1
17	2.85 (dd, 9.5, 5.8)	52.2
18	0.95 (s)	16.7
19	3.74 (d, 11.7), 3.88 (d, 11.7)	59.9
20	-	178.5
21	4.95 (dd, 18.4, 1.6), 5.07 (dd, 18.4, 1.4)	75.3
22	5.93 (s)	117.8
23	-	176.9
Allose (at C-3)		
1'	4.78 (d, 7.9)	99.8
2'	3.28 (dd, 7.9, 3.0)	72.4
3'	4.06 (t, 3.0)	73.1
4'	3.48 (dd, 9.5, 3.0)	69.1
5'	3.69 (m)	75.5
6'	3.66 (dd, 11.3, 5.7), 3.85 (dd, 11.3, 1.8)	63.3

Experimental

General: Optical rotations were measured in DMSO with a Perkin-Elmer 241 polarimeter. ^1H and ^{13}C NMR spectra were recorded on a Bruker Avance DRX III 500 spectrometer (^1H at 500 MHz and ^{13}C at 125 MHz) and on a Bruker Avance III-600 spectrometer equipped with a cryo platform (^1H at 600 MHz and ^{13}C at 150 MHz). 2D-NMR experiments were performed using standard Bruker-microprograms (TopSpin version 3.2 software). HR-ESI-

MS and ESI-MS experiments were performed using a Micromass Q-TOF micro-instrument (Manchester, UK). TLC was carried out on pre-coated silica gel 60 F₂₅₄ (Merck). Vacuum liquid chromatography (VLC) was carried out on LiChroprep RP-18 (40-63 μm , Merck). HPFC was performed on a Grace® Reveleris System using Grace® cartridges (Silica gel or RP-C₁₈). The semi-preparative HPLC was performed on a Dionex apparatus equipped with an ASI-100 automated sample injector, a STH 585 column oven, a LPG-3400A pump, a diode array detector UVD-340U and Chromeleon® software.

Plant material: *Salsola tetragona* Delile was collected in Ghardaïa (Algerian Septentrional Sahara) in May 2011 and was identified by Pr. Gérard De Belair (University of Annaba, Algeria). A voucher specimen (LOST.St05/09) was deposited at the herbarium of the Faculty of Sciences, University frères Mentouri- Constantine.

Extraction and isolation: Air-dried and powdered aerial parts (1980 g) of *S. tetragona* were extracted with 80% MeOH (10 L). After evaporating the MeOH under *vacuum*, the residue was dissolved in water (900 mL) and partitioned with PE, CH_2Cl_2 , EtOAc, and *n*-BuOH, successively (3 x 300 mL). The *n*-BuOH extract (6 g) was subjected to VLC over RP-18 eluted with H_2O -MeOH (6:4, 5:5, 4:6, 3:7, 2:8 and 0:10). Fractions of 150 mL were collected and pooled according to their similarity in TLC profile to give 6 fractions (F1-F6 respectively). F1 (1.2 g) was submitted to RP-18 flash chromatography eluting with MeCN:H₂O (5:95 to 10:0) to afford 19 sub-fractions. Sub-fraction [20-23] (35 mg), was purified by semi-prep HPLC on RP-18 eluted with MeCN:H₂O (6:94 to 1:9), affording compounds **9** (3 mg) and **10** (1.5 mg). F2 (1.5 g) was chromatographed on a silica gel column eluted with EtOAc:MeOH:H₂O (20:2:1), giving 2 main sub-fractions. Sub-fraction [90] (180 mg) afforded **4** (22 mg) by silica gel CC (CHCl_3 :MeOH, 9:1). Sub-fraction [115-119] (223 mg) was subjected to silica gel CC eluting with CHCl_3 :MeOH (8:2) to afford **1** (27 mg) and sub-fraction [35-48] (100 mg), which was purified over silica gel using EtOAc:MeOH:H₂O (20:2:1) to yield compounds **6** (15 mg) and **7** (18 mg). F3 (750 mg) was fractionated by silica gel CC eluted with CHCl_3 :MeOH (0 to 100% of MeOH). The main sub-fraction [59-63] (212 mg) was further separated by silica gel CC (CHCl_3 :MeOH, 0 to 100% of MeOH) and **3** (25 mg) was isolated. F4 (302 mg) was subjected to silica gel CC using CHCl_3 :MeOH (0 to 100% of MeOH) to give 3 sub-fractions. Sub-fraction [72] (111.2 mg) was selected for silica gel CC eluted with CHCl_3 :MeOH (0 to 100% of MeOH) to yield compound **8** (3 mg).

The CH_2Cl_2 extract (2.5 g) was subjected to silica gel CC, eluted with toluene: CHCl_3 (0 to 100% of CHCl_3) and CHCl_3 :EtOAc (0 to 100% of EtOAc). Sub-fraction [27-32] (123 mg) was purified over silica gel using diethyl ether:EtOAc (9:1) to yield compound **11** (1.4 mg). Sub-fraction [160-173] (212 mg) was passed through a silica gel column (CHCl_3 :EtOAc, 9:1) yielding compounds **2** (15 mg) and **5** (5 mg). Sub-fraction [234-257] (162 mg) was chromatographed on silica gel, eluted with (CHCl_3 :MeOH, 9:1) that afforded compound **12** (2 mg).

Acid hydrolysis: The pure compounds were treated with 2 M HCl at 100°C for 1 h. The hydrolysates were extracted with EtOAc and the sugars identified in the aqueous residue by comparison with authentic samples by TLC using silica gel impregnated with 0.2 M NaH_2PO_4 and a solvent system of Me_2CO -H₂O (9:1); the compounds were revealed with aniline malonate. The optical rotation of the purified sugar was measured and compared with an authentic sample of D-allose.

3-O-β-D-Allopyranosylcoroglaucigenin (salsotetragonin) (1)

White amorphous powder.

 $[\alpha]_D^{20}$: -16.9 (c 0.18, DMSO). ^1H NMR (CD₃OD, 600 MHz): Table 1. ^{13}C NMR (CD₃OD, 150 MHz): Table 1.HR-ESI-MS: $[\text{M}+\text{Na}]^+$ m/z 575.2829 (calcd for C₂₉H₄₄O₁₀Na, 575.2832).

Acknowledgments - The authors would like to thank the DGRSDT-MESRS (Algeria) for financial support and the Groupe Isolement et Structure of the Institut de Chimie Moléculaire de Reims (France) for technical help and registering of NMR data and MS.

Supplementary data: Figure S1 and Supplementary Data are included in the Supporting information.

References

- [1] Botschantzev V. (1974) A synopsis of *Salsola* (Chenopodiaceae) from South and South-West Africa. *Kew Bulletin*, **29**, 597-614.
- [2] Ahmad Z, Mehmood S, Fatima I, Malik A, Ifzal R, Afza N, Iqbal L, Latif M, Nizami TA. (2008) Structural determination of salsolins A and B, new antioxidant polyoxygenated triterpenes from *Salsola baryosma* by 1D and 2D NMR spectroscopy. *Magnetic Resonance in Chemistry*, **46**, 94-98.
- [3] Woo WS, Lee EB, Chi HJ. (1977) Biological evaluation of some Saudi Arabian plants. *Journal of Korean Pharmaceutical Sciences*, **21**, 141-145.
- [4] Oueslati MH, Ben JH, Mighri Z, Chriaa J, Abreu PM. (2006) Phytochemical constituents from *Salsola tetrandra*. *Journal of Natural Products*, **69**, 1366-1369.
- [5] Tundis R, Loizzo MR, Statti GA, Menichini F. (2007) Inhibitory effects on the digestive enzyme alpha-amylase of three *Salsola* species (Chenopodiaceae) *in vitro*. *Pharmazie*, **62**, 473-475.
- [6] Xiang Y, Li YB, Zhang J, Li P, Yao YZ. (2007) Studies on chemical constituents of *Salsola collina*. *Zhongguo Zhong Yao Za Zhi*, **32**, 409-413.
- [7] Xiang Y, Li YB, Zhang J, Li P, Yao YZ. (2007) A new alkaloid from *Salsola collina*. *Yao Xue Xue Bao*, **42**, 618-620.
- [8] Jin YS, Du JL, Yang Y, Jin L, Song Y, Zhang W, Chen HS. (2011) Chemical and biologically active constituents of *Salsola collina*. *Chemistry of Natural Compounds*, **47**, 257-260.
- [9] Hamed AI, Masullo M, Sheded MG, Mahalel UA, Tawfik MM, Perrone A, Piacente S. (2011) Triterpene saponins from *Salsola imbricata*. *Phytochemistry Letters*, **4**, 353-356.
- [10] Boudiar T, Lakhall H, Khalfallah A, Kabouche A, Kabouche Z, Brouard I, Bermejo J, Christian B. (2010) A new alkaloid and flavonoids from the aerial parts of *Euphorbia guyaniana*. *Natural Product Communications*, **5**, 35-37.
- [11] Lakhall H, Boudiar T, Kabouche A, Kabouche Z, Touzani R, Christian B. (2010) New sesquiterpene lactone and other constituents from *Centaurea sulphurea* (Asteraceae). *Natural Product Communications*, **5**, 849-850.
- [12] Shaker KH, Morsy N, Zinecker H, Imhoff JF, Schneider B. (2010) Secondary metabolites from *Calotropis procera* (Aiton). *Phytochemistry Letters*, **3**, 212-216.
- [13] Elgamel M, Hani A, Hanna AG, Morsy NAM, Duddeck H, Simon A, Gati T, Toth G. (1999) Complete ^1H and ^{13}C signal assignments of 5α cardenolides isolated from *Calotropis procera* R. B.R. *Journal of Molecular Structure*, **477**, 201-208.
- [14] Hamed AI, Plaza A, Balestrieri ML, Mahalel UA, Springuel IV, Oleszek W, Pizza C, Piacente S. (2006) Cardenolide glycosides from *Pergularia tomentosa* and their proapoptotic activity in Kaposi's sarcoma cells. *Journal of Natural Products*, **69**, 1319-1322.
- [15] Berrehal D, Khalfallah A, Kabouche A, Kabouche Z, Karioti A, Bilia AR. (2010) Flavonoid glycosides from *Randonia africana* Coss. (Resedaceae). *Biochemical Systematics and Ecology*, **38**, 1007-1009.
- [16] Shehab NG, Mahdy A, Khan SA, Noureddin SM. (2011) Chemical constituents and biological activities of *Fagonia indica* Burm F. *Research Journal of Medicinal Plant*, **5**, 531-546.
- [17] Wang XJ, Zhao YX, Jia XH, Ding XB. (2011) Studies on the chemical constituents of *Salsola collina*. *Zhong yao cai*, **34**, 230-231.
- [18] Kanchanapoom T, Kasai R, Yamasaki K. (2002) Iridoid and phenolic diglycosides from *Canthium berberidifolium*. *Phytochemistry*, **61**, 461-464.
- [19] Turki ZA. (2000) Chemotaxonomical studies of the genus *Salsola* (Chenopodiaceae) in Egypt. *Egyptian Journal of Botany*, **38**, 47-61.
- [20] Li Y, An K, Sun J, Wang J, Zhong Y, Sun L. (2011) Chemical constituents of vinegar-processed *Kansui radix*. *Food and Drug*, **13**, 183-186.
- [21] Jolad SD, Bastest RB, Cole JR, Hoffmann JJ, Siahaan TJ, Timmerann BN. (1986) Cardenolides and a lignan from *Asclepias subulata*. *Phytochemistry*, **25**, 2581-2590.
- [22] Murakami T, Emoto A, Matsuda H, Yoshikawa M. (2001) Medicinal foodstuffs. XXI. Structures of new cucurbitane-type triterpene glycosides, goyaglycosides-a, -b, -c, -d, -e, -f, -g, and -h, and new oleanane-type triterpene saponins, goyasaponins I, II, and III, from the fresh fruit of Japanese *Momordica charantia* L. *Chemical and Pharmaceutical Bulletin*, **49**, 54-63.
- [23] Yamauchi T, Abe F, Nish M. (1978) ^{13}C NMR of 5α-cardenolides. *Chemical and Pharmaceutical Bulletin*, **26**, 2894-2896.
- [24] Pan E, Harinantenaina L, Brodie PJ, Callmander M, Rakotonandrasana S, Rakotobe E, Resamison VE, TenDyke K, Shen Y, Suh EM, Kingston DGI. (2011) Cardenolides of *Leptadenia madagascariensis* from the Madagascar dry forest. *Bioorganic and Medicinal Chemistry*, **19**, 422-428.