



**HAL**  
open science

## Triterpenoid Saponins from the Fruits of *Caryocar glabrum*

Abdulmagid Alabdul Magid, Laurence Voutquenne, Christian Moretti,  
Christophe Long, Catherine Lavaud

► **To cite this version:**

Abdulmagid Alabdul Magid, Laurence Voutquenne, Christian Moretti, Christophe Long, Catherine Lavaud. Triterpenoid Saponins from the Fruits of *Caryocar glabrum*. *Journal of Natural Products*, 2006, 69 (2), pp.196-205. 10.1021/np050336s. hal-01996893

**HAL Id: hal-01996893**

**<https://hal.univ-reims.fr/hal-01996893>**

Submitted on 24 Nov 2021

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial 4.0 International License

## **Triterpenoid Saponins from the Fruits of *Caryocar glabrum***

Abdulmagid Alabdul Magid,<sup>†</sup> Laurence Voutquenne,<sup>\*,†</sup> Christian Moretti,<sup>‡</sup> Christophe Long,<sup>§</sup>  
and Catherine Lavaud<sup>†</sup>

*Laboratoire de Pharmacognosie, FRE CNRS 2715, IFR 53 Biomolécules, Bât. 18, BP 1039,  
51687 Reims Cedex 2, France, IRD, Unité N° S84 Biodival, Technoparc, 5 rue du Carbone,  
45072 Orléans Cedex 2, France, and Institut de Recherche Pierre Fabre - ISTMT - UMS  
CNRS 2579 - 3 rue des Satellites, BP 94244, 31432 Toulouse Cedex, France*

\* To whom correspondence should be addressed. Tel.: +33 (0)326918208. Fax:

+33(0)326913596. E-mail: laurence.voutquenne@univ-reims.fr.

<sup>†</sup> Université de Reims.

<sup>‡</sup> IRD.

<sup>§</sup> UMS CNRS 2579.

Twenty-one new triterpenoid saponins, named caryocarosides (**1–21**), glycosides of  $2\beta$ -hydroxyoleanolic acid, hederagenin, bayogenin and gypsogenic acid, have been isolated from the fruits of *Caryocar glabrum* along with nine known triterpenoid saponins (**22–30**) that are described for the first time from a plant in the Caryocaraceae. Their structures were established by 1D and 2D NMR techniques ( $^{13}\text{C}$ , COSY, TOCSY, HSQC, HMBC, and ROESY experiments), ESIMS, and acid hydrolysis. The isolated compounds could be classified into two series: glucosides (**1–8**, **22**, **27**, and **30**) derived from the 3-*O*-monoglucoside and glucuronides (**9–21**, **23–26**, **28**, and **29**) derived from the 3-*O*-monoglucuronide. In twenty-two of the saponins (**1–8**, **12–22**, and **24–26**), a galactose moiety was linked to C-3 of glucuronic acid or glucose moiety. The galactose was substituted in position 3 by a second galactose unit (**6**, **7**, **20**, and **21**) or by a xylose unit (**8**). Seven saponins (**4**, **5**, **16–19**, and **26**) were found to be bidesmosides with one glucose unit linked to C-28 of the aglycon. The hemolytic activity of the major saponins (**2**, **3**, **5**, **12–15**, **17**, **24**, and **28**) was measured on sheep erythrocytes in order to establish structure-activity relationships based on the type of sugar attached to the aglycon and on the structure of this aglycon.

The Caryocaraceae is a small family of magnificent trees comprising only two genera *Anthodiscus* (9 species) and *Caryocar* (16 species), distributed in neotropical and tropical America.<sup>1</sup> The *Caryocar* genus, which is called “pequi”<sup>2,3</sup> by the local Indian tribes, are exploited for the oil content of the pulp and the seeds. The resultant edible fats are of potential commercial value for cooking as a substitute for butter, and used in the cosmetic industry and for home-made soaps.<sup>4,5</sup> Apart from scattered reports about lipid composition,<sup>2,3,5</sup> the phytochemistry of the Caryocaraceae has not been studied exhaustively. A short report about *C. microcarpum* mentioned the presence of a large quantity of gallotanin in addition to ellagic acid, gallic acid and methyl gallate, as well as the occurrence of glycosides of oleanolic acid and its hydroxylated derivatives,  $2\beta$ -hydroxyoleanolic acid, hederagenin and bayogenin, in the leaves.<sup>6-7</sup>

*Caryocar glabrum* (Aubl.) Pers. sp. *glabrum* (local name: “sawa” and “peke'a là”) is a large tree (up to 30 m high) which grows in the primary humid forests of northern South America, occurring wild in the northern Amazonia and in Guyana.<sup>1</sup> The fruit is irregularly globular; approximately 5-8 cm in diameter. The peel is yellowish-grey colored and easy to remove. The fleshy pulp (mesocarp) generally surrounds 1 to 2 kidney-shaped seeds. The fruit and the seed are reported to be edible, fresh or cooked. The fruit and the stem bark have been used traditionally by the Indian tribes of French Guyana, Colombia, Venezuela, and Brazil as a remedy for skin problems and as a fish poison.<sup>7</sup>

We herein report the isolation and structural elucidation of 30 saponins. The compounds **1–8**, **22**, **27**, and **30** are glucoside triterpenoids and compounds **9–21**, **23–26**, **28**, and **29** are glucuronide triterpenoids. The structures of these saponins were established by spectroscopic methods (1D and 2D NMR, ESIMS), acid hydrolysis, and by comparison with literature data for known compounds. To the best of our knowledge, compounds **22–30** are known, whereas the remaining twenty-one isolates were found to be new saponins, tentatively named caryocarosides. The Roman numeral I to V indicate the aglycon type: oleanolic acid (I),

hederagenin (II), bayogenin (III),  $2\beta$ -hydroxyoleanolic acid (IV), and gypsogenic acid (V) and the Arabic number is related to the structure of the glycoside chain. In a programme based on the evaluation of the hemolytic activity of saponins and their structure-activity relationship,<sup>8,9</sup> we have measured the hemolytic activity of selected saponins in order to evaluate the influence of a glucuronic acid as compared to a glucose in the sugar chain. Thus, the hemolytic activity of the ten major saponins (**2**, **3**, **5**, **12–15**, **17**, **24**, and **28**) was measured on sheep erythrocytes and some structure-activity relationships were established.

## Results and Discussion

The fruits of *Caryocar glabrum* were collected in French Guyana at Cayena Island. The dried peel and pulp of fruits were extracted separately with methanol, and the methanol extract was evaporated to dryness. Analytical HPLC and TLC analysis of the two methanol extracts revealed that the chromatographic profiles of the peel and the pulp of fruit were qualitatively identical but quantitatively different with a higher amount of bidesmosides in the pulp. The pulp extract was then purified by a combination of silica gel column or RP-18 column chromatography and finally by semi-preparative HPLC or by preparative TLC to afford 30 saponins (**1–30**).

Acid hydrolysis of the crude extract yielded five aglycons. Among them, three were identified as oleanolic acid, hederagenin and bayogenin by TLC with authentic samples. Structural confirmation was achieved by analysis of 1D and 2D NMR spectra of each purified saponin. <sup>13</sup>C NMR spectra were in accordance with data reported in literature.<sup>10,11</sup> The two remaining aglycons were identified by analysis of their NMR spectral data in each intact saponin as  $2\beta$ -hydroxyoleanolic acid<sup>12</sup> and gypsogenic acid.<sup>10,13</sup> The monosaccharides obtained from the acid hydrolysis were identified as D-xylose, D-glucose, D-galactose and D-glucuronic acid by TLC with authentic samples and measurement of their optical rotation after purification. The

$^1\text{H}$  and  $^{13}\text{C}$  NMR data of intact saponins indicated a  $\beta$  configuration for the anomeric position of each sugars (Tables 1-3).

All of the saponins with oleanolic acid as an aglycon were known compounds. These saponins were identified as 3-*O*- $\beta$ -D-galactopyranosyl-(1 $\rightarrow$ 3)- $\beta$ -D-glucopyranosyl oleanolic acid (arvensoside B) (**22**),<sup>14-16</sup> 3-*O*- $\beta$ -D-glucuronopyranosyl oleanolic acid (glycoside F) (**23**),<sup>14,15,17-25</sup> 3-*O*- $\beta$ -D-galactopyranosyl-(1 $\rightarrow$ 3)- $\beta$ -D-glucuronopyranosyl oleanolic acid (glycoside D) (**24**),<sup>14,15,17</sup> 3-*O*- $\beta$ -D-galactopyranosyl-(1 $\rightarrow$ 3)- $\beta$ -D-methyl glucopyranosiduronate oleanolic acid (**25**),<sup>17,26</sup> and 3-*O*- $\beta$ -D-galactopyranosyl-(1 $\rightarrow$ 3)- $\beta$ -D-glucuronopyranosyl oleanolic acid 28-*O*- $\beta$ -D-glucopyranosyl ester (glycoside C) (**26**).<sup>14,15</sup> The other known saponins were identified as 3-*O*- $\beta$ -D-glucopyranosyl hederagenin (colchiside 4) (**27**),<sup>18,27-29</sup> 3-*O*- $\beta$ -D-glucuronopyranosyl hederagenin (**28**),<sup>18-20</sup> 3-*O*- $\beta$ -D-methyl glucopyranosiduronate hederagenin (**29**),<sup>17,18</sup> and 3-*O*- $\beta$ -D-glucopyranosyl bayogenin (**30**).<sup>28</sup>

The aglycon was identified as 2 $\beta$ -hydroxyoleanolic acid in compounds **9**, **11**, **12**, **15**, **16**, **20**. The negative ESIMS<sup>2</sup> of these compounds showed the same ion fragment at  $m/z$  471 [M – H – glycosidic chain]. The  $^1\text{H}$  NMR spectra exhibited seven signals due to tertiary methyl groups, resonating as singlets at  $\delta$  0.85, 0.93, 0.97, 1.10 (6H), 1.18, and 1.27 ( $\delta \pm 0.02$  ppm). The ethylene H-12 signal was observed at  $\delta$  5.27  $\pm$  0.01 ppm (t,  $J = 3.6 \pm 0.2$  Hz). The two proton signals of a vicinal diol (H-2 and H-3) were observed at  $\delta$  4.22 (q,  $J = 3.3 \pm 0.4$  Hz) and 3.20 (d,  $J = 3.3 \pm 0.4$  Hz), respectively (Tables 2 and 3). The small coupling constant between H-2 and H-3 ( $J_{2,3} = 3.3 \pm 0.4$  Hz) indicated that the two hydroxyl groups adopted a vicinal *cis*- $\beta$  configuration, like in bayogenin. The assignments of other proton and carbon signals of the aglycon were accomplished by analysis of the usual 2D COSY, ROESY, HSQC and HMBC experiments. The  $^1\text{H}$  and  $^{13}\text{C}$  NMR values were in full agreement with those reported in the literature for 2 $\beta$ -hydroxyoleanolic acid.<sup>12</sup>

Caryocaroside V-1 (**1**) was the only saponin containing gypsogenic acid as aglycon. The  $^1\text{H}$  NMR spectrum of **1** exhibited six signals due to tertiary methyl groups, resonating as singlets at  $\delta$  0.83, 0.93, 0.96, 1.00, 1.17, and 1.19, one hydroxymethine signal (H-3) at  $\delta$  4.11 (dd,  $J = 11.8 - 4.5$  Hz), and one ethylene proton signal at  $\delta$  5.27 (t,  $J = 3.6$  Hz). The  $^{13}\text{C}$  NMR spectrum of this compound indicated the presence of two carbonyl groups at  $\delta$  181.9 and 182.0 that were assigned to C-23 and C-28 by analysis of the HMBC experiment (Table 1). The  $^{13}\text{C}$  NMR values of the aglycon were in good agreement with those reported for gypsogenic acid.<sup>10,13</sup> The positive ESIMS of **1** exhibited a molecular ion peak at  $m/z$  811 [ $\text{M} + \text{H}$ ]<sup>+</sup> and a positive fragment at  $m/z$  487 [aglycon + H]<sup>+</sup> attributed to the loss of a disaccharide moiety consisting of two hexoses. Further analysis of the  $^1\text{H}$  and  $^{13}\text{C}$  NMR spectra of **1** revealed the presence of two anomeric protons at  $\delta$  4.37 and 4.52 correlated in the HSQC spectrum with two anomeric carbons at  $\delta$  104.9 and 105.7, respectively (Table 1). Complete assignment of each glycoside proton system was achieved by analysis of COSY and TOCSY experiments. A  $\beta$ -D-glucose unit was identified starting from the anomeric proton at  $\delta$  4.37 (d,  $J = 7.7$  Hz). The second monosaccharide whose anomeric proton resonates at  $\delta$  4.52 (d,  $J = 7.6$  Hz) was identified as a  $\beta$ -D-galactose, characterized by its equatorial proton H-4'' at  $\delta$  3.82 (d,  $J_{3,4} = 3.5$  Hz) (Table 1). The deshielding of C-3' ( $\delta$  87.3) of glucose suggested the linkage site with the galactose (Table 1). In the HMBC experiment, the cross peaks observed between H-1'' of galactose and C-3' of glucose and between H-1' of glucose and C-3 of the gypsogenic acid ( $\delta$  85.5) led to the assignment of caryocaroside V-1 (**1**) as 3-O- $\beta$ -D-galactopyranosyl-(1 $\rightarrow$ 3)- $\beta$ -D-glucopyranosyl gypsogenic acid.

Caryocarosides II-1 (**2**) and III-1 (**3**) exhibited molecular ion peaks [ $\text{M} - \text{H}$ ]<sup>-</sup> at  $m/z$  795 and 811 in the negative ESIMS. The MS<sup>2</sup> experiments of the [ $\text{M} - \text{H}$ ]<sup>-</sup> ion of both **2** and **3** gave the same negative fragments [ $\text{M} - \text{H} - 162$ ]<sup>-</sup> and [ $\text{M} - \text{H} - 2 \times 162$ ]<sup>-</sup> attributed to the successive loss of two hexoses. The  $^1\text{H}$  and  $^{13}\text{C}$  NMR spectra of **2** and **3** indicated two

anomeric carbons at  $\delta$  105.1 and 105.5 for **2** and at  $\delta$  105.2 and 105.8 for **3** which were correlated in the HSQC experiment with anomeric protons at  $\delta$  4.48 (d,  $J = 7.9$  Hz) and 4.52 (d,  $J = 7.7$  Hz) for **2** and 4.52 (d,  $J = 7.8$  Hz) and 4.55 (d,  $J = 7.7$  Hz) for **3**. Comparison of the glycosidic  $^1\text{H}$  and  $^{13}\text{C}$  NMR values for **2** and **3** with those of **1** showed that all these compounds contained the same glycoside chain ( $\beta$ -D-galactopyranosyl-(1 $\rightarrow$ 3)- $\beta$ -D-glucopyranoside). Thus, caryocarosides II-1 (**2**) and III-1 (**3**) were concluded to be 3-*O*- $\beta$ -D-galactopyranosyl-(1 $\rightarrow$ 3)- $\beta$ -D-glucopyranosyl hederagenin and 3-*O*- $\beta$ -D-galactopyranosyl-(1 $\rightarrow$ 3)- $\beta$ -D-glucopyranosyl bayogenin, respectively.

Caryocarosides II-2 (**4**) and III-2 (**5**) gave a molecular ion peak  $[\text{M} - \text{H}]^-$  at  $m/z$  957 and 973 in the negative ESIMS and the  $\text{MS}^2$  experiments of these ions gave the same fragment  $[\text{M} - \text{H} - 162]^-$  at  $m/z$  795 and 811, respectively, suggesting a supplementary hexose unit compared to **2** and **3**. The  $^{13}\text{C}$  NMR spectrum of **4** exhibited three anomeric carbons at  $\delta$  95.7, 105.3, and 105.7, correlated in the HSQC spectrum with three anomeric protons at  $\delta$  5.40, 4.48, and 4.52, respectively (Table 1). Further analysis of the 2D NMR experiments revealed that the sugar chain linked at C-3 of the hederagenin was  $\beta$ -D-galactopyranosyl-(1 $\rightarrow$ 3)- $\beta$ -D-glucopyranoside as in compounds **1**, **2**, and **3**. The values  $\delta_{\text{H}}$  5.40 and  $\delta_{\text{C}}$  95.7 belonged to the anomeric signals of a glycoside ester unit and the  $^{13}\text{C}$  NMR signal due to the C-28 of the aglycon moiety ( $\delta$  178.1) indicated a glycosylation of the carboxyl group. Consequently, compound **4** was a bidesmoside saponin. The COSY and TOCSY experiments led to the assignment of the glycoside ester unit as a  $\beta$ -D-glucose unit. A cross peak was observed in the HMBC experiment between H-1''' of this glucose and C-28 of hederagenin. The above evidences led to the assignment of caryocaroside II-2 (**4**) as 3-*O*- $\beta$ -D-galactopyranosyl-(1 $\rightarrow$ 3)- $\beta$ -D-glucopyranosyl hederagenin 28-*O*- $\beta$ -D-glucopyranosyl ester. Comparison of the  $^1\text{H}$  and  $^{13}\text{C}$  NMR values of the saccharide part in compounds **4** and **5** showed that they contain the same glycoside chains linked at C-3 and C-28 positions. Thus, the structure of



caryocaroside III-2 (**5**) was concluded to be 3-*O*- $\beta$ -D-galactopyranosyl-(1 $\rightarrow$ 3)- $\beta$ -D-glucopyranosyl bayogenin 28-*O*- $\beta$ -D-glucopyranosyl ester.

Caryocarosides II-3 (**6**) and III-3 (**7**) displayed a molecular ion peak  $[M - H]^-$  at  $m/z$  957 and 973 respectively, in the negative ESIMS. The MS<sup>2</sup> experiment gave negative fragments at  $m/z$  795, 633 and 471 for **6** and at  $m/z$  811, 649, and 487 for **7** attributed to the successive losses of three hexoses  $[M - H - 162]^-$ ,  $[M - H - 2 \times 162]^-$ , and  $[M - H - 3 \times 162]^-$ . The <sup>1</sup>H NMR spectrum of **6** exhibited three anomeric protons at  $\delta$  4.48, 4.52, and 4.62, correlated in the HSQC experiment with their respective anomeric carbons at  $\delta$  105.3, 106.3, and 105.3 (Table 1). Complete assignment of each glycoside proton system was achieved by analysis of COSY, TOCSY and ROESY experiments. The sugar units with anomeric protons at  $\delta$  4.62 (d,  $J = 7.8$  Hz) and 4.52 (d,  $J = 7.6$  Hz) corresponded to two  $\beta$ -D-galactose units. The third glycosidic unit with the anomeric proton at  $\delta$  4.48 (d,  $J = 7.9$  Hz) was identified as a  $\beta$ -D-glucose unit (Table 1). The downfield shifts of C-3'' ( $\delta$  84.5) of the inner galactose moiety and C-3' ( $\delta$  87.9) of the glucose moiety, and their correlations in the HMBC spectrum observed between H-1''' ( $\delta$  4.52) of the terminal galactose and C-3'' of the inner galactose, between H-1'' ( $\delta$  4.62) of this galactose and C-3' of the glucose, and between H-1' ( $\delta$  4.48) of the glucose and C-3 ( $\delta$  83.6) of hederagenin showed that caryocaroside II-3 (**6**) is 3-*O*- $\beta$ -D-galactopyranosyl-(1 $\rightarrow$ 3)- $\beta$ -D-galactopyranosyl-(1 $\rightarrow$ 3)- $\beta$ -D-glucopyranosyl hederagenin. Comparison of the <sup>1</sup>H and <sup>13</sup>C NMR values of the trisaccharide chain of compounds **6** and **7** showed that **7** contained the same trisaccharide. Thus, caryocaroside III-3 (**7**) was concluded to be 3-*O*- $\beta$ -D-galactopyranosyl-(1 $\rightarrow$ 3)- $\beta$ -D-galactopyranosyl-(1 $\rightarrow$ 3)- $\beta$ -D-glucopyranosyl bayogenin.

The MS<sup>2</sup> experiment of the  $[M - H]^-$  molecular ion peak of caryocaroside III-4 (**8**) observed at  $m/z$  943 in the negative ESIMS gave a negative fragment at  $m/z$  811  $[M - H - 132]^-$ , suggesting a supplementary terminal pentose unit compared to **7**. Analysis of 2D NMR

experiments of **8** revealed that the nature of the supplementary pentose was a terminal  $\beta$ -D-xylose with anomeric signals at  $\delta_{\text{H}}$  4.53 (d,  $J = 7$  Hz) and  $\delta_{\text{C}}$  105.2 (Table 1). The ROE interactions observed in the ROESY spectrum between H-1''', H-3''', and H-5''' of the xylose unit confirmed the  $\alpha$ -axial orientation of these three protons and the  $\beta$ -anomeric configuration. The sequence of the trisaccharide chain was deduced from the cross peaks observed in the HMBC spectrum between H-1''' of the terminal xylose and C-3'' of the galactose, H-1'' of the galactose and C-3' of the glucose, and H-1' of the glucose and C-3 of bayogenin. This sequence was confirmed by the observation of the ROE interactions between the protons involved in the interglycosidic linkages: H-1''' of xylose/H-3'' of galactose, H-1'' of galactose/H-3' of glucose and H-1' of glucose/H-3 of bayogenin. Based on the foregoing evidence, the structure of caryocaroside III-4 (**8**) was established as 3-*O*- $\beta$ -D-xylopyranosyl-(1 $\rightarrow$ 3)- $\beta$ -D-galactopyranosyl-(1 $\rightarrow$ 3)- $\beta$ -D-glucopyranosyl bayogenin.

The  $^1\text{H}$  NMR spectrum of caryocarosides IV-5 (**9**) and III-5 (**10**) exhibited anomeric proton doublets at  $\delta$  4.46 ( $J = 7.5$  Hz) and 4.51 ( $J = 7.7$  Hz), respectively, which gave correlations with anomeric carbons at  $\delta$  106.5 and 105.3 in the HSQC experiment (Table 2). Analysis of COSY and HSQC experiments identified a  $\beta$ -D-glucuronic acid with the carbonyl C-6' at  $\delta$  173.0 (**9**) and 171.0 (**10**). In addition, the  $[\text{M} - \text{H} - 176]^-$  negative ion observed at  $m/z$  471 (**9**) and 487 (**10**) after  $\text{MS}^2$  fragmentation of the molecular ion  $[\text{M} - \text{H}]^-$ , in the negative ESIMS, confirmed the presence of the glucuronic acid. The cross peaks observed in the HMBC experiments between C-3 of aglycon and H-1' of glucuronic acid and the ROE interactions observed in the ROESY experiments between H-1' of glucuronic acid and H-3 of aglycon indicated that the glucuronic acid was attached to C-3 of 2 $\beta$ -hydroxyoleanolic acid in **9** and to C-3 of bayogenin in **10**. From these data, the structures of caryocarosides IV-5 (**9**) and III-5 (**10**) were established as 3-*O*- $\beta$ -D-glucuronopyranosyl-2 $\beta$ -hydroxyoleanolic acid and 3-*O*- $\beta$ -D-glucuronopyranosyl bayogenin, respectively.

Caryocaroside IV-6 (**11**) showed a molecular ion peak  $[M - H]^-$  at  $m/z$  661 that gave ion fragments at  $m/z$  647  $[M - H - 15]^-$  and 471  $[M - H - (175 + 15)]^-$  in the negative ESIMS<sup>2</sup>, attributed to the losses of a methyl group and a methyl hexosuronate. These data suggested an additional methyl group compared to **9**. This hypothesis was confirmed by the presence of the NMR signal of a methoxy group at  $\delta_H$  3.78 (s, 3H) and  $\delta_C$  52.9 (Table 2). Nearly all of the <sup>13</sup>C NMR signals were superposable with those of **9** except for C-6' of glucuronic acid which exhibited an upfield shift at  $\delta$  171.5. Moreover, in the HMBC experiment, a cross peak was observed between the protons of the methoxy group and the carbonyl C-6' of glucuronic acid. Thus, caryocaroside IV-6 (**11**) is 3-*O*- $\beta$ -D-methyl glucopyranosiduronate-2 $\beta$ -hydroxyoleanolic acid.

The negative ESIMS experiments of caryocarosides IV-7 (**12**) and II-7 (**13**) gave both molecular ion peaks  $[M - H]^-$  at  $m/z$  809, whereas the molecular ion peak was observed at  $m/z$  825 for caryocaroside III-7 (**14**). The MS<sup>2</sup> experiments of the  $[M - H]^-$  ion of these saponins gave the same ion fragment  $[M - H - 162 - 176]^-$  attributed to the loss of a disaccharide consisting of a hexose and a hexosuronic acid. The <sup>1</sup>H and <sup>13</sup>C NMR spectra of **12** confirmed the presence of two sugar residues with their anomeric carbons at  $\delta$  104.6 and 104.9 correlating in the HSQC experiment with the anomeric protons at  $\delta$  4.40 (d,  $J = 8.1$  Hz) and 4.33 (d,  $J = 7.7$  Hz) (Table 2). Analysis of 2D NMR experiments revealed the presence of a terminal  $\beta$ -D-galactose ( $\delta_H$  4.33) and a  $\beta$ -D-glucuronic acid ( $\delta_H$  4.40) monosubstituted at position C-3' ( $\delta_C$  85.7). The HMBC spectrum showed cross peaks between H-1" of galactose and C-3' of glucuronic acid and between H-1' of glucuronic acid and C-3 of 2 $\beta$ -hydroxyoleanolic acid. This evidence led to the assignment of caryocaroside IV-7 (**12**) as 3-*O*- $\beta$ -D-galactopyranosyl-(1 $\rightarrow$ 3)- $\beta$ -D-glucuronopyranosyl-2 $\beta$ -hydroxyoleanolic acid. Comparison of the <sup>1</sup>H and <sup>13</sup>C NMR spectral data of the disaccharide moieties of **13** and **14** with those of **12** showed that hederagenin (**13**) and bayogenin (**14**) were both substituted at C-

3 by the same disaccharide chain (Table 2). Thus, the structure of caryocaroside II-7 (**13**) was elucidated as 3-*O*- $\beta$ -D-galactopyranosyl-(1 $\rightarrow$ 3)- $\beta$ -D-glucuronopyranosyl hederagenin and caryocaroside III-7 (**14**) as 3-*O*- $\beta$ -D-galactopyranosyl-(1 $\rightarrow$ 3)- $\beta$ -D-glucuronopyranosyl bayogenin.

As observed for caryocaroside IV-6 (**11**), the  $^1\text{H}$  and  $^{13}\text{C}$  NMR spectra of caryocaroside IV-8 (**15**) showed an additional methoxy group compared to **12** (Table 2). The MS<sup>2</sup> experiment of its  $[\text{M} - \text{H}]^-$  molecular ion at  $m/z$  823, gave fragments at  $m/z$  791  $[\text{M} - \text{H} - \text{CH}_3\text{OH}]^-$  and 471  $[\text{M} - \text{H} - 162 - (175 + 15)]^-$  due to the elimination of the methoxy group and to the loss of a disaccharide consisting of a hexose and a methyl hexosuronate. The methoxy group observed in the  $^1\text{H}$  NMR at  $\delta_{\text{H}}$  3.80 (s, 3H) was correlated in the HMBC spectrum with the carbonyl C-6' of glucuronic acid. Thus, caryocaroside IV-8 (**15**) is 3-*O*- $\beta$ -D-galactopyranosyl-(1 $\rightarrow$ 3)- $\beta$ -D-methyl glucopyranosiduronate-2 $\beta$ -hydroxyoleanolic acid.

In the negative ESIMS, the molecular ion peaks  $[\text{M} - \text{H}]^-$  of caryocarosides IV-9 (**16**) and II-9 (**17**) were both observed at  $m/z$  971, whereas, in caryocaroside III-9 (**18**) the molecular ion peak was observed at  $m/z$  987. The MS<sup>2</sup> experiment of these ions gave fragments suggesting an additional hexose compared to **12**, **13**, and **14**, respectively. Analysis of the  $^1\text{H}$  and  $^{13}\text{C}$  NMR spectra of compounds **16**, **17** and **18** revealed that the C-3 of the aglycon was substituted by the disaccharide [ $\beta$ -D-galactopyranosyl-(1 $\rightarrow$ 3)- $\beta$ -D-glucuronopyranoside] as in compounds **12**, **13**, and **14**. The additional hexose unit was identified as a terminal  $\beta$ -D-glucopyranose with anomeric signals at  $\delta_{\text{H}}$  5.40 and  $\delta_{\text{C}}$  95.7 (Table 3), linked to C-28 of the aglycon as deduced from the cross peaks observed in the HMBC spectrum between its H-1''' and the C-28 ( $\delta$  178.1) of the aglycon. Based on the foregoing evidence, the structures of caryocarosides IV-9, II-9, and III-9 were concluded to be 3-*O*- $\beta$ -D-galactopyranosyl-(1 $\rightarrow$ 3)- $\beta$ -D-glucuronopyranosyl-2 $\beta$ -hydroxyoleanolic acid 28-*O*- $\beta$ -D-glucopyranosyl ester (**16**), 3-*O*- $\beta$ -D-galactopyranosyl-(1 $\rightarrow$ 3)- $\beta$ -D-glucuronopyranosyl hederagenin 28-*O*- $\beta$ -D-glucopyranosyl

ester (**17**), and 3-*O*- $\beta$ -D-galactopyranosyl-(1 $\rightarrow$ 3)- $\beta$ -D-glucuronopyranosyl bayogenin 28-*O*- $\beta$ -D-glucopyranosyl ester (**18**), respectively.

Comparison of the  $^1\text{H}$  and  $^{13}\text{C}$  NMR and ESIMS of caryocaroside II-10 (**19**) with those of II-9 (**16**) showed that **19** also possessed an additional methoxy group. Furthermore, in the HMBC experiment, a cross peak was observed between the protons of this methyl ( $\delta_{\text{H}}$  3.80) and the carbonyl C-6' ( $\delta_{\text{C}}$  171.0) of glucuronic acid. Thus, caryocaroside II-10 (**19**) was elucidated as 3-*O*- $\beta$ -D-galactopyranosyl-(1 $\rightarrow$ 3)- $\beta$ -D-methylglucopyranosiduronate hederagenin 28-*O*- $\beta$ -D-glucopyranosyl ester.

The negative ESIMS of caryocarosides IV-11 (**20**) and II-11 (**21**) showed that they were isomers to **16** and **17**. The  $^{13}\text{C}$  NMR spectrum of **20** and **21** showed three anomeric carbon signals at  $\delta$  105.1, 106.3 (2C) for **20** and 105.2, 105.4 and 106.3 for **21**. Analysis of the 2D NMR experiments allowed full identification of a terminal  $\beta$ -D-galactose ( $\delta_{\text{H}}$  4.53 and  $\delta_{\text{C}}$  106.3), a  $\beta$ -D-galactose ( $\delta_{\text{H}}$  4.66 or 4.64 and  $\delta_{\text{C}}$  105.1 or 105.2) monosubstituted at position C3'' ( $\delta_{\text{C}}$  84.5), and a  $\beta$ -D-glucuronic acid monosubstituted at position C-3' ( $\delta_{\text{C}}$  85.9 or 87.3) (Table 3). The sequence of the trisaccharide chain was deduced from the cross peaks observed in the HMBC experiments between H-1''' of the terminal galactose and C-3'' of the inner galactose, between H-1'' of the inner galactose and C-3' of the glucuronic acid and between H-1' of the glucuronic acid and C-3 of the aglycon. This evidence led to the assignment of caryocaroside IV-11 (**20**) as 3-*O*- $\beta$ -D-galactopyranosyl-(1 $\rightarrow$ 3)- $\beta$ -D-galactopyranosyl-(1 $\rightarrow$ 3)- $\beta$ -D-glucuronopyranosyl-2 $\beta$ -hydroxyoleanolic acid and caryocaroside II-11 (**21**) as 3-*O*- $\beta$ -D-galactopyranosyl-(1 $\rightarrow$ 3)- $\beta$ -D-galactopyranosyl-(1 $\rightarrow$ 3)- $\beta$ -D-glucuronopyranosyl hederagenin.

The ability of the methanol extracts of the pulp and the peel of *C. glabrum* fruits, and of the ten major isolated saponins (**2**, **3**, **5**, **12–15**, **17**, **24**, and **28**) to lyse sheep erythrocytes

(10% suspension phosphate buffer) were evaluated *in vitro* using the method previously described.<sup>8</sup> The tested saponins were generally less active than the dialyzed saponin mixture from Sigma<sup>®</sup> (Sigma<sup>®</sup>D) used as a reference (Table 4). The hemolytic activity of the monodesmosidic saponins **3** and **13** was much higher than the corresponding bidesmosidic saponins **5** and **17**. Comparison of the hemolytic activity of saponins **28** and **13**, showed that the disaccharide saponin **13** was more active than the corresponding monosaccharide **28**. These results confirmed the fact that bidesmoside saponins are generally less hemolytic than monodesmoside saponins and that hemolytic activity increases with the number of sugar units linked at position 3 of the aglycon.<sup>8</sup> In our previous study, we established that hemolytic activity increased with the polarity of ring A (hydroxyl group at C-23 and/or C-2). This observation was confirmed with saponins containing a  $\beta$ -D-glucose linked at position C-3 of the aglycon. Caryocaroside III-1 (**3**) was three fold more active than caryocaroside II-1 (**2**). To evaluate the influence of a glucose and a glucuronic acid on the hemolytic activity, we compared the activity of saponins **2** and **3** with **13** and **14**, respectively. With hederagenin as aglycon (**2** and **13**), the presence of a glucuronic acid slightly enhanced hemolytic activity, but with bayogenin (**3** and **14**), we observed the reverse effect; glucuronic acid decreased the hemolytic activity. It is known that the effect of sugar residue is not transferable from one aglycon to one another.<sup>30</sup> With saponins including a  $\beta$ -D-glucuronic acid linked at position 3 of the aglycon (**24**, **12**, **13** and **14**), the hemolytic activity decreased when the polarity of ring A increased. We suggest that the addition of a hydroxyl group of ring A (at C-23 and/or C-2) might interfere with the free carboxylic acid group of the glucuronic acid in the hemolytic mechanism. Caryocaroside IV-7 (**12**) was five times more hemolytic than caryocaroside IV-8 (**15**), which contains a methoxy group attached to C-6' of the glucuronic acid. The esterified glucuronic acid decreased the polarity of the saponin and led to a decrease in the hemolytic activity, confirming the important role of polarity of ring A on hemolytic activity.

## Experimental Section

**General Experimental procedures.** Optical rotations were measured in MeOH or H<sub>2</sub>O with a Perkin-Elmer 241 polarimeter. <sup>1</sup>H and <sup>13</sup>C NMR spectra were recorded in CD<sub>3</sub>OD or DMSO-*d*<sub>6</sub> on a Bruker Avance DRX - 500 spectrometer (<sup>1</sup>H at 500 MHz and <sup>13</sup>C at 125 MHz), and 2D - NMR experiments were performed using standard Bruker microprograms (XWIN-NMR version 2.6 software). ESIMS and MS-MS experiments were performed using a Bruker Esquire-LC ion trap mass spectrometer. The samples were introduced by infusion in a solution of MeOH. TLC were carried out on precoated silica gel 60 F<sub>254</sub> (Merck), with CHCl<sub>3</sub>-MeOH-H<sub>2</sub>O (14:6:1), and spots were visualized by spraying with 50% H<sub>2</sub>SO<sub>4</sub>. Column chromatography (CC) was carried out on Kieselgel 60 (63-200 μm, Merck) or LiChroprep RP-18 (40-63 μm, Merck). HPLC was performed on a Dionex apparatus equipped with an ASI-100 automated sample injector, a STH 585 column oven, a P580 pump, a diode array detector UVD 340S and the Chromeleon<sup>®</sup> software. Prepacked C<sub>18</sub> reversed phase column (201SPTM, 4.6 x 250 mm, 5 μm, 90 Å, Dionex, vydac, France) was used for analytical HPLC with a binary gradient elution (solvent A: H<sub>2</sub>O – TFA, 0.025 %, solvent B: MeCN) and a flow rate of 1 mL.min<sup>-1</sup>, the chromatogram was monitored at 205 and 210 nm. Prepacked C<sub>18</sub> reversed phase column (201SP510, 10 x 250 mm, 5 μm, 90 Å, Dionex, vydac, France) was used for semi-preparative HPLC with a binary gradient elution (solvent A: H<sub>2</sub>O – TFA 0.0025%, solvent B: MeCN) and a flow rate of 3 mL.min<sup>-1</sup>, the chromatogram was monitored at 205 and 210 nm.

**Plant Material.** The fruits of *C. glabrum* were collected in two locality of French Guyana: the Amirande forest near Matoury, and the Ecerex forest station near Sinnamary in October 2001. The species was identified by M.F. Prevost of the botany laboratory of the IRD Centre of Cayena (French Guyana), and a herbarium specimen (Prevost MFP 4864) was deposited in the Herbarium of Guyana.<sup>31</sup>

**Hemolytic Assays.** This assay was performed as described previously.<sup>8</sup> The 10% sheep erythrocyte suspension (10%) was obtained by dilution of a commercial 50% suspension from Biomerieux<sup>®</sup>, Lyon with phosphate buffer saline (PBS). Saponins **2**, **3**, **5**, **12–15**, **17**, **24**, and **28** were prepared in triplicate with concentrations ranging from 1 to 500  $\mu\text{g/mL}$  in PBS. 25  $\mu\text{L}$  of erythrocyte suspension were added to 1 mL of the sample and rapidly stirred. Absorbance of the supernatant was measured at 540 nm after 60 min of incubation and centrifuged for 5 min at 3000 rpm.  $\text{HD}_{50}$  and  $\text{HD}_{100}$  values were the concentrations of sample which causes 50 and 100% of hemolysis. The hemolytic activity was measured with regard to a dialysed saponin from Sigma<sup>®</sup> used as reference standard, which caused 100% hemolysis at 25  $\mu\text{g/mL}$ .

**Extraction and Isolation.** The air-dried powdered pulp (mesocarp) (35 g) and peel (pericarp) (84 g) of fruit were boiled under reflux in methanol (1 and 2 L, respectively) for 3 h. After cooling and filtration, the methanol extract was evaporated to provide the saponin mixture as a brown residue (20 g; yield 57% for the pulp and 26 g; yield 31% for the peel). The saponin mixture was analyzed by TLC on silica gel using the mixture of solvent  $\text{CHCl}_3$ -MeOH- $\text{H}_2\text{O}$  (14:6:1) and by analytical HPLC over  $\text{C}_{18}$  with the gradient elution program 30 to 50% B for 75 min.

A part of the methanol extract of the pulp (6 g) was subjected to column chromatography on silica gel (240 g, 2.5 x 38 cm) using a gradient of  $\text{CHCl}_3$ -MeOH- $\text{H}_2\text{O}$  (95:5:0 to 60:40:7) to give 191 fractions of 250 mL. Fractions 56-58 eluted with  $\text{CHCl}_3$ -MeOH (8:2) were purified by preparative TLC in  $\text{CHCl}_3$ -MeOH (75:25) to give **27** (19 mg) and **30** (11 mg). Fraction 63 eluted with  $\text{CHCl}_3$ -MeOH (8:2), contains compound **2** (103 mg). Fractions 64-69 eluted with  $\text{CHCl}_3$ -MeOH (8:2) were purified by preparative TLC in  $\text{CHCl}_3$ -MeOH- $\text{H}_2\text{O}$  (70:30:3) to give **3** (28 mg) and **22** (6 mg). Fractions 83-86, eluted with  $\text{CHCl}_3$ -MeOH (7:3),



were purified on a silica gel column chromatography using the gradient CHCl<sub>3</sub>-MeOH (85:15 to 7:3), and fractions eluted with CHCl<sub>3</sub>-MeOH (85:15) were purified further by reversed-phase C<sub>18</sub> column chromatography using a gradient of MeOH-H<sub>2</sub>O (5:5 to 7:3) to give **4** (11 mg) and **7** (5 mg). Fractions [87-90], eluted with CHCl<sub>3</sub>-MeOH (7:3), were purified by RP-18 column chromatography, eluting with MeOH-H<sub>2</sub>O (5:5 to 7:3), to give **5** (29 mg). Fractions 91-100 eluted with CHCl<sub>3</sub>-MeOH-H<sub>2</sub>O (70:30:1) were passed through an ion exchange IRN 77 (H<sup>+</sup>) Amberlite<sup>®</sup> resin column before purification by RP-18 column chromatography, eluting with MeOH-H<sub>2</sub>O (5:5 to 8:2), to give **6** (5 mg), **8** (13 mg) and **29** (11 mg). Fractions 101-105, eluted with CHCl<sub>3</sub>-MeOH-H<sub>2</sub>O (70:30:1), were purified by RP-18 column chromatography, using a gradient of MeOH-H<sub>2</sub>O (5:5 to 8:2), to give **23** (12 mg) and **28** (30 mg). Fractions eluted with MeOH-H<sub>2</sub>O (5:5) were further purified by semi-preparative HPLC with a linear gradient (40 to 55% B) for 15 min to afford 2 mg of **1** (*R<sub>t</sub>* = 12.4 min), and fractions eluted with MeOH-H<sub>2</sub>O (8:2) were purified by semi-preparative HPLC with a linear gradient (30 to 43% B) for 30 min to afford 7 mg of **9** (*R<sub>t</sub>* = 12.4 min). Fractions 119-124, eluted with CHCl<sub>3</sub>-MeOH-H<sub>2</sub>O (70:30:1), were purified by RP-18 column chromatography eluting with MeOH-H<sub>2</sub>O (45:55 to 8:2), to give **12** (17 mg), **13** (58 mg), and **24** (21 mg), while fractions eluted with MeOH-H<sub>2</sub>O (45:55) were purified by semi-preparative HPLC with a linear gradient (35 to 41% B) for 25 min, to give 4 mg of **10** (*R<sub>t</sub>* = 10.5 min). Fractions 125-137, eluted with CHCl<sub>3</sub>-MeOH-H<sub>2</sub>O (70:30:2), were purified by RP-18 column chromatography using a gradient of MeOH-H<sub>2</sub>O (35:65 to 8:2), to give **25** (11 mg). Fractions eluted with MeOH-H<sub>2</sub>O (7:3) were further purified by semi-preparative HPLC, with a linear gradient (30 to 43% B) for 30 min, to give 8 mg of **14** (*R<sub>t</sub>* = 23.6 min) and 2 mg of **19** (*R<sub>t</sub>* = 19.9 min), and fractions eluted with MeOH-H<sub>2</sub>O (8:2) were further purified by silica gel column chromatography to give **11** (2 mg), **15** (18 mg), and **24** (6 mg). Fractions 152-155, eluted with CHCl<sub>3</sub>-MeOH-H<sub>2</sub>O (70:30:5), were purified by RP-18 column chromatography, eluting with MeOH-H<sub>2</sub>O (3:7 to 8:2), to give **16** (3 mg) and **17** (18 mg). Fractions eluted with

MeOH-H<sub>2</sub>O (6:4) were then purified by semi-preparative HPLC with a linear gradient of 30 to 38% B for 30 min, to give 3 mg of **26** ( $R_t = 25.9$  min). Fractions 156-170, eluted with CHCl<sub>3</sub>-MeOH-H<sub>2</sub>O (70:30:5), were purified by RP-18 column chromatography, using a gradient of MeOH-H<sub>2</sub>O (3:7 to 8:2). The fraction eluted with MeOH-H<sub>2</sub>O (3:7) was further purified by a semi-preparative HPLC with a linear gradient (30 to 35% B) for 20 min to give 4 mg of **18** ( $R_t = 12.9$  min) and 7 mg of **21** ( $R_t = 13.9$  min). Fractions eluted with MeOH-H<sub>2</sub>O (4:6) was then purified by semi-preparative HPLC with a linear gradient (40 to 50% B) for 20 min to give 4 mg of **20** ( $R_t = 15.5$  min).

**Acid hydrolysis of Saponin Mixture.** An aliquot of the crude saponin mixture (1 g) was refluxed with 60 mL of 2 N HCl for 4.5 h. The sapogenin mixture was extracted with EtOAc (3 x 30 mL), washed with H<sub>2</sub>O, and evaporated to dryness. Oleanolic acid, hederagenin and bayogenin were identified from the sapogenin residue with authentic sample by TLC in CHCl<sub>3</sub>-MeOH (98:2). The acid aqueous layer was neutralized with 0.5 M NaOH and freeze-dried. Four sugars were identified and compared with authentic samples by TLC using MeCOEt-isoPrOH-Me<sub>2</sub>CO-H<sub>2</sub>O (20:10:7:6) as xylose, glucose, galactose and glucuronic acid. After preparative TLC of the sugar mixture (100 mg) in this solvent, the optical rotation of each purified sugar was measured.

**Caryocaroside V-1 (1):** White powder;  $[\alpha]_D^{20} + 16.7$  ( $c$  0.17, MeOH); <sup>1</sup>H NMR of the aglycon (CD<sub>3</sub>OD, 500 MHz), see Table 1; <sup>13</sup>C NMR of the aglycon (CD<sub>3</sub>OD, 125 MHz)  $\delta$  12.0 (C-24), 16.2 (C-25), 17.6 (C-26), 21.9 (C-6), 24.0 (C-30), 24.0 (C-16), 24.5 (C-11), 26.4 (C-27), 26.4 (C-2), 28.9 (C-15), 31.6 (C-20), 33.6 (C-29), 33.8 (C-7), 34.0 (C-22), 35.0 (C-21), 37.4 (C-10), 39.7 (C-1), 40.8 (C-8), 42.7 (C-18), 42.9 (C-14), 47.2 (C-19), 47.6 (C-17), 49.0 (C-9), 52.9 (C-5), 54.2 (C-4), 85.8 (C-3), 123.4 (C-12), 145.2 (C-13), 181.9 (C-23),

182.0 (C-28);  $^1\text{H}$  and  $^{13}\text{C}$  NMR of the glycosidic part, see Table 1; ESIMS (positive-ion mode)  $m/z$  833  $[\text{M} + \text{Na}]^+$ , 811  $[\text{M} + \text{H}]^+$ , 487  $[\text{M} + \text{Na} - \text{Glc} - \text{Gal}]^+$ .

**Caryocaroside II-1 (2):** White powder;  $[\alpha]_{\text{D}}^{20} + 31.9$  ( $c$  0.83, MeOH);  $^1\text{H}$  NMR of the aglycon ( $\text{CD}_3\text{OD}$ , 500 MHz), see Table 1;  $^{13}\text{C}$  NMR of the aglycon ( $\text{CD}_3\text{OD}$ , 125 MHz)  $\delta$  13.4 (C-24), 16.4 (C-25), 17.7 (C-26), 18.8 (C-6), 24.0 (C-16), 24.0 (C-30), 24.5 (C-11), 26.2 (C-2), 26.5 (C-27), 28.7 (C-15), 31.5 (C-20), 33.3 (C-7), 33.7 (C-22), 33.9 (C-29), 34.9 (C-21), 37.6 (C-10), 39.3 (C-1), 40.4 (C-8), 42.6 (C-18), 42.9 (C-14), 43.7 (C-4), 47.2 (C-19), 47.6 (C-17), 48.2 (C-5), 48.8 (C-9), 65.1 (C-23), 83.7 (C-3), 123.4 (C-12), 145.2 (C-13), 182.2 (C-28);  $^1\text{H}$  and  $^{13}\text{C}$  NMR of the glycosidic part, see Table 1; ESIMS (negative-ion mode)  $m/z$  795  $[\text{M} - \text{H}]^-$ ; ESIMS-MS  $\text{MS}^2$  (795)  $m/z$  633  $[\text{M} - \text{H} - \text{Gal}]^-$ , 471  $[\text{M} - \text{H} - \text{Gal} - \text{Glc}]^-$ ; ESIMS (positive-ion mode)  $m/z$  835  $[\text{M} + \text{K}]^+$ , 819  $[\text{M} + \text{Na}]^+$ ; ESIMS-MS  $\text{MS}^2$  (819)  $m/z$  775  $[\text{M} + \text{Na} - \text{CO}_2]^+$ , 657  $[\text{M} + \text{Na} - \text{Gal}]^+$ , 613  $[\text{M} + \text{Na} - \text{CO}_2 - \text{Gal}]^+$ .

**Caryocaroside III-1 (3):** White powder;  $[\alpha]_{\text{D}}^{20} + 24.9$  ( $c$  1, MeOH);  $^1\text{H}$  NMR of the aglycon ( $\text{CD}_3\text{OD}$ , 500 MHz), see Table 1;  $^{13}\text{C}$  NMR of the aglycon ( $\text{CD}_3\text{OD}$ , 125 MHz)  $\delta$  14.7 (C-24), 17.5 (C-25), 18.3 (C-26), 18.7 (C-6), 24.4 (C-30), 24.5 (C-16), 24.7 (C-11), 26.6 (C-27), 29.2 (C-15), 31.8 (C-20), 33.6 (C-7), 34.0 (C-29), 34.4 (C-22), 35.5 (C-21), 37.6 (C-10), 40.6 (C-8), 43.0 (C-4), 43.3 (C-14), 43.5 (C-18), 44.4 (C-1), 47.5 (C-17), 48.1 (C-19), 48.2 (C-5), 49.4 (C-9), 65.7 (C-23), 71.2 (C-2), 84.0 (C-3), 122.5 (C-12), 146.8 (C-13), 185.9 (C-28);  $^1\text{H}$  and  $^{13}\text{C}$  NMR of the glycosidic part, see Table 1; ESIMS (negative-ion mode)  $m/z$  811  $[\text{M} - \text{H}]^-$ ; ESIMS-MS  $\text{MS}^2$  (811)  $m/z$  649  $[\text{M} - \text{H} - \text{Gal}]^-$ , 487  $[\text{M} - \text{H} - \text{Gal} - \text{Glc}]^-$ ; ESIMS (positive-ion mode)  $m/z$  835  $[\text{M} + \text{Na}]^+$ ; ESIMS-MS  $\text{MS}^2$  (835)  $m/z$  791  $[\text{M} + \text{Na} - \text{CO}_2]^+$ , 673  $[\text{M} + \text{Na} - \text{Gal}]^+$ , 629  $[\text{M} + \text{Na} - \text{CO}_2 - \text{Gal}]^+$ .

**Caryocaroside II-2 (4):** White powder;  $[\alpha]_D^{20} + 20.7$  (*c* 0.75, MeOH);  $^1\text{H}$  NMR of the aglycon ( $\text{CD}_3\text{OD}$ , 500 MHz), see Table 1;  $^{13}\text{C}$  NMR chemical shift values of the aglycon ( $\text{CD}_3\text{OD}$ , 125 MHz) were identical to those described for **2** ( $\pm 0.6$  ppm) except for C-28 at  $\delta$  178.0;  $^1\text{H}$  and  $^{13}\text{C}$  NMR of the glycosidic part, see Table 1; ESIMS (negative-ion mode)  $m/z$  957  $[\text{M} - \text{H}]^-$ ; ESIMS-MS  $\text{MS}^2$  (957)  $m/z$  795  $[\text{M} - \text{H} - \text{Gal}]^- = [\text{M} - \text{H} - \text{Glc}]^-$ ,  $\text{MS}^3$  (795)  $m/z$  634  $[\text{M} - \text{H} - \text{Glc} - \text{Gal}]^-$ , 471  $[\text{M} - \text{H} - 2 \text{Glc} - \text{Gal}]^-$ ; ESIMS (positive-ion mode)  $m/z$  981  $[\text{M} + \text{Na}]^+$ , 997  $[\text{M} + \text{K}]^+$ ; ESIMS-MS  $\text{MS}^2$  (981)  $m/z$  819  $[\text{M} + \text{Na} - (\text{Glc or Gal})]^+$ ,  $\text{MS}^3$  (819)  $m/z$  775  $[\text{M} + \text{Na} - \text{Glc} - \text{CO}_2]^+$ , 613  $[\text{M} + \text{Na} - \text{Glc} - \text{CO}_2 - \text{Gal}]^+$ .

**Caryocaroside III-2 (5):** White powder;  $[\alpha]_D^{20} + 28.8$  (*c* 1, MeOH);  $^1\text{H}$  NMR of the aglycon ( $\text{CD}_3\text{OD}$ , 500 MHz), see Table 1;  $^{13}\text{C}$  NMR chemical shift values of the aglycon ( $\text{CD}_3\text{OD}$ , 125 MHz) were identical to those described for **3** ( $\pm 0.6$  ppm) except for C-28 at  $\delta$  178.0;  $^1\text{H}$  and  $^{13}\text{C}$  NMR of the glycosidic part, see Table 2; ESIMS (negative-ion mode)  $m/z$  973  $[\text{M} - \text{H}]^-$ ; ESIMS-MS  $\text{MS}^2$  (973)  $m/z$  811  $[\text{M} - \text{H} - \text{Gal}]^- = [\text{M} - \text{H} - \text{Glc}]^-$ ; ESIMS (positive-ion mode)  $m/z$  1013  $[\text{M} + \text{K}]^+$ , 997  $[\text{M} + \text{Na}]^+$ ; ESIMS-MS  $\text{MS}^2$  (997)  $m/z$  835  $[\text{M} + \text{Na} - (\text{Gal or Glc})]^+$ ,  $\text{MS}^3$  (835)  $m/z$  791  $[\text{M} + \text{Na} - \text{CO}_2 - (\text{Gal or Glc})]^+$ , 673  $[\text{M} + \text{Na} - \text{Gal} - \text{Glc}]^+$ , 629  $[\text{M} + \text{Na} - \text{Gal} - \text{CO}_2 - \text{Glc}]^+$ .

**Caryocaroside II-3 (6):** White powder;  $[\alpha]_D^{20} + 15.6$  (*c* 0.42, MeOH);  $^1\text{H}$  NMR of the aglycon ( $\text{CD}_3\text{OD}$ , 500 MHz), see Table 1;  $^{13}\text{C}$  NMR chemical shift values of the aglycon ( $\text{CD}_3\text{OD}$ , 125 MHz) were identical to those described for **2** ( $\pm 0.6$  ppm);  $^1\text{H}$  and  $^{13}\text{C}$  NMR of the glycosidic part, see Table 1; ESIMS (negative-ion mode)  $m/z$  957  $[\text{M} - \text{H}]^-$ ; ESIMS-MS  $\text{MS}^2$  (957)  $m/z$  795  $[\text{M} - \text{H} - \text{Gal}]^-$ , 633  $[\text{M} - \text{H} - \text{Gal} - \text{Gal}]^-$ , 471  $[\text{M} - \text{H} - 2 \text{Gal} - \text{Glc}]^-$ .

**Caryocaroside III-3 (7):** White powder;  $[\alpha]_D^{20} + 15$  (*c* 0.42, MeOH);  $^1\text{H}$  NMR of the aglycon ( $\text{CD}_3\text{OD}$ , 500 MHz), see Table 1;  $^{13}\text{C}$  NMR chemical shift values of the aglycon ( $\text{CD}_3\text{OD}$ , 125 MHz) were identical to those described for **3** ( $\pm 0.4$  ppm);  $^1\text{H}$  and  $^{13}\text{C}$  NMR of the glycosidic part, see Table 1; ESIMS (negative-ion mode)  $m/z$  973  $[\text{M} - \text{H}]^-$ ; ESIMS-MS  $\text{MS}^2$  (973)  $m/z$  811  $[\text{M} - \text{H} - \text{Gal}]^-$ , 649  $[\text{M} - \text{H} - 2 \text{Gal}]^-$ , 487  $[\text{M} - \text{H} - 2 \text{Gal} - \text{Glc}]^-$ .

**Caryocaroside III-4 (8):** White powder;  $[\alpha]_D^{20} + 17.5$  (*c* 0.42, MeOH);  $^1\text{H}$  NMR of the aglycon ( $\text{CD}_3\text{OD}$ , 500 MHz), see Table 1;  $^{13}\text{C}$  NMR chemical shift values of the aglycon ( $\text{CD}_3\text{OD}$ , 125 MHz) were identical to those described for **3** ( $\pm 0.4$  ppm);  $^1\text{H}$  and  $^{13}\text{C}$  NMR of the glycosidic part, see Table 1; ESIMS (negative-ion mode)  $m/z$  943  $[\text{M} - \text{H}]^-$ ; ESIMS-MS  $\text{MS}^2$  (943)  $m/z$  811  $[\text{M} - \text{H} - \text{Xyl}]^-$ , 649  $[\text{M} - \text{H} - \text{Xyl} - \text{Gal}]^-$ , 487  $[\text{M} - \text{H} - \text{Xyl} - \text{Gal} - \text{Glc}]^-$ ; ESIMS (positive-ion mode)  $m/z$  967  $[\text{M} + \text{Na}]^+$ ; ESIMS-MS  $\text{MS}^2$  (967)  $m/z$  923  $[\text{M} + \text{Na} - \text{CO}_2]^+$ , 835  $[\text{M} + \text{Na} - \text{Xyl}]^+$ , 673  $[\text{M} + \text{Na} - \text{Xyl} - \text{Gal}]^+$ ,  $\text{MS}^3$  (923)  $m/z$  791  $[\text{M} + \text{Na} - \text{CO}_2 - \text{Xyl}]^+$ , 629  $[\text{M} + \text{Na} - \text{CO}_2 - \text{Xyl} - \text{Gal}]^+$ .

**Caryocaroside IV-5 (9):** White powder;  $[\alpha]_D^{20} + 26.2$  (*c* 0.58, MeOH);  $^1\text{H}$  NMR of the aglycon ( $\text{CD}_3\text{OD}$ , 500 MHz), see Table 2;  $^{13}\text{C}$  NMR of the aglycon ( $\text{CD}_3\text{OD}$ , 125 MHz)  $\delta$  18.5 (C-24), 16.8 (C-25), 17.8 (C-26), 19.1 (C-6), 24.0 (C-30), 24.0 (C-16), 24.5 (C-11), 26.4 (C-27), 28.7 (C-15), 29.9 (C-23), 31.6 (C-20), 33.4 (C-29), 33.8 (C-22), 34.0 (C-7), 34.9 (C-21), 37.7 (C-10), 39.4 (C-4), 40.6 (C-8), 42.7 (C-18), 43.0 (C-14), 44.5 (C-1), 47.2 (C-19), 47.6 (C-17), 49.3 (C-9), 57.0 (C-5), 71.2 (C-2), 91.1 (C-3), 123.9 (C-12), 145.2 (C-13), 181.9 (C-28);  $^1\text{H}$  and  $^{13}\text{C}$  NMR of the glycosidic part, see Table 2; ESIMS (negative-ion mode)  $m/z$  647  $[\text{M} - \text{H}]^-$ ; ESIMS-MS  $\text{MS}^2$  (647)  $m/z$  471  $[\text{M} - \text{H} - \text{GlcA}]^-$ ; ESIMS (positive-ion mode)  $m/z$  671  $[\text{M} + \text{Na}]^+$ , 687  $[\text{M} + \text{K}]^+$ ; ESIMS-MS  $\text{MS}^2$  (671)  $m/z$  495  $[\text{M} + \text{Na} - \text{GlcA}]^+$ , 451  $[\text{M} + \text{Na} - \text{GlcA} - \text{CO}_2]^+$ .

**Caryocaroside III-5 (10):** White powder;  $[\alpha]_D^{20} + 16.8$  (*c* 0.25, MeOH);  $^1\text{H}$  NMR of the aglycon ( $\text{CD}_3\text{OD}$ , 500 MHz), see Table 2;  $^{13}\text{C}$  NMR chemical shift values of the aglycon ( $\text{CD}_3\text{OD}$ , 125 MHz) were identical to those described for **5** ( $\pm 0.2$  ppm);  $^1\text{H}$  and  $^{13}\text{C}$  NMR of the glycosidic part, see Table 2; ESIMS (negative-ion mode)  $m/z$  663  $[\text{M} - \text{H}]^-$ ; ESIMS-MS  $\text{MS}^2$  (663)  $m/z$  487  $[\text{M} - \text{H} - \text{GlcA}]^-$ ; ESIMS (positive-ion mode)  $m/z$  709  $[\text{M} - \text{H} + 2 \text{Na}]^+$ , 687  $[\text{M} + \text{Na}]^+$ ; ESIMS-MS  $\text{MS}^2$  (709)  $m/z$  691  $[\text{M} - \text{H} + 2 \text{Na} - \text{H}_2\text{O}]^+$ , 533  $[\text{M} - \text{H} + 2 \text{Na} - \text{GlcA}]^+$ .

**Caryocaroside IV-6 (11):** White powder;  $[\alpha]_D^{20} + 12$  (*c* 0.17, MeOH);  $^1\text{H}$  NMR of the aglycon ( $\text{CD}_3\text{OD}$ , 500 MHz), see Table 2;  $^{13}\text{C}$  NMR chemical shift values of the aglycon ( $\text{CD}_3\text{OD}$ , 125 MHz) were identical to those described for **9** ( $\pm 0.4$  ppm);  $^1\text{H}$  and  $^{13}\text{C}$  NMR of the glycosidic part, see Table 2; ESIMS (negative-ion mode)  $m/z$  661  $[\text{M} - \text{H}]^-$ ; ESIMS-MS  $\text{MS}^2$  (661)  $m/z$  647  $[\text{M} - \text{H} - \text{CH}_3]^-$ , 471  $[\text{M} - \text{H} - (\text{CH}_3\text{GlcA})]^-$ ; ESIMS (positive-ion mode)  $m/z$  701  $[\text{M} + \text{K}]^+$ , 685  $[\text{M} + \text{Na}]^+$ ; ESIMS-MS  $\text{MS}^2$  (685)  $m/z$  495  $[\text{M} + \text{Na} - \text{Me} - \text{GlcA}]^+$ .

**Caryocaroside IV-7 (12):** White powder;  $[\alpha]_D^{20} + 8.4$  (*c* 0.17, MeOH);  $^1\text{H}$  NMR of the aglycon ( $\text{DMSO-}d_6$ , 500 MHz), see Table 2;  $^{13}\text{C}$  NMR of the aglycon ( $\text{DMSO-}d_6$ , 125 MHz)  $\delta$  17.2 (C-25), 18.7 (C-26), 19.5 (C-24), 19.2 (C-6), 24.9 (C-30), 23.6 (C-16), 24.5 (C-11), 27.0 (C-27), 28.8 (C-15), 30.6 (C-23), 31.9 (C-20), 34.4 (C-29), 34.0 (C-22), 34.0 (C-7), 35.2 (C-21), 37.7 (C-10), 39.3 (C-4), 40.3 (C-8), 42.6 (C-18), 43.0 (C-14), 44.8 (C-1), 47.7 (C-19), 47.7 (C-17), 49.0 (C-9), 52.0 (C-5), 69.6 (C-2), 90.0 (C-3), 123.0 (C-12), 146.0 (C-13), 179.7 (C-28);  $^1\text{H}$  and  $^{13}\text{C}$  NMR of the glycosidic part, see Table 2; ESIMS (negative-ion mode)  $m/z$  809  $[\text{M} - \text{H}]^-$ ; ESIMS-MS  $\text{MS}^2$  (809)  $m/z$  647  $[\text{M} - \text{H} - \text{Gal}]^-$ , 471  $[\text{M} - \text{H} - \text{Gal} - \text{GlcA}]^-$ ; ESIMS (positive-ion mode)  $m/z$  855  $[\text{M} - \text{H} + 2\text{Na}]^+$ ; ESIMS-MS  $\text{MS}^2$  (855)  $m/z$  693  $[\text{M} - \text{H}$

+ 2 Na – Gal]<sup>+</sup>, 517 [M – H + 2Na – Gal – GlcA]<sup>+</sup>, MS<sup>3</sup> (693) *m/z* 675 [M – H + 2 Na – Gal – H<sub>2</sub>O]<sup>+</sup>, 517 [M – H + 2 Na – Gal – GlcA]<sup>+</sup>.

**Caryocaroside II-7 (13):** White powder; [ $\alpha$ ]<sub>D</sub><sup>20</sup> + 11.4 (*c* 0.44, MeOH); <sup>1</sup>H NMR of the aglycon (DMSO-*d*<sub>6</sub>, 500 MHz), see Table 2; <sup>13</sup>C NMR of the aglycon (DMSO-*d*<sub>6</sub>, 125 MHz)  $\delta$  14.3 (C-24), 17.0 (C-25), 18.4 (C-26), 18.7 (C-6), 24.1 (C-16), 24.7 (C-30), 24.4 (C-11), 26.4 (C-2), 27.0 (C-27), 28.7 (C-15), 31.8 (C-20), 33.6 (C-7), 33.5 (C-22), 34.2 (C-29), 34.8 (C-21), 37.5 (C-10), 39.5 (C-1), 40.4 (C-8), 42.4 (C-18), 42.8 (C-14), 43.8 (C-4), 47.0 (C-17), 47.2 (C-19), 47.5 (C-5), 48.6 (C-9), 63.9 (C-23), 79.6 (C-3), 123.0 (C-12), 145.4 (C-13), 182.0 (C-28); <sup>1</sup>H and <sup>13</sup>C NMR of the glycosidic part, see Table 2; ESIMS (negative-ion mode) *m/z* 809 [M – H]<sup>-</sup>; ESIMS-MS MS<sup>2</sup> (809) *m/z* 471 [M – H – Gal – GlcA]<sup>-</sup>; ESIMS (positive-ion mode) *m/z* 847 [M + K]<sup>+</sup>.

**Caryocaroside III-7 (14):** White powder; [ $\alpha$ ]<sub>D</sub><sup>20</sup> + 26.2 (*c* 0.67, MeOH); <sup>1</sup>H NMR of the aglycon (CD<sub>3</sub>OD, 500 MHz), see Table 2; <sup>13</sup>C NMR chemical shift values of the aglycon (CD<sub>3</sub>OD, 125 MHz) were identical to those described for **3** ( $\pm$  0.6 ppm); <sup>1</sup>H and <sup>13</sup>C NMR of the glycosidic part, see Table 2; ESIMS (negative-ion mode) *m/z* 825 [M – H]<sup>-</sup>; ESIMS-MS MS<sup>2</sup> (825) *m/z* 487 [M – H – Gal – GlcA]<sup>-</sup>; ESIMS (positive-ion mode) *m/z* 871 [M – H + 2 Na]<sup>+</sup>, 849 [M + Na]<sup>+</sup>; ESIMS-MS MS<sup>2</sup> (871) *m/z* 709 [M – H + 2 Na – Gal]<sup>+</sup>, 533 [M – H + 2 Na – Gal – GlcA]<sup>+</sup>, MS<sup>3</sup> (709) *m/z* 691 [M – H + 2 Na – Gal – H<sub>2</sub>O]<sup>+</sup>, 533 [M – H + 2 Na – Gal – GlcA]<sup>+</sup>.

**Caryocaroside IV-8 (15):** White powder; [ $\alpha$ ]<sub>D</sub><sup>20</sup> + 30 (*c* 1, MeOH); <sup>1</sup>H NMR of the aglycon (CD<sub>3</sub>OD, 500 MHz), see Table 2; <sup>13</sup>C NMR chemical shift values of the aglycon (CD<sub>3</sub>OD, 125 MHz) were identical to those described for **9** ( $\pm$  0.2 ppm). <sup>1</sup>H and <sup>13</sup>C NMR of

the glycosidic part, see Table 2; ESIMS (negative-ion mode)  $m/z$  823  $[M - H]^-$ ; ESIMS-MS  $MS^2$  (8423)  $m/z$  791  $[M - H - CH_3OH]^-$ , 601  $[M - H - Gal - (COOCH_3)]^-$ , 471  $[M - H - Gal - CH_3GlcA]^-$ ; ESIMS (positive-ion mode)  $m/z$  863  $[M + K]^+$ , 847  $[M + Na]^+$ ; ESIMS-MS  $MS^2$  (847)  $m/z$  685  $[M + Na - Gal]^+$ , 641  $[M + Na - Gal - CO_2]^+$ .

**Caryocaroside IV-9 (16):** White powder;  $[\alpha]^{20}_D + 12$  ( $c$  0.5, MeOH);  $^1H$  NMR of the aglycon ( $CD_3OD$ , 500 MHz), see Table 3;  $^{13}C$  NMR chemical shift values of the aglycon ( $CD_3OD$ , 125 MHz) were identical to those described for **9** ( $\pm 0.6$  ppm) except for C-28 at  $\delta$  178.0;  $^1H$  and  $^{13}C$  NMR of the glycosidic part : see Table 3; ESIMS (negative-ion mode)  $m/z$  971  $[M - H]^-$ ; ESIMS-MS  $MS^2$  (971)  $m/z$  809  $[M - H - Gal]^- = [M - H - Glc]^-$ ,  $MS^3$  (809)  $m/z$  647  $[M - H - Gal - Glc]^-$ , 471  $[M - H - Gal - Glc - GlcA]^-$ ; ESIMS (positive-ion mode)  $m/z$  1017  $[M - H + 2Na]^+$ ; ESIMS-MS  $MS^2$  (1017)  $m/z$  855  $[M - H + 2 Na - (Glc or Gal)]^+$ ,  $MS^3$  (855)  $m/z$  693  $[M - H + 2 Na - Glc - Gal]^+$ , 517  $[aglycon - H + 2 Na]^+$ .

**Caryocaroside II-9 (17):** White powder;  $[\alpha]^{20}_D + 11.9$  ( $c$  1, MeOH);  $^1H$  NMR of the aglycon ( $CD_3OD$ , 500 MHz), see Table 3;  $^{13}C$  NMR chemical shift values of the aglycon ( $CD_3OD$ , 125 MHz) were identical to those described for **4** ( $\pm 0.6$  ppm);  $^1H$  and  $^{13}C$  NMR of the glycosidic part : see Table 3; ESIMS (negative-ion mode)  $m/z$  971  $[M - H]^-$ ; ESIMS-MS  $MS^2$  (971)  $m/z$  809  $[M - H - Gal]^- = [M - H - Glc]^-$ ,  $MS^3$  (809)  $m/z$  647  $[M - H - Gal - Glc]^-$ , 471  $[M - H - Gal - Glc - GlcA]^-$ ; ESIMS (positive-ion mode)  $m/z$  1009  $[M - H + K]^+$ ; ESIMS-MS  $MS^2$  (1009)  $m/z$  847  $[M - H + K - (Glc or Gal)]^+$ , 685  $[M - H + K - Glc - Gal]^+$ ,  $MS^3$  (847)  $m/z$  803  $[M - H + K - (Glc or Gal) - CO_2]^+$ , 685  $[M - H + K - Glc - Gal]^+$ , 641  $[M - H + K - Glc - Gal - CO_2]^+$ .

**Caryocaroside III-9 (18):** White powder;  $[\alpha]^{20}_D + 15.6$  ( $c$  1, MeOH);  $^1H$  NMR of the aglycon ( $CD_3OD$ , 500 MHz), see Table 3;  $^{13}C$  NMR chemical shift values of the aglycon



(CD<sub>3</sub>OD, 125 MHz) were identical to those described for **5** ( $\pm 0.4$  ppm); <sup>1</sup>H and <sup>13</sup>C NMR of the glycosidic part : see Table 3; ESIMS (negative-ion mode)  $m/z$  987 [M – H]<sup>–</sup>; ESIMS-MS MS<sup>2</sup> (987)  $m/z$  825 [M – H – Gal]<sup>–</sup> = [M – H – Glc]<sup>–</sup>, MS<sup>3</sup> (825)  $m/z$  663 [M – H – Gal – Glc]<sup>–</sup>, 487 [M – H – Gal – Gal – GlcA]<sup>–</sup>; ESIMS (positive-ion mode)  $m/z$  1027 [M + K]<sup>+</sup>, 1011 [M + Na]<sup>+</sup>; ESIMS-MS MS<sup>2</sup> (1027)  $m/z$  863 [M + K – (Glc or Gal)]<sup>+</sup>, MS<sup>3</sup> (863)  $m/z$  819 [M + K – (Glc or Gal) – CO<sub>2</sub>]<sup>+</sup>, 701 [M + K – Glc – Gal]<sup>+</sup>, 657 [M + K – Glc – Gal – CO<sub>2</sub>]<sup>+</sup>.

**Caryocaroside II-10 (19):** White powder;  $[\alpha]_D^{20} + 10.8$  ( $c$  0.17, MeOH); <sup>1</sup>H NMR of the aglycon (CD<sub>3</sub>OD, 500 MHz), see Table 3; <sup>13</sup>C NMR chemical shift values of the aglycon (CD<sub>3</sub>OD, 125 MHz) were identical to those described for **5** ( $\pm 0.4$  ppm); <sup>1</sup>H and <sup>13</sup>C NMR of the glycosidic part : see Table 3; ESIMS (positive-ion mode)  $m/z$  1009 [M + Na]<sup>+</sup>, 847 [M + Na – (Glc or Gal)]<sup>+</sup>; ESIMS-MS MS<sup>2</sup> (1009)  $m/z$  979 [M + Na – CH<sub>3</sub>OH]<sup>+</sup>, 847 [M + Na – (Glc or Gal)]<sup>+</sup>, 685 [M + Na – Glc – Gal]<sup>+</sup>, MS<sup>3</sup> (847)  $m/z$  685 [M + Na – Glc – Gal]<sup>+</sup>, 641 [M + Na – Glc – Gal – CO<sub>2</sub>]<sup>+</sup>.

**Caryocaroside IV-11 (20):** White powder;  $[\alpha]_D^{20} + 12$  ( $c$  0.29, MeOH); <sup>1</sup>H NMR of the aglycon (CD<sub>3</sub>OD, 500 MHz), see Table 3; <sup>13</sup>C NMR chemical shift values of the aglycon (CD<sub>3</sub>OD, 125 MHz) were identical to those described for **9** ( $\pm 0.4$  ppm); <sup>1</sup>H and <sup>13</sup>C NMR of the glycosidic part : see Table 3; ESIMS (negative-ion mode)  $m/z$  971 [M – H]<sup>–</sup>; ESIMS-MS MS<sup>2</sup> (971)  $m/z$  809 [M – H – Gal]<sup>–</sup>; 647 [M – H – Gal – Gal]<sup>–</sup>, 471 [M – H – 2 Gal – GlcA]<sup>–</sup>.

**Caryocaroside II-11 (21):** White powder;  $[\alpha]_D^{20} + 11.5$  ( $c$  0.42, MeOH); <sup>1</sup>H NMR of the aglycon (CD<sub>3</sub>OD, 500 MHz), see Table 3; <sup>13</sup>C NMR chemical shift values of the aglycon (CD<sub>3</sub>OD, 125 MHz) were identical to those described for **2** ( $\pm 0.4$  ppm); <sup>1</sup>H and <sup>13</sup>C NMR of

the glycosidic part : see Table 3; ESIMS (negative-ion mode)  $m/z$  971 [M – H]<sup>-</sup>; ESIMS-MS MS<sup>2</sup> (971)  $m/z$  809 [M – H – Gal]<sup>-</sup>, 647 [M – H – 2 Gal]<sup>-</sup>, 471 [M – H – 2 Gal – GlcA]<sup>-</sup>.


**Acknowledgment.** The authors are very grateful to M.F. Prevost of the Botany Department, Research Institute for the Development (IRD), Cayena (Guyana), for help with the identification of the plant. We thank Isabelle Pouny, ISTMT Research Institut Pierre Fabre, for recording some ESIMS spectra and Karen Plé for revising the English of the manuscript.

## References and Notes

- (1) Prance, G. T. *Opera-Botanica* **1987**, *92*, 179–184.
- (2) Dresen, H.; Prasad, R. B. N.; Gülz, P.-G. *Z. Naturforsch. C* **1989**, *44*, 739–742.
- (3) Marx, F.; Andrade, E. H.-A.; Maia, J. G. *Z. Lebensm. Unters.-Forsch. A* **1997**, *204*, 442–444.
- (4) Kelly, D.; Bessiere, J.; Crimmins, J.; and Renard, S. *Söfw J.* **2003**, *129*, 2–6.
- (5) Passos, X. S.; Castro, A. C. M.; Pires, J. S.; Garcia, A. C.- F.; Campos, F.-C; Fernandes, O. F. L.; Paula, J. R.; Ferreira, H. D.; Santos, S. C.; Ferri, P. H.; Silva, M. D. R. R. *Pharm. Biol.* **2003**, *41*, 319–324.
- (6) Kawanishi, K.; Raffauf, R.-F. *J. Nat. Prod.* **1986**, *49*, 1167–1168.
- (7) Grenand, P.; Moretti, C. ; Jacquemin, H.; Prévost, M. F. In *Pharmacopées Traditionnelles en Guyane, Créoles, Wayasi, Palikur*; IRD Editions: Paris, 2004, 293–298.
- (8) Voutquenne, L.; Lavaud, C.; Massiot, G.; Le Men-Olivier, L. *Pharm. Biol.* **2002**, *40*, 253–262.
- (9) Chwalek, M.; Ple, K.; Voutquenne-Nazabadioko, L. *Chem. Pharm. Bull.* **2004**, *52*, 965–971.

- (10) Mahato S. B.; Kundu, A. P. *Phytochemistry* **1994**, *35*, 1319–1324.
- (11) Tan, N.; Zhou, J.; Zhao, S. *Phytochemistry* **1999**, *52*, 153–192.
- (12) (a) Thiilborg, S. T.; Christensen, S. B.; Cornett, C.; Olsen, C. E.; Lemmich, E. *Phytochemistry* **1994**, *36*, 753–759. (b) Bialy, Z.; Jurzysta, M.; Mella, M.; Tava, A. *J. Agric. Food Chem.* **2004**, *52*, 1095–1099.
- (13) Lavaud, C.; Voutquenne, L.; Massiot, G.; Le Men-Olivier, L.; Das, B. C.; Laprévotte, O.; Serani, L.; Delaude, C.; Becchi, M. *Phytochemistry* **1998**, *47*, 441–449.
- (14) (a) Vidal-Ollivier, E.; Balansard, G.; Faure, R.; Babadjamian, A. *J. Nat. Prod.* **1989**, *52*, 1156–1159. (b) Yoshikawa, M.; Murakami, T.; Kishi, A.; Kageura, T.; Matsuda, H. *Chem. Pharm. Bull.* **2001**, *49*, 863–870.
- (15) (a) Chemli, R.; Babadjamian, A.; Faure, R.; Boukef, K.; Balansard, G.; Vidal, E. *Phytochemistry* **1987**, *26*, 1785–1788. (b) Pizza, C.; Zhong-Liang, Z.; De Tommasi, N. *J. Nat. Prod.* **1987**, *50*, 927–931.
- (16) Abdel-Sattar, E. *Pharm. Biol.* **2001**, *39*, 440–444.
- (17) Hu, M.; Ogawa, K.; Sashida, Y.; Xiao, P. G. *Phytochemistry* **1995**, *39*, 179–184.
- (18) Kizu, H.; Kitayama, S.; Nakatani, F.; Tomimori, T.; Namba, T. *Chem. Pharm. Bull.* **1985**, *33*, 3324–3329.
- (19) Mshvildadze, V.; Elias, R.; Faure, R.; Debrauwer, L.; Dekanosidze, G.; Kemertelidze, E.; Balansard, G. *Chem. Pharm. Bull.* **2001**, *49*, 752–754.
- (20) Domon, B.; Hostettmann, K. *Helv. Chim. Acta* **1983**, *66*, 422–428.
- (21) Lavaud, C.; Beauvière, S.; Massiot, G.; Le Men-Olivier, L.; Bourdy, G. *Phytochemistry* **1996**, *43*, 189–194.
- (22) Paphassarang, S.; Raynaud, J.; Lussignol, M.; Becchi, M. *Phytochemistry* **1989**, *28*, 1539–1541.

- (23) Marquina, S.; Maldonado, N.; Garduño-Ramirez, M. L.; Aranda, E.; Villarreal, M. L.; Navarro, V.; Bye, R.; Delgado, G.; Alvarez, L. *Phytochemistry* **2001**, *56*, 93–97.
- (24) Zhu, N.; Sheng, S.; Sang, S.; Jhoo, J. W.; Bai, N.; Karwe, M. V.; Rosen, R. T.; Ho, C. T. *J. Agric. Food Chem.* **2002**, *50*, 865–867.
- (25) Ridout, C. L.; Price, K. R.; Dijoux, M.-G.; Lavaud, C. *J. Agric. Food Chem.* **1994**, *42*, 279–282.
- (26) Sakai, S.; Katsumata, M.; Satoh, Y.; Nagasao, M.; Miyakoshi, M.; Ida, Y.; Shoji, J. *Phytochemistry* **1994**, *35*, 1319–1324.
- (27) Mshvildadze, V.; Favel, A.; Delmas F.; Elias, R.; Faure, R.; Decanosidze, G.; Kemertelidze, E.; Balansard, G. *Pharmazie* **2000**, *55*, 325–326.
- (28) Marston, A.; Gafner, F.; Dossaji, S. F.; Hostettmann, K. *Phytochemistry* **1988**, *27*, 1325–1326.
- (29) Abdel-Kader, M. S.; Bahler, B. D.; Malone, S.; Werkhoven, M. C. M.; Wisse, J. H.; Neddermann, K. M.; Bursuker, I.; Kingston, D. G. I. *J. Nat. Prod.* **2000**, *63*, 923–926.
- (30) (a) Takechi, M.; Uno C.; Tanaka, Y. *Phytochemistry* **1996**, *41*, 121–123. (b) Ullah, N.; Seebacher W.; Weis, R.; Jerenitsch J.; Rauchensteiner, K.; Haslinger, E. *Monatsh. Chem.* **2000**, *131*, 787–794.
- (31) Herbar de Guyane available at <http://www.cayenne.ird.fr/aublet2>.



	<b>R<sub>1</sub></b>	<b>R<sub>2</sub></b>		<b>R<sub>3</sub></b>	<b>R<sub>4</sub></b>
<b>1</b>	H	COOH		Gal-(1→3)-Glc-	H
<b>2</b>	H	CH <sub>2</sub> OH		Gal-(1→3)-Glc-	H
<b>3</b>	OH	CH <sub>2</sub> OH		Gal-(1→3)-Glc-	H
<b>4</b>	H	CH <sub>2</sub> OH		Gal-(1→3)-Glc-	Glc
<b>5</b>	OH	CH <sub>2</sub> OH		Gal-(1→3)-Glc-	Glc
<b>6</b>	H	CH <sub>2</sub> OH	Gal-(1→3)-Gal-(1→3)-Glc-		H
<b>7</b>	OH	CH <sub>2</sub> OH	Gal-(1→3)-Gal-(1→3)-Glc-		H
<b>8</b>	OH	CH <sub>2</sub> OH	Xyl-(1→3)-Gal-(1→3)-Glc-		H
<b>9</b>	OH	CH <sub>3</sub>		GlcA-	H
<b>10</b>	OH	CH <sub>2</sub> OH		GlcA-	H
<b>11</b>	OH	CH <sub>3</sub>	6-O-methyl-GlcA-		H
<b>12</b>	OH	CH <sub>3</sub>	Gal-(1→3)-GlcA-		H
<b>13</b>	H	CH <sub>2</sub> OH	Gal-(1→3)-GlcA-		H
<b>14</b>	OH	CH <sub>2</sub> OH	Gal-(1→3)-GlcA-		H
<b>15</b>	OH	CH <sub>3</sub>	Gal-(1→3)-6-O-methyl-GlcA-		H
<b>16</b>	OH	CH <sub>3</sub>	Gal-(1→3)-GlcA-		Glc
<b>17</b>	H	CH <sub>2</sub> OH	Gal-(1→3)-GlcA-		Glc
<b>18</b>	OH	CH <sub>2</sub> OH	Gal-(1→3)-GlcA-		Glc
<b>19</b>	H	CH <sub>2</sub> OH	Gal-(1→3)-6-O-methyl-GlcA-		Glc
<b>20</b>	OH	CH <sub>3</sub>	Gal-(1→3)-Gal-(1→3)-GlcA-		H
<b>21</b>	H	CH <sub>2</sub> OH	Gal-(1→3)-Gal-(1→3)-GlcA-		H
<b>24</b>	H	CH <sub>3</sub>	Gal-(1→3)-GlcA-		H
<b>28</b>	H	CH <sub>2</sub> OH		GlcA-	H

**Table 1.** <sup>1</sup>H and <sup>13</sup>C NMR Data of Compounds **1–8** (CD<sub>3</sub>OD).

	<b>1</b>		<b>2</b>		<b>3</b>		<b>4</b>		<b>5</b>		<b>6</b>		<b>7</b>		<b>8</b>	
	<sup>1</sup> H	<sup>13</sup> C	<sup>1</sup> H	<sup>13</sup> C	<sup>1</sup> H	<sup>13</sup> C	<sup>1</sup> H	<sup>13</sup> C	<sup>1</sup> H	<sup>13</sup> C	<sup>1</sup> H	<sup>13</sup> C	<sup>1</sup> H	<sup>13</sup> C	<sup>1</sup> H	<sup>13</sup> C
aglycon																
2	1.71 (m)	26.4	1.77 (m)	26.2	4.34 (q, 3.1)	71.2	1.78 (m)	26.3	4.32 (q, 2.9)	70.7	1.77 (td, 14, 4)	26.4	4.34 (q, 3.5)	71.1	4.34 (q, 3.5)	71.2
2	1.99 (m)	-	1.94 (dq, 13.5, 4.4)	-	-	-	1.96 (dq, 14.1, 3.8)	-	-	-	1.95 (m)	-	-	-	-	-
3	4.11 (dd, 11.8, 4.5)	85.8	3.67 (d, 8.5, 4.7)	83.7	3.64 (d, 3.8)	84.0	3.67 (dd, 12.2, 4)	83.5	3.52 (d, 3.6)	84.1	3.65 (m)	83.6	3.64 (d, 3.5)	84.0	3.64 (d, 3.5)	84.0
12	5.27 (t, 3.6)	123.4	5.26 (t, 3.6)	123.4	5.25 (t, 3.6)	122.5	5.27 (t, 3.7)	123.8	5.30 (t, 3.4)	123.5	5.27 (t, 3.3)	122.3	5.25 (t, 3.5)	122.4	5.25 (t, 3.6)	122.4
18	2.87 (dd, 14.2, 4)	42.7	2.86 (dd, 13.7, 4.2)	42.6	2.93 (dd, 13.7, 4.5)	43.5	2.88 (dd, 13.1, 3.4)	42.6	2.87 (dd, 13.7, 4.2)	42.3	2.93 (dd, 13.6, 3.2)	43.5	2.93 (dd, 13.1, 3.5)	43.5	2.94 (dm, 13)	43.5
23	-	181.9	3.31 (d, 11.4)	65.1	3.26 (d, 11..3)	65.7	3.27 (d, 11.8)	65.0	3.25 (d, 11.3)	65.8	3.31 (d, 11.1)	65.0	3.26 (d, 12.2)	65.7	3.26 (d, 11.5)	65.8
23	-	-	3.65 (d, 11.4)	-	3.64 (d, 11.5)	-	3.66 (d, 11.8)	-	3.61 (d, 11.4)	-	3.63 (d, 11.1)	-	3.64 (d, 12)	-	3.64 (d, 11.5)	-
24	1.17 (s)	12.0	0.73 (s)	13.4	0.96 (s)	14.7	0.73 (s)	13.4	0.95 (s)	14.5	0.73 (s)	13.4	0.96 (s)	14.7	0.96 (s)	14.7
25	1.00 (s)	16.2	1.00 (s)	16.4	1.30 (s)	17.5	1.01 (s)	16.5	1.28 (s)	17.4	1.00 (s)	16.5	1.30 (s)	17.5	1.29 (s)	17.5
26	0.83 (s)	17.6	0.83 (s)	17.7	0.91 (s)	18.3	0.82 (s)	17.8	0.81 (s)	17.6	0.92 (s)	18.3	0.92 (s)	18.7	0.92 (s)	18.3
27	1.19 (s)	26.4	1.19 (s)	26.5	1.17 (s)	26.6	1.19 (s)	26.3	1.16 (s)	26.4	1.17 (s)	26.6	1.17 (s)	26.6	1.17 (s)	26.6
28	-	182.0	-	182.2	-	185.9	-	178.1	-	178.0	-	186.0	-	186.3	-	180.2
29	0.93 (s)	33.6	0.92 (s)	33.6	0.90 (s)	34.0	0.96 (s)	33.5	0.91 (s)	33.5	0.90 (s)	34.0	0.90 (s)	34.0	0.90 (s)	34.0
30	0.96 (s)	24.0	0.96 (s)	24.0	0.98 (s)	24.4	0.96 (s)	24.0	0.94 (s)	23.9	0.98 (s)	24.5	0.98 (s)	24.6	0.98 (s)	24.4
<i>β</i> -D-glucose (at C-3)																
1'	4.37 (d, 7.7)	104.9	4.48 (d, 7.9)	105.1	4.52 (d, 7.8)	105.2	4.48 (d, 7.9)	105.3	4.49 (d, 7.9)	104.6	4.48 (d, 7.9)	105.3	4.51 (d, 7.9)	105.2	4.51 (d, 7.5)	105.2
2'	3.33 (t, 8.4)	74.3	3.40 (dd, 8.9, 7.9)	74.7	3.50 (dd, 9, 7.8)	74.7	3.39 (dd, 8.9, 7.9)	74.8	3.50 (t, 8.4)	74.2	3.40 (dd, 8.9, 7.9)	74.7	3.51 (dd, 8.9, 7.9)	74.6	3.50 (dd, 9, 7.5)	74.8
3'	3.35 (t, 9)	87.3	3.57 (t, 8.9)	88.0	3.60 (t, 8.9)	87.8	3.57 (t, 9)	88.1	3.58 (t, 8.3)	87.8	3.59 (t, 8.9)	87.9	3.62 (t, 8.9)	87.8	3.61 (t, 9)	87.7
4'	3.42 (t, 8.8)	70.0	3.45 (dd, 9.6, 8.9)	69.8	3.51 (dd, 9.6, 8.9)	69.5	3.45 (dd, 9.6, 9)	69.9	3.53 (t, 8.7)	69.2	3.45 (dd, 9.6, 8.9)	69.8	3.52 (t, 9)	69.5	3.52 (t, 8.9)	69.5
5'	3.28 (ddd, 10, 5.1, 2)	77.4	3.34 (ddd, 9.6, 5.1, 2.4)	77.1	3.35 (ddd, 9.6, 4.6, 2.3)	77.3	3.33 (m)	77.3	3.33 (m)	76.9	3.33 (m)	77.3	3.35 (m)	77.3	3.35 (m)	77.3
6'a	3.70 (dd, 11.6, 4.7)	62.6	3.72 (dd, 12, 5.1)	62.5	3.74 (dd, 12, 4.6)	62.2	3.72 (dd, 11.5, 4.3)	62.6	3.74 (dd, 12, 4.5)	61.9	3.72 (m)	62.6	3.75 (m)	62.2	3.74 (dd, 12, 4.5)	62.2
6'b	3.87 (dd, 12, 2.3)	-	3.87 (dd, 12, 2.4)	-	3.83 (dd, 12, 2.3)	-	3.87 (dd, 11.5, 2)	-	3.83 (dd, 12, 3.6)	-	3.87 (dd, 12, 2)	-	3.84 (m)	-	3.83 (dd, 12, 2)	-
<i>β</i> -D-galactose (at C-3')																
1''	4.52 (d, 7.6)	105.7	4.52 (d, 7.7)	105.5	4.55 (d, 7.7)	105.8	4.52 (d, 7.7)	105.7	4.50 (d, 7.8)	105.4	4.62 (d, 7.8)	105.3	4.63 (d, 7.8)	105.3	4.62 (d, 7.5)	106.3
2''	3.60 (dd, 9.8, 7.7)	73.1	3.63 (dd, 9.7, 7.7)	72.9	3.64 (dd, 9.6, 7.8)	73.0	3.63 (dd, 9.6, 7.7)	73.0	3.64 (dd, 9.6, 7.8)	72.7	3.80 (dd, 9.4, 7.6)	72.2	3.81 (dd, 9, 8)	72.1	3.80 (t, 8)	72.2
3''	3.51 (dd, 9.8, 3.6)	74.7	3.54 (dd, 9.7, 3.3)	74.6	3.54 (dd, 9.6, 3.4)	74.7	3.53 (dd, 9.6, 3.3)	74.7	3.54 (dd, 9.6, 3.2)	74.4	3.68 (dd, 9.4, 2.9)	84.5	3.69 (dd, 9.3, 3)	84.6	3.67 (dd, 7, 3)	84.1
4''	3.82 (d, 3.5)	70.3	3.83 (dd, 3.3, 0.1)	70.2	3.83 (d, 3.4)	70.3	3.82 (d, 3.3)	70.3	3.83 (d, 3.2)	69.9	4.13 (d, 3)	70.2	4.15 (d, 3.3)	69.8	4.04 (dd, 3, 0.5)	69.8
5''	3.56 (m)	77.1	3.60 (ddd, 9.6, 5.8, 1)	77.0	3.60 (dd, 7.7, 4.4)	77.2	3.59 (ddd, 7.7, 3.3, 1)	77.2	3.58 (m)	76.8	3.62 (m)	76.7	3.63 (m)	76.7	3.61 (m)	76.7
6''a	3.70 (dd, 11.6, 4.6)	62.6	3.72 (dd, 11.6, 4.5)	62.5	3.71 (dd, 11.5, 4.4)	62.6	3.71 (dd, 11.4, 3.3)	62.5	3.71 (dd, 11.5, 4)	62.3	3.71 (dd, 11.5, 2.7)	62.6	3.74 (dd, 11.5, 2.7)	62.6	3.70 (dd, 11, 4)	62.5
6''b	3.79 (dd, 11.6, 7.6)	-	3.81 (dd, 11.6, 7.8)	-	3.80 (dd, 11.5, 7.7)	-	3.80 (dd, 11.4, 7.7)	-	3.81 (dd, 11.5, 7.8)	-	3.81 (dd, 11.5, 7.4)	-	3.80 (dd, 11.5, 5)	-	3.80 (dd, 11, 8)	-
sugar (at C-28 or C-3'')							<i>β</i> -D-glucose (at C-28)		<i>β</i> -D-glucose (at C-28)		<i>β</i> -D-galactose (at C-3'')		<i>β</i> -D-galactose (at C-3'')		<i>β</i> -D-xylose (at C-3'')	
1'''							5.40 (d, 8.1)	95.7	5.40 (d, 8.1)	95.4	4.52 (d, 7.6)	106.3	4.53 (d, 7.8)	106.9	4.53 (d, 7)	105.2
2'''							3.34 (dd, 9.3, 8.1)	73.9	3.35 (t, 8.5)	73.5	3.64 (m)	73.0	3.64 (m)	72.9	3.32 (t, 9)	75.2
3'''							3.43 (t, 9.3)	78.3	3.45 (dd, 9.1, 8.5)	77.9	3.52 (dd, 9.7, 3.3)	74.6	3.53 (dd, 9.7, 3.6)	74.8	3.36 (t, 9)	77.6
4'''							3.38 (t, 9.3)	71.1	3.40 (t, 9.3)	70.8	3.82 (d, 3.4)	69.8	3.85 (d, 3)	70.2	3.52 (m)	71.2
5'''/5'''a							3.37 (m)	78.7	3.37 (m)	78.2	3.56 (m)	76.8	3.57 (m)	76.8	3.24 (t, 11.5)	66.9
5'''b															3.89 (dd, 11.5, 5)	-
6'''a							3.70 (dd, 11.4, 3)	62.4	3.71 (dd, 11.6, 3.4)	62.2	3.75 (m)	62.5	3.74 (m)	62.6		
6'''b							3.84 (dd, 11.4, 2)	-	3.83 (dd, 11.6, 2.5)	-	3.75 (m)	-	3.74 (m)	-		

**Table 2.** <sup>1</sup>H and <sup>13</sup>C NMR Data of Compounds **9** – **11**, **14** – **15** (CD<sub>3</sub>OD), and **12** – **13** (DMSO-*d*<sub>6</sub>).

	<b>9</b>		<b>10</b>		<b>11</b>		<b>12</b>		<b>13</b>		<b>14</b>		<b>15</b>	
	<sup>1</sup> H	<sup>13</sup> C	<sup>1</sup> H	<sup>13</sup> C	<sup>1</sup> H	<sup>13</sup> C	<sup>1</sup> H	<sup>13</sup> C	<sup>1</sup> H	<sup>13</sup> C	<sup>1</sup> H	<sup>13</sup> C	<sup>1</sup> H	<sup>13</sup> C
Aglycon														
2	4.22 (q, 3.3)	71.2	4.32 (q, 3.4)	71.1	4.17 (q, 2.9)	71.2	4.08 (q, 3.7)	69.6	1.59 (m)	26.4	4.28 (q, 3.3)	71.5	4.08 (q, 3.3)	69.9
2	-	-	-	-	-	-	-	-	1.78 (m)	-	-	-	-	-
3	3.20 (d, 3.7)	91.1	3.63 (d, 3.5)	83.8	3.20 (d, 3.7)	91.0	3.04 (d, 2.2)	90.0	3.58 (dd, 13.3, 4)	79.6	3.66 (d, 3.3)	83.9	3.02 (d, 3.4)	89.6
12	5.28 (t, 3.6)	123.9	5.28 (t, 3.7)	123.7	5.27 (t, 3.6)	123.6	5.12 (t, 3.7)	123.0	5.16 (t, 3.5)	123.0	5.27 (t, 3.4)	123.7	5.12 (t, 3.7)	122.2
18	2.87 (dd, 14, 3)	42.7	2.87 (dd, 13.4, 4.4)	42.8	2.88 (dd, 13.5, 4)	42.9	2.87 (dd, 13.4, 3)	42.6	2.76 (dd, 12.1, 3)	42.4	2.87 (dd, 14, 4.1)	42.7	2.87 (dd, 13.9, 4)	41.3
23	1.10 (s)	29.9	3.24 (d, 11.3)	65.3	1.10 (s)	29.9	0.99 (s)	30.6	3.04 (d, 10.5)	63.9	3.25 (d, 11.3)	65.3	1.10 (s)	28.4
23	-	-	3.64 (d, 11.4)	-	-	-	-	-	3.48 (d, 11.1)	-	3.64 (d, 11.3)	-	-	-
24	1.10 (s)	18.5	0.97 (s)	14.6	1.10 (s)	18.5	0.97 (s)	19.5	0.97 (s)	14.3	0.97 (s)	14.6	1.10 (s)	17.1
25	1.27 (s)	16.8	1.30 (s)	17.4	1.27 (s)	16.8	1.16 (s)	17.2	0.89 (s)	17.0	1.30 (s)	17.4	1.27 (s)	15.4
26	0.85 (s)	17.8	0.85 (s)	17.8	0.86 (s)	17.8	0.74 (s)	18.7	0.73 (s)	18.4	0.85 (s)	17.8	0.85 (s)	16.4
27	1.18 (s)	26.4	1.17 (s)	26.5	1.18 (s)	26.4	1.06 (s)	27.0	1.10 (s)	27.0	1.20 (s)	26.5	1.18 (s)	25.1
28	-	181.9	-	182.0	-	181.8	-	179.7	-	182.0	-	182.0	-	181.0
29	0.93 (s)	33.4	0.93 (s)	33.6	0.93 (s)	33.6	0.84 (s)	34.4	0.87 (s)	34.2	0.93 (s)	33.6	0.93 (s)	32.1
30	0.97 (s)	24.0	0.97 (s)	24.0	0.97 (s)	24.0	0.80 (s)	24.9	0.87 (s)	24.7	0.97 (s)	24.0	0.96 (s)	22.5
<i>β</i> -D-glucuronic acid (at C-3)														
1'	4.46 (d, 7.5)	106.5	4.51 (d, 7.7)	105.3	4.47 (d, 7.5)	106.7	4.40 (d, 8.1)	104.6	4.38 (d, 7.9)	104.2	4.60 (d, 7.8)	105.5	4.53 (d, 7.8)	104.2
2'	3.36 (dd, 9.2, 7.7)	75.0	3.33 (dd, 9.1, 7.7)	75.1	3.36 (t, 7.6)	75.0	3.27 (t, 8.4)	74.2	3.18 (t, 8.5)	73.5	3.56 (dd, 9.5, 8.4)	74.4	3.59 (t, 7.8)	73.0
3'	3.41 (t, 9.2)	77.6	3.42 (t, 9.1)	77.8	3.40 (dd, 9.5, 7.6)	77.4	3.50 (m)	85.7	3.44 (t, 9.5)	86.3	3.65 (m)	86.5	3.66 (m)	84.8
4'	3.53 (t, 9.5)	73.2	3.49 (t, 9.1)	73.3	3.54 (t, 9.5)	73.2	3.30 (t, 9.2)	71.8	3.32 (t, 10.1)	72.0	3.64 (t, 9)	73.0	3.66 (m)	71.6
5'	3.83 (d, 9.7)	76.0	3.77 (d, 10.2)	76.1	3.90 (d, 9.6)	76.5	3.38 (d, 9.9)	75.0	3.38 (d, 9.5)	76.1	3.90 (d, 9)	76.0	3.95 (d, 9.6)	74.6
6'	-	173.0	-	171.0	-	171.5	-	174.2	-	175.0	-	174.2	-	169.7
-OCH <sub>3</sub>	-	-	-	-	3.78 (s)	52.9	-	-	-	-	-	-	3.80 (s)	51.5
<i>β</i> -D-galactose (at C-3')														
1''							4.33 (d, 7.7)	104.9	4.36 (d, 7.9)	105.1	4.55 (d, 7.7)	105.6	4.57 (d, 7.7)	105.0
2''							3.46 (dd, 9.5, 2.7)	71.8	3.47 (t, 8.4)	71.7	3.63 (dd, 9.6, 7.7)	71.9	3.64 (dd, 9.6, 3.4)	70.4
3''							3.33 (dd, 8.5, 3.3)	74.2	3.30 (dd, 9.7, 3)	74.2	3.55 (dd, 9.7, 3.2)	74.7	3.53 (dd, 8.5, 3.3)	73.2
4''							3.62 (d, 3.3)	69.6	3.63 (d, 3)	69.5	3.83 (d, 3.2)	70.4	3.80 (d, 3.3)	68.9
5''							3.42 (m)	77.2	3.44 (m)	77.3	3.59 (ddd, 7.7, 4.3, 3.2)	77.2	3.65 (m)	75.7
6''a							3.51 (m)	61.9	3.51 (m)	61.8	3.71 (dd, 11.5, 4.3)	62.7	3.70 (dd, 11.8, 4.3)	61.2
6''b							3.51 (m)	-	3.51 (m)	-	3.81 (dd, 11.5, 7.7)	-	3.79 (dd, 11.8, 5.5)	-

**Table 3.** <sup>1</sup>H and <sup>13</sup>C NMR Data of Compounds **16** – **21** (CD<sub>3</sub>OD).


	<b>16</b>		<b>17</b>		<b>18</b>		<b>19</b>		<b>20</b>		<b>21</b>	
	<sup>1</sup> H	<sup>13</sup> C	<sup>1</sup> H	<sup>13</sup> C	<sup>1</sup> H	<sup>13</sup> C	<sup>1</sup> H	<sup>13</sup> C	<sup>1</sup> H	<sup>13</sup> C	<sup>1</sup> H	<sup>13</sup> C
Aglycon												
2	4.21 ( <i>q</i> , 3.7)	71.3	1.77 ( <i>m</i> )	26.2	4.28 ( <i>q</i> , 3.1)	71.5	1.77 ( <i>m</i> )	26.4	4.22 ( <i>m</i> )	70.9	1.77 ( <i>m</i> )	26.4
2	-	-	1.97 ( <i>m</i> )	-	-	-	1.95 ( <i>m</i> )	-	-	-	1.89 ( <i>ddd</i> , 13.4, 2.9, 1.3)	-
3	3.22 ( <i>d</i> , 3.7)	91.2	3.70 ( <i>dd</i> , 11, 2.7)	82.2	3.65 ( <i>d</i> , 3.4)	83.9	3.63 ( <i>dd</i> , 11, 2.7)	83.5	3.22 ( <i>d</i> , 3.3)	91.2	3.68 ( <i>dd</i> , 9.7, 3.3)	83.2
12	5.28 ( <i>t</i> , 3.8)	123.9	5.27 ( <i>t</i> , 3.6)	123.8	5.28 ( <i>t</i> , 3.5)	123.8	5.27 ( <i>t</i> , 3.6)	123.8	5.28 ( <i>t</i> , 3.4)	124.0	5.26 ( <i>t</i> , 3.5)	123.8
18	2.88 ( <i>dd</i> , 14, 4.9)	42.6	2.87 ( <i>dd</i> , 13.4, 4.3)	42.6	2.87 ( <i>dd</i> , 13.5, 4.2)	42.6	2.87 ( <i>dd</i> , 13.4, 4.3)	42.6	2.87 ( <i>dd</i> , 14, 3)	42.7	2.87 ( <i>dd</i> , 13.6, 4.2)	42.8
23	1.09 ( <i>s</i> )	29.8	3.27 ( <i>d</i> , 11.4)	64.9	3.25 ( <i>d</i> , 11.3)	65.3	3.27 ( <i>d</i> , 11.4)	64.7	1.10 ( <i>s</i> )	29.9	3.30 ( <i>d</i> , 11.3)	64.8
23	-	-	3.66 ( <i>d</i> , 11.4)	-	3.63 ( <i>d</i> , 11.3)	-	3.66 ( <i>d</i> , 11.4)	-	-	-	3.64 ( <i>d</i> , 11.3)	-
24	1.09 ( <i>s</i> )	18.5	0.72 ( <i>s</i> )	13.4	0.96 ( <i>s</i> )	14.7	0.72 ( <i>s</i> )	13.3	1.10 ( <i>s</i> )	18.6	0.72 ( <i>s</i> )	13.4
25	1.27 ( <i>s</i> )	16.9	1.00 ( <i>s</i> )	16.5	1.30 ( <i>s</i> )	17.5	1.00 ( <i>s</i> )	16.5	1.27 ( <i>s</i> )	16.8	1.00 ( <i>s</i> )	16.4
26	0.83 ( <i>s</i> )	17.8	0.82 ( <i>s</i> )	17.8	0.83 ( <i>s</i> )	17.8	0.82 ( <i>s</i> )	17.8	0.85 ( <i>s</i> )	17.8	0.84 ( <i>s</i> )	17.8
27	1.18 ( <i>s</i> )	26.3	1.19 ( <i>s</i> )	26.3	1.19 ( <i>s</i> )	26.4	1.19 ( <i>s</i> )	26.4	1.18 ( <i>s</i> )	26.4	1.20 ( <i>s</i> )	26.2
28	-	178.0	-	178.1	-	178.1	-	178.1	-	181.9	-	182.0
29	0.93 ( <i>s</i> )	33.5	0.93 ( <i>s</i> )	33.5	0.93 ( <i>s</i> )	33.5	0.93 ( <i>s</i> )	33.5	0.93 ( <i>s</i> )	33.6	0.93 ( <i>s</i> )	33.6
30	0.96 ( <i>s</i> )	24.0	0.96 ( <i>s</i> )	24.0	0.95 ( <i>s</i> )	24.0	0.96 ( <i>s</i> )	24.0	0.97 ( <i>s</i> )	24.0	0.96 ( <i>s</i> )	24.0
β-D-glucuronic acid (at C-3)												
1'	4.53 ( <i>d</i> , 7.8)	106.3	4.51 ( <i>d</i> , 7.9)	104.6	4.60 ( <i>d</i> , 7.9)	105.5	4.54 ( <i>d</i> , 7.9)	105.6	4.53 ( <i>d</i> , 7.7)	106.3	4.54 ( <i>d</i> , 7.9)	105.4
2'	3.57 ( <i>dd</i> , 9.5, 7.8)	74.5	3.47 ( <i>dd</i> , 9.1, 8)	74.6	3.56 ( <i>t</i> , 8.2)	74.4	3.43 ( <i>dd</i> , 9.1, 8)	74.6	3.59 ( <i>dd</i> , 10.1, 7.5)	74.6	3.47 ( <i>dd</i> , 9.7, 7.5)	74.6
3'	3.64 ( <i>t</i> , 9.2)	86.4	3.67 ( <i>t</i> , 9)	86.7	3.65 ( <i>m</i> )	86.5	3.64 ( <i>t</i> , 9)	86.8	3.67 ( <i>m</i> )	85.9	3.64 ( <i>m</i> )	87.3
4'	3.65 ( <i>t</i> , 9.2)	71.9	3.57 ( <i>dd</i> , 9.9, 8.9)	72.1	3.65 ( <i>m</i> )	71.8	3.63 ( <i>t</i> , 8.9)	71.8	3.67 ( <i>m</i> )	71.2	3.63 ( <i>m</i> )	71.2
5'	3.90 ( <i>d</i> , 9.4)	76.0	3.66 ( <i>d</i> , 9.9)	76.8	3.91 ( <i>d</i> , 8.7)	75.4	3.91 ( <i>d</i> , 9.7)	76.3	3.80 ( <i>m</i> )	nd	3.83 ( <i>d</i> , 9)	76.6
6'	-	172.2	-	176.7	-	172.0	-	171.0	-	174.2	-	174.2
-OCH <sub>3</sub>												
							3.80 ( <i>s</i> )	52.9				
β-D-galactose (at C-3')												
1''	4.57 ( <i>d</i> , 7.7)	105.6	4.60 ( <i>d</i> , 7.8)	105.5	4.56 ( <i>d</i> , 7.7)	105.6	4.53 ( <i>d</i> , 7.7)	105.6	4.66 ( <i>d</i> , 7.9)	105.1	4.64 ( <i>d</i> , 7.8)	105.2
2''	3.63 ( <i>dd</i> , 9.7, 7.7)	73.1	3.65 ( <i>dd</i> , 9.8, 7.8)	72.8	3.63 ( <i>dd</i> , 10.1, 7.7)	73.0	3.63 ( <i>dd</i> , 9.6, 7.7)	73.0	3.81 ( <i>dd</i> , 9.6, 7.9)	72.1	3.80 ( <i>dd</i> , 9.8, 8)	72.1
3''	3.53 ( <i>dd</i> , 9.7, 3.3)	74.7	3.53 ( <i>dd</i> , 9.8, 3.4)	74.7	3.53 ( <i>dd</i> , 10.1, 3.4)	74.7	3.57 ( <i>dd</i> , 9.6, 3.3)	74.7	3.69 ( <i>dd</i> , 9.6, 3.2)	84.5	3.68 ( <i>dd</i> , 9.5, 3.1)	84.5
4''	3.82 ( <i>dd</i> , 3.3, 0.8)	70.4	3.82 ( <i>dd</i> , 3.4, 1.2)	70.5	3.82 ( <i>dm</i> , 3.4)	70.4	3.81 ( <i>d</i> , 3.4)	70.3	4.14 ( <i>d</i> , 2.8)	69.9	4.13 ( <i>d</i> , 3.1)	69.9
5''	3.58 ( <i>ddd</i> , 7.8, 4.1, 0.8)	77.2	3.61 ( <i>ddd</i> , 7.8, 4.4, 1.2)	77.2	3.59 ( <i>dd</i> , 7.8, 4.5)	77.2	3.59 ( <i>ddd</i> , 7.8, 4.4, 1.2)	77.2	3.63 ( <i>dd</i> , 7.1, 5)	76.8	3.63 ( <i>m</i> )	76.8
6''a	3.70 ( <i>dd</i> , 11.5, 4.1)	62.7	3.70 ( <i>dd</i> , 11.9, 4.5)	62.6	3.70 ( <i>dd</i> , 11.8, 4.5)	62.7	3.70 ( <i>dd</i> , 11.9, 4.5)	62.2	3.71 ( <i>dd</i> , 10.8, 5)	62.6	3.71 ( <i>dd</i> , 11.9, 4.3)	62.6
6''b	3.80 ( <i>dd</i> , 11.5, 7.8)		3.82 ( <i>m</i> )		3.81 ( <i>dd</i> , 11.8, 7.8)		3.82 ( <i>brd</i> , 11.8)		3.82 ( <i>dd</i> , 10.8, 7.1)		3.77 ( <i>m</i> )	
sugar at C-28 or C-3''	β-D-glucose (at C-28)		β-D-glucose (at C-28)		β-D-glucose (at C-28)		β-D-glucose (at C-28)		β-D-galactose (at C-3'')		β-D-galactose (at C-3'')	
1'''	5.40 ( <i>d</i> , 8.1)	95.7	5.40 ( <i>d</i> , 8.1)	95.7	5.40 ( <i>d</i> , 8.2)	95.7	5.40 ( <i>d</i> , 8.1)	95.7	4.53 ( <i>d</i> , 7.6)	106.3	4.53 ( <i>d</i> , 7.6)	106.3
2'''	3.38 ( <i>dd</i> , 9, 8.1)	73.9	3.34 ( <i>dd</i> , 9.1, 8.1)	73.9	3.34 ( <i>t</i> , 8.6)	73.9	3.35 ( <i>dd</i> , 9.1, 8.1)	73.9	3.64 ( <i>dd</i> , 9.7, 7.7)	73.0	3.64 ( <i>dd</i> , 9.5, 7.7)	73.3
3'''	3.43 ( <i>t</i> , 9)	78.3	3.43 ( <i>t</i> , 9.1)	78.3	3.43 ( <i>t</i> , 8.6)	78.3	3.43 ( <i>t</i> , 9.1)	78.3	3.52 ( <i>dd</i> , 9.7, 3.3)	74.6	3.52 ( <i>dd</i> , 9.5, 3.4)	74.6
4'''	3.37 ( <i>t</i> , 9.5)	71.1	3.38 ( <i>t</i> , 9.5)	71.1	3.38 ( <i>t</i> , 9.5)	71.1	3.38 ( <i>t</i> , 9.5)	71.1	3.84 ( <i>d</i> , 3.3)	70.2	3.84 ( <i>dd</i> , 3.4, 0.5)	70.3
5'''	3.37 ( <i>m</i> )	78.7	3.37 ( <i>m</i> )	78.7	3.37 ( <i>m</i> )	78.7	3.37 ( <i>m</i> )	78.7	3.56 ( <i>ddd</i> , 7, 5, 1)	76.8	3.56 ( <i>ddd</i> , 7.3, 5, 0.5)	76.8
6'''a	3.70 ( <i>dd</i> , 11.5, 4.1)	62.4	3.70 ( <i>dd</i> , 11.5, 4.4)	62.4	3.70 ( <i>dd</i> , 11.8, 4.2)	62.4	3.70 ( <i>dd</i> , 11.5, 4.4)	61.9	3.72 ( <i>dd</i> , 11.3, 5)	62.6	3.72 ( <i>dd</i> , 11.5, 5)	62.6
6'''b	3.83 ( <i>dd</i> , 11.5, 1.8)		3.84 ( <i>m</i> )		3.84 ( <i>brd</i> , 11.8)		3.84 ( <i>brd</i> , 11.8)		3.77 ( <i>dd</i> , 11.3-7.2)		3.77 ( <i>m</i> )	



**Table 4.** Hemolytic activity of the ten major saponins and the methanol extract of the pulp (MEP1) and the peel (MEP2).

Compound	HD <sub>50</sub>		HD <sub>100</sub>	
	$\mu\text{M}$	$\mu\text{g/mL}$	$\mu\text{M}$	$\mu\text{g/mL}$
<b>2</b>	56	45	94	75
<b>3</b>	18	15	24	25
<b>5</b>	-	-	>513	>500 [10%] <sup>a</sup>
<b>28</b>	93	60	116	75
<b>24</b>	19	15	31	25
<b>12</b>	32	26	49	40
<b>13</b>	45	37	93	75
<b>14</b>	66	55	121	100
<b>15</b>	157	130	303	250
<b>17</b>	309	300	514	500
<b>MEP1</b>	-	25	-	100
<b>MEP2</b>	-	25	-	100
<b>Sigma<sup>®</sup> D</b>	-	7.5	-	25


<sup>a</sup> hemolytic percentage observed at this concentration



## Table of Contents Graphic

### Triterpenoid Saponins from the Fruits of *Caryocar glabrum*

Abdulmagid Alabdul Magid, Laurence Voutquenne,\* Christian Moretti, Christophe Long, and Catherine Lavaud



- 2**  $R_1 = H, R_2 = CH_2OH, R_3 = CH_2OH$
- 3**  $R_1 = OH, R_2 = CH_2OH, R_3 = CH_2OH$
- 24**  $R_1 = H, R_2 = CH_3, R_3 = COOH$
- 12**  $R_1 = OH, R_2 = CH_3, R_3 = COOH$
- 13**  $R_1 = H, R_2 = CH_2OH, R_3 = COOH$
- 14**  $R_1 = OH, R_2 = CH_2OH, R_3 = COOH$