



HAL
open science

Acetylated glucuronide triterpene bidesmosidic saponins from *Symplocos glomerata*

Pierre Waffo-Téguo, Laurence Voutquenne, Odile Thoison, Vincent
Dumontet, van Hung Nguyen, Catherine Lavaud

► **To cite this version:**

Pierre Waffo-Téguo, Laurence Voutquenne, Odile Thoison, Vincent Dumontet, van Hung Nguyen, et al.. Acetylated glucuronide triterpene bidesmosidic saponins from *Symplocos glomerata*. *Phytochemistry*, 2004, 65 (6), pp.741-750. hal-01996956

HAL Id: hal-01996956

<https://hal.univ-reims.fr/hal-01996956>

Submitted on 24 Nov 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.


**Acetylated glucuronide bidesmosidic triterpene saponins from
*Symplocos glomerata***

Pierre Waffo-Teguo^a, Laurence Voutquenne^a Odile Thoison^b, Vincent Dumontel^b, Catherine Lavaud

^a *Laboratoire de Pharmacognosie, UMR CNRS 6013 Bât. 18 BP 1039, 51079, Reims Cedex, France.*

^b *ICSN, UPR CNRS, avenue de la Terrasse, 91198 Gif/Yvette Cedex, France.*

Eight oleanolic acid 3-*O*-glucuronide bidesmosidic saponins were isolated from the stem bark of *Symplocos glomerata* along with one bidesmosidic saponin of morolic acid and two known saponins, salsoloside C and copteroside E. The structures were established by spectroscopic methods.



Acetylated glucuronide triterpene bidesmosidic saponins from *Symplocos glomerata*

Pierre Waffo-Teguo*^a, Laurence Voutquenne^a, Odile Thoison^b, Vincent Dumontet^b, Catherine Lavaud^a

^a Laboratoire de Pharmacognosie, UMR CNRS 6013 Bât. 18, BP 1039, 51079 Reims Cedex, France

^b ICSN, UPR 2031, avenue de la Terrasse, 91198 Gif/Yvette Cedex, France.

* Corresponding author

Dr. Pierre Waffo-Teguo

Laboratoire de Pharmacognosie

UMR CNRS 6013, Bât. 18,

BP 1039

51097 Reims Cedex, France.

Tel : 33 (0) 3 26 91 82 08

Fax : 33 (0) 3 26 91 35 96

E-mail : pierre.waffo-teguo@univ-reims.fr

Abstract

Nine new bidesmosidic 3-*O*-glucuronide **oleanane triterpenoid** saponins were isolated from the stem bark of *Symplocos glomerata* King along with two known saponins, salsoloside C and copteroside E, and two major lignans, (-)-pinoresinol and (-)-pinoresinol-4'-*O*- β -D-glucopyranoside. The structures of the new saponins were established using one- and two-dimensional NMR **spectroscopy** and mass spectrometry as, 3-*O*-[β -D-xylopyranosyl(1 \rightarrow 4)-[2-*O*-**acetyl**]- β -D-glucuronopyranosyl]-28-*O*-[β -D-glucopyranosyl]- oleanolic acid, 3-*O*-[β -D-xylopyranosyl(1 \rightarrow 4)-[3-*O*-**acetyl**]- β -D-glucuronopyranosyl]-28-*O*-[β -D-glucopyranosyl]- oleanolic acid, 3-*O*-[β -D-xylopyranosyl (1 \rightarrow 4)-[2,3-*O*-**diacetyl**]- β -D-glucuronopyranosyl]-28-*O*-[β -D-glucopyranosyl]-oleanolic acid, 3-*O*-[α -L-arabinopyranosyl(1 \rightarrow 4)- β -D-glucuronopyranosyl]-28-*O*-[β -D-glucopyranosyl]- oleanolic acid, 3-*O*-[α -L-arabinopyranosyl (1 \rightarrow 4)-[2-*O*- **acetyl**]- β -D-glucuronopyranosyl]-28-*O*-[β -D-glucopyranosyl]- oleanolic acid, 3-*O*-{[β -D-xylopyranosyl (1 \rightarrow 2)]-[β -D-xylopyranosyl (1 \rightarrow 4)]-[3-*O*-**acetyl**]- β -D-glucuronopyranosyl}-28-*O*-[β -D-glucopyranosyl]- oleanolic acid, 3-*O*-{[β -D-glucopyranosyl (1 \rightarrow 2)]-[β -D-xylopyranosyl (1 \rightarrow 4)]-[3-*O*-**acetyl**]- β -D-glucuronopyranosyl}-28-*O*-[β -D-glucopyranosyl] - oleanolic acid, 3-*O*-{[β -D-glucopyranosyl (1 \rightarrow 2)]-[α -L-arabinofuranosyl (1 \rightarrow 4)]-[3-*O*-**acetyl**]- β -D-glucuronopyranosyl}-28-*O*-[β -D-glucopyranosyl]- oleanolic acid, and 3 β -*O*-[β -D-xylopyranosyl(1 \rightarrow 4)-[2-*O*-**acetyl**]- β -D-glucuronopyranosyl]-28-*O*-[β -D-glucopyranosyl]- morolic acid. The EtOH and EtOAc extracts of the stem bark showed no cytotoxic activity. At a concentration of 370 μ g/ml, the saponin mixture showed haemolytic activity and caused 50% haemolysis of a 10% suspension of sheep erythrocytes.

Keywords: *Symplocos glomerata*; Symplocaceae; Triterpenoid saponins; Oleanolic acid; Morolic acid; Haemolysis.

1. Introduction

The genus *Symplocos* of the Symplocaceae family consists of 300-500 species, and is most common in tropical and subtropical Asia, Malaysia and America (Hegnauer, 1973). Previously, flavonoids (Tschesche et al., 1980, Tanaka et al., 1980, 1982, Lin et al., 1996), iridoids (Iida et al., 1990), lignans (Ishida et al., 2001), steroids (Frotan et al., 1983) and triterpenoids (Ali et al., 1990) have been found as constituents of this genus. Previous phytochemical studies revealed the presence of triterpenoid saponins in *Symplocos spicata* (Higuchi et al., 1982). *Symplocos* species are used for antimicrobial activity (Khan et al., 2001), and for the treatment of diarrhoea, dysentery, eye diseases, hemorrhagic gingivitis, menorrhagia and uterine disorders (Ali et al., 1990), female diseases (Ahmad et al., 2003), bowel complaints and ulcers (Dhaon et al., 1989).

The stem bark of *S. glomerata* King was collected in Pà Co, Mai Châu (Hoà Binh) in Vietnam. The chemical constituents and biological activity of this plant have not been previously investigated. This species was selected in a screening program for potential cytotoxic compounds from plants growing in Vietnam. The ethanolic and ethyl acetate extracts of the stem bark were tested *in vitro* against KB cells, but showed no cytotoxic activity. Primary chemical studies on the ethanolic and ethyl acetate extracts of leaves, stem bark and fruits, showed the presence of saponins. The present paper reports the isolation and the structural elucidation of one bidesmosidic morolic acid saponin and eight bidesmosidic saponins possessing oleanolic acid as aglycone, along with two known saponins, salsolide C and copteroside E.

2. Results and discussion

The dried and powdered stem bark of *S. glomerata* was extracted with boiling 80% methanol. The aqueous methanolic extract was concentrated and precipitated into acetone to give saponin mixture A. The filtrate was concentrated and again precipitated in diethyl ether to give saponin mixture B. Each crude saponin precipitate was dialysed. Saponin mixture A and B showed the same contents by TLC analysis, and saponin mixture B, was chromatographed on a silica gel column. Purification by reversed phase C-18 column chromatography and semi-preparative HPLC over C-18 reversed phase afforded eleven saponins (**1-11**). In addition to saponins, two lignans, (-)-pinoresinol and (-)-pinoresinol-4'-*O*- β -D-glucopyranoside were also isolated from saponin mixture B. The spectroscopic data of these compounds were found to be identical with those reported in the literature (Cuenca and Catalan, 1991; Nishibe et al., 1984; Casabuono and Pomilio, 1994).

Acid hydrolysis of saponin mixture A gave only oleanolic acid, purified by silica gel column chromatography. The sugars were identified as D-glucose, D-glucuronic acid, L-arabinose, and D-xylose by TLC and their absolute configurations were determined by the measurement of optical rotation after separation by prep. TLC.

Oleanolic acid was identified as the aglycone of compounds **1-10** on the basis of its $^1\text{H-NMR}$, $^{13}\text{C-NMR}$, COSY, ROESY, HSQC, and HMBC spectra. The ^{13}C chemical shifts for C-28 between δ 176.2 and 176.6, and for C-3 between δ 89.7 and 91 suggested that sugars were connected to these points (Tables 1 and 2). Moreover, the $^{13}\text{C-NMR}$ shifts of the aglycone part of compounds **1-10** were in good agreement with those reported for bidesmosidic saponins of oleanolic acid (Tan et al., 1999).

Saponin **1**, molecular formula $\text{C}_{47}\text{H}_{74}\text{O}_{18}$ (ESI-MS $^-$: m/z 925 [M-H] $^-$) and saponin **7**, molecular formula $\text{C}_{52}\text{H}_{82}\text{O}_{22}$ (ESI-MS $^-$: m/z 1057 [M-H] $^-$), were identified as salsolide C, 3-*O*-[β -D-

xylopyranosyl-(1→4)- β -D-glucuronopyranosyl]-28-O-[β -D-glucopyranosyl]-oleanolic acid, and copteroside E, 3-O-[[β -D-xylopyranosyl (1→2)]-[β -D-xylopyranosyl (1→4)]- β -D-glucuronopyranosyl]-28-O-[β -D-glucopyranosyl]- oleanolic acid, respectively, on the basis of their spectral data. Compound **1** was previously isolated from *Salsola micranthera* (Annaev et al., 1983a), *Aralia cordata* (Kawai et al., 1989), *Melanthera scandens* (Penders and Delaude, 1994) and *Aralia continentalis* (Kim and Kang., 1998), while compound **7** was isolated from *Climacoptera transoxana* (Annaev et al., 1983b) and *Verbesina suncho* (Cerda-Garcia-Rojas et al., 2000) .

The positive ESI-MS of compound **2** gave a quasi-molecular ion peak at m/z 991 $[M+Na]^+$ while, in the negative mode, a quasi-molecular ion was detected at m/z 967 $[M-H]^-$ indicating a M_r of 968 amu in agreement with a molecular formula of $C_{49}H_{76}O_{19}$ and suggesting the presence of an acetyl group compared to saponin **1**. The 1H and ^{13}C NMR spectra of **2** showed signals for a single acetyl ester at δ_H 2.08 and δ_C 19.8 (CH_3) and 171.1 (CO). The MS^2 experiment of the $[M+Na]^+$ ion, gave positive fragments at m/z 859 $[M+Na-132]^+$, 829 $[M+Na-162]^+$, 697 $[M+Na-132-162]^+$ and 623 $[M+Na-132-176-CH_3CO_2H]^+$, attributed to the loss of a terminal pentose, a terminal hexose, both terminal sugars and a diglycoside chain consisting of a pentose and an acetylated uronic acid, respectively. These results suggested that the saponin **2** contained three sugars units, one of which is an acetylated uronic acid.

The three anomeric carbons were detected at 94.2, 103.8 and 105.2 in the ^{13}C -NMR spectrum, attached to doublets at δ 5.4, 4.28 and 4.51, respectively in the HSQC experiment. Complete assignments of each glycosidic proton system were achieved by analysis of COSY and TOCSY experiments. The units with anomeric protons at δ 5.4 ($J = 8.1$) and 4.28 ($J = 7.4$) corresponded to an hexose and a pentose with their methylene carbons at δ 60.9 and 65.6, respectively, and were identified as β -D-glucose and β -D-xylose (Table 1). The third glycosidic unit with anomeric proton at δ 4.51 ($J = 7.8$) was identified as a β -D-glucuronic

acid which possessed a deshielded H-3' at δ 5.03 indicating the position of acetylation. The deshielding of C-4' (δ 77.2) of glucuronic acid suggested the point of linkage of the xylose (Table 1).

Sequencing of the glycosidic chains in saponin **2** was achieved by analysis of HMBC and ROESY experiments. The HMBC spectrum showed cross peaks between C-28 (δ 176.6) of oleanolic acid and H-1'' of the glucose ester (δ 5.40), between C-3 (δ 89.9) of oleanolic acid and H-1' of esterified glucuronic acid (δ 4.51), and between C-4' (δ 77.2) of this glucuronic acid and H-1''' of the terminal xylose unit (δ 4.28). Thus, saponin **2** is 3-*O*-[β -D-xylopyranosyl-(1 \rightarrow 4)-[3-*O*-acetyl]- β -D-glucuronopyranosyl]-28-*O*-[β -D-glucopyranosyl]-oleanolic acid or 3'-*O*-acetylsalsolide C.

The ESI-MS of compound **3** gave quasi-molecular ion peaks at m/z 991 [M+Na]⁺ (positive mode) and m/z 967 [M-H]⁻ (negative mode) indicating a M_r of 968 amu in agreement with a molecular formula of C₄₉H₇₆O₁₉ isomeric with saponin **2**. The MS² experiment of the [M+Na]⁺ ion, gave the same positive fragments described for **2**. The MS⁴ experiment of the ion at m/z 641 gave an ion fragment at m/z 479 [genin+Na]⁺ due to the loss of the glycosidic chains. The presence of three sugar residues was confirmed from the observation of signals for three anomeric positions at δ_C 94.2, 102.9 and 103.4 and δ_H 5.4 ($J = 8.2$), 4.63 ($J = 8$) and 4.4 ($J = 7.5$). Inspection of the spectroscopic data indicated that saponin **3** had the same glycosidic part as saponin **2**: a β -D-glucopyranosyl ester (δ_H 5.4), a terminal β -D-xylopyranosyl (δ_H 4.40) and a 4-substituted β -D-glucuronopyranosyl (δ_H 4.63). These results suggested that compound **3** is a regioisomer of **2** with the acetate attached to C-2 rather than to C-3 (see Table 1). Thus, saponin **3** is 3-*O*-[β -D-xylopyranosyl-(1 \rightarrow 4)-[2-*O*-acetyl]- β -D-glucuronopyranosyl]-28-*O*-[β -D-glucopyranosyl]-oleanolic acid or 2'-*O*-acetyl salsolide C. The negative ESI-MS of saponin **4** gave a quasi-molecular ion at m/z 1009 [M-H]⁻, indicating a molecular formula of C₅₁H₇₈O₂₀ (M_r 1010). The MS/MS fragmentation indicated a

supplementary acetyl unit compared to saponins **2** and **3**. The ^1H and ^{13}C NMR data of compound **4** were closely comparable to those of **2** and **3** except for the signals of glucuronic acid and the presence of two acetate methyls (δ_{H} 2.01 and 2.02) (Table 1). The β -D-glucuronic acid possessed two deshielded protons H-2' and H-3' at δ 4.9 and 5.19, respectively, indicating the position of the acetates. Thus saponin **4** is 3-*O*-[β -D-xylopyranosyl-(1 \rightarrow 4)-[2,3-*O*-diacetyl]- β -D-glucuronopyranosyl]-28-*O*-[β -D-glucopyranosyl]-oleanolic acid or 2',3'-*O*-diacetylsalsolide C.

The positive ESI-MS of compound **5** gave a quasi-molecular ion peak at m/z 971 [$\text{M}+2\text{Na}-\text{H}$] $^+$ while, in the negative mode, a quasi-molecular ion was detected at m/z 925 [$\text{M}-\text{H}$] $^-$ indicating a M_r of 926 amu in agreement with a molecular formula of $\text{C}_{47}\text{H}_{74}\text{O}_{18}$. The MS^2 experiment of the [$\text{M}+2\text{Na}-\text{H}$] $^+$ ion, gave positive fragments at m/z 839 [$\text{M}+\text{Na}-\text{H}-132$] $^+$ and 809 [$\text{M}+2\text{Na}-\text{H}-162$] $^+$, attributed to the loss of a terminal pentose and a terminal hexose. The MS^3 fragmentation of the ion at m/z 809 gave an ion fragment at m/z 677 [$\text{M}+2\text{Na}-\text{H}-162-132$] $^+$ corresponding to the loss of both hexose and pentose. The MS^4 fragmentation of the ion at m/z 677 gave a product ion at m/z 501 [genin+2Na-H] $^+$ after the loss of an uronic acid unit. This analysis suggested that saponin **5** contained three sugars units and was isomeric of saponin **1**.

Three anomeric carbons were detected at δ 94.2, 103.4 and 105.4 in the ^{13}C -NMR spectrum, attached to doublets at δ 5.4, 4.35 and 4.44, respectively in HSQC experiment (Table 1). Analysis of 2D experiments (COSY, TOCSY and HSQC) revealed a β -D-glucopyranosyl ester (δ_{H} 5.4, δ 94.2) attached to C-28 of the genin, a 4-substituted β -D-glucuronic acid, and of a terminal α -L-arabinose. The cross peaks observed in the HMBC experiment between C-28 (δ 176.6) of oleanolic acid and H-1'' of glucose (δ 5.4), between C-3 (δ 89.8) of the genin and H-1' of glucuronic acid (δ 4.44) and between C-4' (δ 79.7) of this glucuronic acid and H-1''' (δ 4.35) of the α -L-arabinose unit, showed that compound **5** is 3-*O*-[α -L-

arabinopyranosyl-(1→4)- β -D-glucuronopyranosyl]-28-*O*-[β -D-glucopyranosyl]- oleanolic acid.

Saponin **6** exhibited an intense quasi-molecular ion peak at m/z 1013 $[M+2Na-H]^+$ while, in the negative mode, a quasi-molecular ion was detected at m/z 967 $[M-H]^-$ indicating a M_r of 968 ($C_{49}H_{76}O_{19}$). This result suggested an additional acetyl group compared with saponin **5**. The MS² experiment of the $[M+2Na-H]^+$ ion gave positive fragments at m/z 953 $[M+2Na-H-60]^+$, 881 $[M+2Na-H-132]^+$ and 851 $[M+2Na-H-162]^+$, attributed to the loss of acetic acid, a terminal pentose and a terminal hexose, respectively. The 2D experiments (COSY, TOCSY and HSQC) permitted the assignment of all the ¹H and ¹³C signals of the sugars, identified as a terminal β -D-glucopyranosyl (δ_H 5.37, δ_C 94.2), a terminal α -L-arabinopyranosyl (δ_H 4.34, δ_C 103.3) and a 4-substituted β -D-glucuronic acid (δ_H 4.61, δ_C 102.9). Relative to **5**, the identical value observed for C-4' of β -D-glucuronic acid at δ 79.7 ppm confirmed that the α -L-arabinopyranosyl was attached at this position and the deshielded value of H-2' (δ 4.8) suggested that the glucuronic acid was acetylated at position 2. HMBC correlations showed that the sugars were attached in the same way in both saponins **5** and **6**. Consequently, the structure of saponin **6** is 3-*O*-[α -L-arabinopyranosyl-(1→4)-[2-*O*-acetyl]- β -D-glucuronopyranosyl]-28-*O*-[β -D-glucopyranosyl]-oleanolic acid.

The ESI-MS of saponin **8** gave a quasi-molecular ion peaks at m/z 1145 $[M+2Na-H]^+$ (positive mode) and at m/z 1099 $[M-H]^-$ (negative mode) indicating a M_r of 1100 amu ($C_{54}H_{84}O_{23}$), and suggesting a supplementary pentose unit relative to saponin **2**. Saponin **8** was shown to contain four glycosidic residues and one acetate group from the HSQC spectrum. The anomeric proton signals at δ 5.4, 4.65, 4.37 and 4.29 had correlations with carbon signals at δ 94.2, 103.8, 104 and 103.8, respectively. Analysis of 2D experiments permitted the assignment of the four sugars: a β -D-glucopyranosyl ester (δ_H 5.4, δ 94.2) attached to C-28 of oleanolic acid, two terminal β -D-xyloses and one 2,4-disubstituted β -D-

glucuronic acid. The cross peaks observed in the ROESY experiment between H-3 (δ 3.17) of the genin and H-1' of glucuronic acid (δ 4.65) and between H-2' (δ 3.71) and H-4' (δ 3.94) of this glucuronic acid and H-1''' of the first β -D-xylose (δ 4.37) and H-1'''' of the second β -D-xylose (δ 4.29), respectively, showed that the saponin **8** was an acetylated copteroside E **7**. The acetyl group was readily assigned to C-3 of the glucuronic acid (see Table 2). Thus, saponin **8** is 3-*O*-{[β -D-xylopyranosyl-(1 \rightarrow 2)]-[β -D-xylopyranosyl-(1 \rightarrow 4)]-[3-*O*-acetyl]- β -D-glucuronopyranosyl}-28-*O*-[β -D-glucopyranosyl]- oleanolic acid. This saponin corresponded to the 3'-*O*-acetyl derivative of the known copteroside E (Annaev et al., 1983b). The negative ESI-MS of saponin **9** gave a quasi-molecular ion peak at m/z 1129 [M-H]⁻ indicating a M_r of 1130 amu (C₅₅H₈₆O₂₄), and suggesting a supplementary hexose unit compared to saponin **2**. The analysis of MS/MS fragmentation showed that this supplementary hexose unit replaced the pentose unit found in saponin **8**.

Analysis of 2D experiments (COSY, TOCSY, HSQC and HMBC) allowed the identification of the glycosidic part of saponin **8**: a β -D-glucopyranosyl ester (δ_H 5.4, δ 94.2) attached to C-28 of oleanolic acid, a terminal β -D-xylose, and a 2,4-disubstituted β -D-glucuronic acid acetylated in position 3 (deshielded H-3' at δ 5.86). The anomeric proton of the fourth sugar appeared at δ 5.13 ($J = 7.6$). This was identified as a second β -D-glucopyranosyl unit with its hydroxymethyl group at δ_C 63.2 (Table 2). The HMBC experiment showed correlations between C-3 (δ 89.7) of the genin and H-1' of glucuronic acid (δ 4.85) and between C-2' (δ 77.2) and C-4' (δ 78.9) of glucuronic acid and H-1''' of the second β -D-glucose (δ 5.13) and H-1'''' of β -D-xylose (δ 5.03), respectively. Thus, compound **9** is 3-*O*-{[β -D-glucopyranosyl-(1 \rightarrow 2)]-[β -D-xylopyranosyl-(1 \rightarrow 4)]-[3-*O*-acetyl]- β -D-glucuronopyranosyl}-28-*O*-[β -D-glucopyranosyl]- oleanolic acid. This saponin corresponded to the 3'-*O*-acetyl derivative of the known salsolose E (Annaev et al., 1984).

The positive ESI-MS of compound **10** gave a quasi-molecular ion peak at m/z 1153 $[M+Na]^+$ while, in the negative mode, a quasi-molecular ion was detected at m/z 1129 $[M-H]^-$ indicating a M_r of 1130 amu ($C_{55}H_{86}O_{24}$) as for saponin **9**. The MS^2 experiment of the $[M+Na]^+$ ion, gave positive fragments at m/z 1021 $[M+Na-132]^+$ and 991 $[M+Na-162]^+$, attributed to the loss of a terminal pentose and the loss of a terminal hexose, and at m/z 535 corresponding to the loss of the triglycosidic chain at position 3 of oleanolic acid. Comparison (not comparison!) of the NMR spectra of compound **10** and saponin **9** indicated that **10** possessed an α -L-arabinofuranose instead of a β -D-xylose (Table 2). This α -L-arabinofuranose was characterized by the chemical shifts of its anomeric proton (δ_H 4.96) and carbon (δ_C 109.4).

Sequencing of the sugar chains in saponin **10** was achieved by analysis of HMBC experiments, which showed cross peaks between C-28 (δ 176.6) of oleanolic acid and H-1'' of the glucose ester (δ 5.41), and between H-1' (δ 4.69), H-2' (δ 3.86), H-3' (δ 5.27) and H-4' (δ 3.9) of the trisubstituted glucuronic acid with C-3 (δ 91) of oleanolic acid, C-1''' (δ 102.8) of glucose, the carbonyl (δ 171.1) of the acetyl unit and C-1'''' (δ 109.4) of arabinose, respectively. Thus, saponin **10** was concluded to be 3-*O*-{ $[\beta$ -D-glucopyranosyl (1 \rightarrow 2)] [α -L-arabinofuranosyl-(1 \rightarrow 4)]-[3-*O*-acetyl]- β -D-glucuronopyranosyl]-28-*O*-[β -D-glucopyranosyl]-oleanolic acid.

The negative ESI-MS of compound **11** gave a quasi-molecular ion at m/z 967 $[M-H]^-$ indicating a molecular formula of $C_{49}H_{76}O_{19}$ (M_r 968) isomeric with saponins **2**, **3** and **6**. The MS^2 experiment of the positive $[M+2Na-H]^+$ molecular ion at m/z 1013, gave positive fragments at m/z 953 $[M+2Na-H-60]^+$ and 851 $[M+2Na-H-162]^+$, attributed to the loss of acetic acid and a terminal hexose. The MS^3 experiment of the $[M+2Na-H-162]^+$ ion, gave positive fragments at m/z 791 $[M+2Na-H-162-60]^+$ and 719 $[M+2Na-H-162-132]^+$, attributed to the loss of acetic acid, and a terminal pentose. The analysis of 1D- and 2D-NMR

experiments (^1H , ^{13}C , COSY, TOCSY, HSQC and HMBC) of saponin **11** demonstrated that the glycosidic chains of compounds **3** and **11** were identical (Table 1). The ^1H NMR spectrum of **11** showed a trisubstituted olefinic proton at δ 5.16 (H-19) correlated in the HMBC spectrum with C-13 (δ 40.8), C-18 (δ 137), C-20 (δ 31.5), C-29 (δ 29.3) and C-30 (δ 27.9), indicating that the double bond is between C-18 and C-19 instead of C-12 and C-13. The ^{13}C NMR chemical shifts of the signals due to the aglycone part of saponin **11** were in close agreement with those reported for morolic acid (Yoshikawa et al., 1997). In the ROESY spectrum, the Overhauser interaction of the axial proton H-13 at δ 2.29 (*dd*, $J = 11.4, 2.5$) with the methyl group at δ 1.03 (CH_3 -26), assigned from HMBC correlations, indicated that H-13 is in a β -axial position. Thus, the structure of saponin **11** is $3\beta\text{-O-}[\beta\text{-D-xylopyranosyl}(1\rightarrow4)\text{-}2\text{-O-acetyl-}\beta\text{-D-glucuronopyranosyl}]\text{-}28\text{-O-}[\beta\text{-D-glucopyranosyl-}]\text{-morolic acid}$.

The haemolytic activity of the purified saponin mixture and of pure saponins **1**, **2**, **3**, **8**, **9** and **10** was assessed on sheep erythrocytes using the method previously described (Voutquenne et al., 2002). The isolated quantities of saponins **4**, **5**, **6** and **7** were not sufficient to allow evaluation of biological activity. Saponin mixture B was active and 50% haemolysis was obtained at 370 $\mu\text{g/ml}$. Amongst the tested compounds, only saponin **2** exhibited haemolytic activity with an HD_{50} at 216 μM (209.1 $\mu\text{g/ml}$).

3. Experimental

3.1 General experimental procedures

^1H and ^{13}C NMR spectra were recorded on a Bruker Avance DRX 500 (^1H at 500 MHz and ^{13}C at 125 MHz). 2D experiments were performed using standard Bruker microprograms. ESI-MS and MS-MS experiments were recorded on a Finnigan LCQ deca quadrupole ion trap mass spectrometer (Finnigan MAT, San Jose, USA). The samples were introduced by direct infusion in a solution of MeOH at a rate of $5\ \mu\text{L}\ \text{min}^{-1}$. Optical rotations were determined in MeOH or pyridine with a Perkin-Elmer 241 polarimeter. TLC was performed on pre-coated silica gel 60 F₂₅₄ Merck and detection was achieved by H₂SO₄ 50%. CC were carried out on Kieselgel 60 (63-200 mesh) Merck or LiChroprep RP-18 (40-63 μm) Merck. HPLC was performed on a DIONEX apparatus equipped with an ASI-100 autosampler, a P580 pump, a diode array detector UVD 340S and a Chromeleon[®] software. C18 reversed phase column (201SP, 250×10mm, 5 μm , Dionex, Jouyenosap, France) was used for semi-preparative HPLC with a binary gradient eluent (solvent A, H₂O (pH 2.4 with TFA); solvent B, MeCN) and a flow rate $5\ \text{mL}\ \text{min}^{-1}$; the chromatogram was monitored at 205 nm.

3.2. Plant material

Stem bark of *S. glomerata* King. (VN076) was collected in PàCo, Mai chau (Hoa Binh), Vietnam in april 1996 by Vincent Dumontet. The specimen of the plant (VN 076) is deposited in the herbarium of CNST of Hanoi (Vietnam).

3.3. Extraction and isolation

Dried and finely powdered stem bark of the plant (1 kg) was macerated in 20% aq. MeOH (15 l) for 2 h and then boiled for 3 h. The hydromethanolic extract was filtered, and then evaporated to yield a residue (125 g). This residue was suspended in MeOH (300 ml), then poured dropwise into 1.5 l of Me₂CO. The resulting ppt. was filtered and dried over KOH *in vacuo* (42.8 g). The filtrate was evaporated, dried, suspended in MeOH and precipitated into

Et₂O and the ppt. was filtered and dried over KOH *in vacuo* (62.5 g). The two dried ppt. (62.5 g) were dissolved in H₂O and dialysed against H₂O in seamless cellulose tubing **with** agitation during 48 h. The contents of the tubes were freeze-dried to afford 10.5g **of** saponin mixt. A (from ppt. Me₂CO) and 10.7 g **of** saponin mixt. B (from ppt.Et₂O).

An aliquot of saponin mixt. B (10g) was fractionated on silica gel CC, using a gradient of CHCl₃/MeOH (100/0 to 0/100). Frs. [9-16] = I eluted with CHCl₃/MeOH (98/2) was purified by preparative TLC in CHCl₃/MeOH (9/1) to give (-)-pinoresinol (20mg). Frs [46-62] = II eluted with CHCl₃/MeOH (95/5) **yielded (-)-pinoresinol-4'-O-β-D-glucopyranoside** (1.5 g). Frs [131-134] = III eluted with CHCl₃/MeOH (7/3) and containing saponins **1-4** and **11** was purified on a reversed-phase RP18 CC using a gradient of MeOH/H₂O (6/4 to 8/2): Frs. [6-18] eluted with MeOH/H₂O (6/4) was purified by semi-prep. HPLC with the elution prog.: 30-50%B (0-20 min) and 50% B (20-25 min) to give saponin **1** (rt =14 min, 11 mg) and saponin **2** (rt = 16 min, 10 mg); Frs [22-46] eluted with MeOH/H₂O (6/4) was purified by semi-prep. HPLC 45%B (0-15min) to give saponins **3** (rt = 8.78 min, 14 mg) and **4** (rt = 10 min, 5mg); saponin **11** (rt = 9.32 min, 5 mg) was purified from Frs. [53-54] eluted with MeOH/H₂O (7/3) by semi-prep. HPLC using the same prog. as above. Frs. [135-139] = IV eluted with CHCl₃/MeOH (7/3) and containing saponins **8, 9** and **10** was chromatographed on a reversed-phase RP18 CC using a gradient of MeOH/H₂O (5/5 to 7/3): Frs. [35-57] eluted with MeOH/H₂O (6/4) was purified by semi-prep. HPLC, using a linear gradient 30 to 35% B in 40 min. to yield saponins **9** (rt = 10.90 min, 23 mg), **10** (rt = 12.58 min, 14 mg) and **8** (rt = 13.52 min, 18 mg). Frs. [145-157] = V eluted with CHCl₃/MeOH (7/3) was purified on a RP18 CC using a gradient of MeOH/H₂O (5/5 to 7/3) to give saponins **5, 6** and **7**: Frs. [42-45] eluted with MeOH/H₂O (6/4) was purified by semi-prep. HPLC, using a linear gradient 30 to 50% B in 30 min to give saponins **5** (rt = 8.33 min, 6 mg) and **7** (rt = 9.61 min, 8 mg). Saponin **6** (rt =

14.19 min, 9 mg) eluted with MeOH/H₂O (7/3) from Frs. [58-61] was finally purified by semi-prep. HPLC using the same prog. as above.

3.4. saponin 2

White powder [α]²¹_D +9 (MeOH, *c* 0.5); ¹H and ¹³C NMR (CD₃OD) of the glycosidic part: see Table 1; ¹³C NMR (CD₃OD) of the aglycone δ 14.5 (C-25), 15.5 (C-24), 16.2 (C-26), 17.8 (C-6), 22.5 (C-30), 22.5 (C-16), 23.1 (C-11), 24.8 (C-27), 25.5 (C-2), 27 (C-23), 27.4 (C-15), 30.1 (C-20), 31.7 (C-22), 32 (C-29), 32.5 (C-7), 33.4 (C-21), 36.4 (C-10), 38.3 (C-1), 38.7 (C-4), 39.2 (C-8), 41.1 (C-18), 41.6 (C-14), 45.8 (C-19), 46.5 (C-17), 47.7 (C-9), 55.5 (C-5), 89.9 (C-3), 122.3 (C-12), 143.4 (C-13), 176.6 (C-28); ESI-MS (negative ion mode) *m/z* 967 [M-H]⁻; ESI-MS-MS: MS² (967) *m/z* 817 [(M-H)-132-H₂O]⁻, 805 [(M-H)-162]⁻, 757 [(M-H)-132-H₂O-CH₃COOH]⁻, 611 [(M-H)-132-162-CH₃CO-H₂O]⁻, MS³ (757) *m/z* 595 [(M-H)-132-H₂O-CH₃CO₂H-162]⁻; ESI-MS (positive ion mode) *m/z* 991 [M+Na]⁺; ESI-MS-MS: MS² (991) *m/z* 859 [(M+Na)-132]⁺, 829[(M+Na)-162]⁺, 697 [(M+Na)-132-162]⁺, 641 [(M+Na)-132-176-CH₃CO₂H]⁺, 623 [(M+Na)-132-176-CH₃CO₂H-H₂O]⁺, MS⁴ (623) *m/z* 461 [(genin+Na)-H₂O]⁺.

3.5. saponin 3

White powder [α]²¹_D 0 (MeOH, *c* 0.46); ¹H and ¹³C NMR (CD₃OD) of the glycosidic part: see Table 1, ¹³C NMR chemical shift values of aglycone are identical to those described for saponin 2; ESI-MS (negative ion mode) *m/z* 967 [M-H]⁻; ESI-MS-MS: MS² (967) *m/z* 817 [(M-H)-132-H₂O]⁻, 655 [(M-H)-132-162-H₂O]⁻, MS³ (817) *m/z* 655 [(M-H)-132-162-H₂O]⁻; ESI-MS (positive ion mode) *m/z* 991 [M+Na]⁺; ESI-MS-MS : MS² (991) *m/z* 859 [(M+Na)-132]⁺, 829 [(M+Na)-162]⁺, 641 [(M+Na)-132-176-CH₃CO₂H]⁺, MS³ (859) *m/z* 623 [(M+Na)-132-176-CH₃CO₂H-H₂O]⁺, MS³ (641) *m/z* 479 [genin+Na]⁺.

3.6. saponin 4

White powder $[\alpha]_D^{21} +1.5$ (MeOH, c 0.2); ^1H and ^{13}C NMR (CD_3OD) of the glycosidic part: see Table 1, ^{13}C NMR chemical shift values of the aglycone are identical to those described for saponin **2** to ± 0.2 ppm; ESI-MS (negative ion mode) m/z 1009 $[\text{M}-\text{H}]^-$; ESI-MS-MS: MS^2 (1009) m/z 859 $[(\text{M}-\text{H})-132-\text{H}_2\text{O}]^-$, 799 $[(\text{M}-\text{H})-132-\text{H}_2\text{O}-\text{CH}_3\text{CO}_2\text{H}]^-$, 653 $[(\text{M}-\text{H})-132-162-\text{CO}_2-\text{H}_2\text{O}]^-$, 637 $[(\text{M}-\text{H})-132-162-\text{H}_2\text{O}-\text{CH}_3\text{CO}_2\text{H}]^-$, 593 $[(\text{M}-\text{H})-132-162-\text{CO}_2-\text{H}_2\text{O}-\text{CH}_3\text{CO}_2\text{H}]^-$, 577 $[(\text{M}-\text{H})-132-162-\text{H}_2\text{O}-2\text{CH}_3\text{CO}_2\text{H}]^-$; ESI-MS (positive ion mode) m/z 1055 $[\text{M}+2\text{Na}-\text{H}]^+$, 1033 $[\text{M}+\text{Na}]^+$; ESI-MS-MS : MS^2 (1055) m/z 893 $[(\text{M}+2\text{Na}-\text{H})-162]^+$, MS^3 (893) m/z 875 $[(\text{M}+2\text{Na}-\text{H})-162-\text{H}_2\text{O}]^+$, 833 $[(\text{M}+2\text{Na}-\text{H})-162-\text{CH}_3\text{CO}_2\text{H}]^+$, 639 $[(\text{M}+2\text{Na}-\text{H})-132-162-\text{CH}_3\text{CO}_2\text{H}-\text{CO}_2]^+$.

3.7. saponin 5

White powder $[\alpha]_D^{21} +4.7$ (MeOH, c 0.38); ^1H and ^{13}C NMR (CD_3OD) of the glycosidic part: see Table 1, ^{13}C NMR chemical shift values of the aglycone are identical to those described for saponin **2**; ESI-MS (negative ion mode) m/z 925 $[\text{M}-\text{H}]^-$; ESI-MS-MS: MS^2 (925) m/z 775 $[(\text{M}-\text{H})-132-\text{H}_2\text{O}]^-$, 613 $[(\text{M}-\text{H})-132-162-\text{H}_2\text{O}]^-$, MS^4 (613) m/z 455 $[\text{genin}-\text{H}]^-$; ESI-MS (positive ion mode) m/z 971 $[\text{M}+2\text{Na}-\text{H}]^+$; ESI-MS-MS : MS^2 (971) m/z 839 $[(\text{M}+2\text{Na}-\text{H})-132]^+$, 809 $[(\text{M}+2\text{Na}-\text{H})-162]^+$, MS^3 (809) m/z 791 $[(\text{M}+2\text{Na}-\text{H})-162-\text{H}_2\text{O}]^+$, 677 $[(\text{M}+2\text{Na}-\text{H})-162-132]^+$, MS^4 (677) m/z 659 $[(\text{M}+2\text{Na}-\text{H})-162-132-\text{H}_2\text{O}]^+$, 501 $[\text{genin}+2\text{Na}-\text{H}]^+$.

3.8. saponin 6

White powder $[\alpha]_D^{21} +8$ (MeOH, c 0.55); ^1H and ^{13}C NMR (CD_3OD) of the glycosidic part: see Table 2, ^{13}C NMR chemical shift values of the aglycone are identical to those described for saponin **2** to ± 0.3 ppm; ESI-MS (negative ion mode) m/z 967 $[\text{M}-\text{H}]^-$; ESI-MS-MS: MS^2 (967) m/z 817 $[(\text{M}-\text{H})-132-\text{H}_2\text{O}]^-$, 655 $[(\text{M}-\text{H})-132-162-\text{H}_2\text{O}]^-$, 611 $[(\text{M}-\text{H})-132-162-\text{CO}_2]^-$, MS^4 (655) m/z 595 $[(\text{M}-\text{H})-132-\text{H}_2\text{O}-\text{CH}_3\text{CO}_2\text{H}]^-$, 455 $[\text{genin}-\text{H}]^-$; ESI-MS (positive ion mode) m/z 1013 $[\text{M}+2\text{Na}-\text{H}]^+$; ESI-MS-MS : MS^2 (1013) m/z 953 $[(\text{M}+2\text{Na}-\text{H})-\text{CH}_3\text{CO}_2\text{H}]^+$, 881 $[(\text{M}+2\text{Na}-\text{H})-132]^+$, 851 $[(\text{M}+2\text{Na}-\text{H})-162]^+$, MS^3 (851) m/z 791 $[(\text{M}+2\text{Na}-\text{H})-162-$

$\text{CH}_3\text{CO}_2\text{H}]^+$, 719 $[(\text{M}+2\text{Na}-\text{H})-132-162]^+$, MS^4 (791) m/z 773 $[(\text{M}+2\text{Na}-\text{H})-162-\text{CH}_3\text{CO}_2\text{H}-\text{H}_2\text{O}]^+$, 659 $[(\text{M}+2\text{Na}-\text{H})-162-132-\text{CH}_3\text{CO}_2\text{H}]^+$, 501 $[\text{genin}+2\text{Na}-\text{H}]^+$.

3.9. saponin 8

White powder $[\alpha]_{\text{D}}^{21} +1.4$ (Pyridine, c 0.416); ^1H and ^{13}C NMR (CD_3OD) of the glycosidic part: see Table 2, ^{13}C NMR chemical shift values of the aglycone are identical to those described for saponin 2 to ± 0.3 ppm; ESI-MS (negative ion mode) m/z 1099 $[\text{M}-\text{H}]^-$, 743 $[(\text{M}-\text{H})-132-162-\text{CH}_3\text{CO}_2\text{H}-2\text{H}]^-$; ESI-MS (positive ion mode) m/z 1145 $[\text{M}+2\text{Na}-\text{H}]^+$; ESI-MS-MS: MS^2 (1145) m/z 995 $[(\text{M}+2\text{Na}-\text{H})-132-\text{H}_2\text{O}]^+$, 983 $[(\text{M}+2\text{Na}-\text{H})-162]^+$, 935 $[(\text{M}+2\text{Na}-\text{H})-132-\text{H}_2\text{O}-\text{CH}_3\text{CO}_2\text{H}]^+$, 869 $[(\text{M}+2\text{Na}-\text{H})-132-\text{H}_2\text{O}-\text{CO}_2\text{Na}-\text{CH}_3\text{CO}_2\text{H}+\text{H}]^+$, 983 $[(\text{M}+2\text{Na}-\text{H})-132-162-\text{H}_2\text{O}-\text{CO}_2\text{Na}+\text{H}]^+$, 729 $[(\text{M}+2\text{Na}-\text{H})-132-162-\text{H}_2\text{O}-\text{CO}_2\text{Na}-\text{CH}_3\text{CO}_2\text{H}+\text{H}]^+$, MS^3 (869) m/z 737 $[(\text{M}+2\text{Na})-132-162-\text{H}_2\text{O}-\text{CO}_2\text{Na}-\text{CH}_3\text{CO}_2\text{H}+\text{H}]^+$.

3.10. saponin 9

White powder $[\alpha]_{\text{D}}^{21} +7.7$ (Pyridine, c 0.21); ^1H and ^{13}C NMR ($\text{C}_5\text{D}_5\text{N}$) of the glycosidic part: see Table 2; ^{13}C NMR chemical shift values of the aglycone are identical to those described in the literature (Tan et al., 1999); ESI-MS (negative ion mode) m/z 1129 $[\text{M}-\text{H}]^-$; ESI-MS-MS: MS^2 (1129) m/z 979 $[(\text{M}-\text{H})-132-\text{H}_2\text{O}]^-$, 919 $[(\text{M}-\text{H})-132-\text{H}_2\text{O}-\text{CH}_3\text{CO}_2\text{H}]^-$, 773 $[(\text{M}-\text{H})-132-162-\text{CO}_2-\text{H}_2\text{O}]^-$, 713 $[(\text{M}-\text{H})-132-162-\text{CO}_2-\text{H}_2\text{O}-\text{CH}_3\text{CO}_2\text{H}]^-$, MS^3 (919) m/z 757 $[(\text{M}-\text{H})-132-\text{H}_2\text{O}-\text{CH}_3\text{CO}_2\text{H}-162]^-$; ESI-MS (positive ion mode) m/z 1175 $[\text{M}+2\text{Na}-\text{H}]^+$, 1153 $[\text{M}+\text{Na}]^+$; ESI-MS-MS : MS^2 (1153) m/z 1021 $[(\text{M}+\text{Na})-132]^+$, 991 $[(\text{M}+\text{Na})-162]^+$, 965 $[(\text{M}+2\text{Na}-\text{H})-132-\text{H}_2\text{O}-\text{CH}_3\text{CO}_2\text{H}]^+$, 899 $[(\text{M}+2\text{Na}-\text{H})-132-\text{H}_2\text{O}-\text{CH}_3\text{CO}_2\text{H}-\text{CO}_2\text{Na}+\text{H}]^+$ MS^3 (899) m/z 737 $[(\text{M}+2\text{Na}-\text{H})-132-162-\text{H}_2\text{O}-\text{CH}_3\text{CO}_2\text{H}-\text{CO}_2\text{Na}+\text{H}]^+$, MS^4 (737) m/z 575 $[(\text{M}+2\text{Na}-\text{H})-132-162-162-\text{H}_2\text{O}-\text{CH}_3\text{CO}_2\text{H}-\text{CO}_2\text{Na}+\text{H}]^+$.

3.11. saponin 10

White powder $[\alpha]_D^{21}$ -10.8 (MeOH, *c* 0.61); ^1H and ^{13}C NMR (CD_3OD) of the glycosidic part: see Table 2, ^{13}C NMR chemical shift values of the aglycone are identical to those described for saponin 2 to ± 0.3 ppm; ESI-MS (negative ion mode) m/z 1129 $[\text{M}-\text{H}]^-$, 773 $[(\text{M}-\text{H})-132-162-\text{CO}_2-\text{H}_2\text{O}]^-$; ESI-MS (positive ion mode) m/z 1153 $[\text{M}+\text{Na}]^+$; ESI-MS-MS : MS^2 (1153) m/z 1021 $[(\text{M}+\text{Na})-132]^+$, 991 $[(\text{M}+\text{Na})-162]^+$, 535 $[\text{triglycosidic chain}+\text{CH}_3\text{CO}_2\text{H}+\text{Na}]^+$, MS^3 (535) m/z 475 $[(\text{triglycosidic chain}+\text{Na})]^+$.

3.12. saponin II

$[\alpha]_D^{21}$ -8.1 (MeOH, *c* 0.28); ^1H and ^{13}C NMR (CD_3OD) of the glycosidic part: see Table 1, ^{13}C NMR (CD_3OD) of the aglycone δ 14.1 (C-27), 15.1 (C-26), 15.3 (C-24), 15.7 (C-25), 17.7 (C-6), 19.7 (C-11), 25.6 (C-2), 25.7 (C-12), 26.7 (C-23), 27.9 (C-30), 29 (C-15), 29.4 (C-29), 31.5 (C-20), 32.6 (C-16), 33 (C-21), 33 (C-22), 34.3 (C-7), 36.6 (C-10), 38.5 (C-1), 38.6 (C-4), 40.5 (C-8), 40.8 (C-13), 42.2 (C-14), 48.4 (C-17), 51.1 (C-9), 55.6 (C-5), 89.7 (C-3), 132.5 (C-19), 137 (C-18), 175.4 (C-28); ESI-MS (negative ion mode) m/z 967 $[\text{M}-\text{H}]^-$; ESI-MS-MS: MS^2 (967) m/z 817 $[(\text{M}-\text{H})-132-\text{H}_2\text{O}]^-$, 655 $[(\text{M}-\text{H})-132-162-\text{H}_2\text{O}]^-$; ESI-MS (positive ion mode) m/z 1013 $[\text{M}+2\text{Na}-\text{H}]^+$; ESI-MS-MS : MS^2 (1013) m/z 953 $[(\text{M}+2\text{Na}-\text{H})-\text{CH}_3\text{CO}_2\text{H}]^+$, 851 $[(\text{M}+2\text{Na}-\text{H})-162]^+$, MS^3 (851) m/z 833 $[(\text{M}+2\text{Na}-\text{H})-162-\text{H}_2\text{O}]^+$, 807 $[(\text{M}+2\text{Na}-\text{H})-162-\text{CO}_2]^+$, 791 $[(\text{M}+2\text{Na}-\text{H})-162-\text{CH}_3\text{CO}_2\text{H}]^+$, 773 $[(\text{M}+2\text{Na}-\text{H})-162-\text{CH}_3\text{CO}_2\text{H}-\text{H}_2\text{O}]^+$, 719 $[(\text{M}+2\text{Na}-\text{H})-132-162]^+$, MS^4 (791) m/z 659 $[(\text{M}+2\text{Na}-\text{H})-162-132-\text{CH}_3\text{CO}_2\text{H}]^+$, 640 $[(\text{M}+2\text{Na}-\text{H})-162-132-\text{CH}_3\text{CO}_2\text{H}-\text{H}_2\text{O}]^+$.

3.13. Acid hydrolysis of saponins

The crude saponin mixt. A (1g) was refluxed with 30 ml of 2N HCl for 4 h 30. The sapogenin mixture was extracted with EtOAc (3 \times 15 ml), washed with H_2O , dried over Na_2SO_4 and evapd to dryness. Oleanolic acid was identified from the sapogenin residue with an authentic sample by TLC in $\text{CHCl}_3/\text{MeOH}$ (98/2). The acid aq. layer was neutralised with 0.5 M KOH and freeze-dried. Three sugars were identified with authentic samples by TLC in

MeCOEt/iso-PrOH/Me₂CO/H₂O (20/10/7/6) as glucuronic acid, glucose, xylose and arabinose. After prep. TLC of the sugar mixt. in this solvent, the optical rotation of each purified sugar was measured. [valeurs??]

3.14. Haemolytic activity

This assay was performed as described previously (Voutquenne et al., 2002). Sheep erythrocyte suspension (10%) was obtained by dilution of a commercial 50% suspension from Biomerieux[®] Lyon with phosphate buffer saline (PBS). Saponin mixt. was dissolved in PBS. 25 µl of erythrocyte suspension were added to 1 ml of the sample and rapidly stirred. Absorbance of the supernatant was measured at 540 nm after 60 min of incubation and centrifuged at 3000 rpm for 5 min. HD₅₀ and HD₁₀₀ were the concentrations of sample which cause 50% and 100% of heamolysis, respectively.


Acknowledgment: The authors are grateful to Dr. T. Sévenet, Institut de Chimie des Substances Naturelles CNRS (Gif sur Yvette) for [the Mass Spectrometry facility and for providing plant material.](#)

References

- Ahmad, V. U., Abbasi, M. A., Hussain, H., Akhtar, M. N., Farooq, U., Fatima, N., Choudhary, M. I., 2003. Phenolic glycosides from *Symplocos racemosa*: natural inhibitors of phosphodiesterase I. *Phytochemistry* 63, 217-220.
- Ali, M., Bhutani, K. K., Srivastava, T. N., 1990. Triterpenoids from *Symplocos racemosa* bark. *Phytochemistry* 29, 3601-3604.
- Annaev, C., Isamukhamedova, M., Abubakirov, N. K., 1983a. Triterpene glycosides of *Salsola micranthera*. I. Structure of salsolosides C and D. *Khim. Prir. Soedin.* 6, 727-732, *Chem. Nat. Compd. (Engl. Transl.)* 19, 691-695.
- Annaev, C., Isaev, M. I., Abubakirov, N. K., 1983b. Triterpene glycosides from *Climacoptera transoxana*. III. Structure of copteroside E and F. *Khim. Prir. Soedin.* 5, 596-601, *Chem. Nat. Compd. (Engl. Transl.)* 19, 560-564.
- Annaev, C., Isamukhamedova, M., Abubakirov, N. K., 1984. Triterpene glycosides of *Salsola micranthera*. II. Structure of salsoloside E. *Khim. Prir. Soedin.* 20, 65-69; *Chem. Nat. Compd. (Engl. Transl.)* 20, 60-64.
- Casabueno, A. C., Pomilio, A. B., 1994. Lignans and a stilbene from *Festuca argentina*. *Phytochemistry* 35, 479-483.
- Cerda-Garcia-Rojas, C.M., Zamorano, G., Chavez, M.I., Catalan, C.A.N., Joseph-Nathan, P., 2000. ¹H and ¹³C NMR study of copteroside E derivatives. *Magn. Reson. Chem.* 38, 494-499.
- Cuenca, M. D. R., Catalan, C. A. N., 1991. Monoterpenes and lignans from *Mikania saltensis*. *J. Nat. Prod.* 54, 1162-1164.
- Dhaon, R., Jain, G. K., Sarin, J. P. S., Khanna, N. M., 1989. Symposide (I) a new anti-fibrinolytic glycoside from *Symplocos racemosa* Roxb. *Indian J. Chem.* 28B, 982-983.

- Frotan, M. H., Acharya, S. B., Frotan, R., Pathak, N. K. R., Biswas, M., 1983. Pharmacological investigations on alpha spinasterol isolated from *Symplocos spicata*. Indian J. Pharmacol. 15, 197-201.
- Hegnauer, R., 1973. Symplocaceae. In Hegnauer, R. (Ed), Chemotaxonomie der Pflanzen, VI, Birkhäuser Verlag, Basel and stuttgart, pp. 478-481.
- Higuchi, R., Kawasaki, T., Biswas, M., Pandey, V. B., Dasgupta, B., 1982. Triterpenoid saponins from the stem bark of *Symplocos spicata*. Phytochemistry 21, 907-910.
- Iida, J., Hayashi, M., Murata, T., Ono, M., Inoue, K., Fujita, T., 1990. Identification of a new minor iridoid glycoside in *Symplocos glauca* by thermospray liquid chromatography-mass spectrometry. J. Chromatogr. 515 503-508.
- Ishida, J., Wang, H-K., Oyama, M., Cosentino, M. L., Hu, C-Q., Lee, K-H., 2001. Anti-AIDS agents. 46.¹ Anti-HIV activity of Harman, an anti-HIV principle from *Symplocos setchuensis*, and its derivatives. J. Nat. Prod. 64, 958-960.
- Kawai, H., Nishida, M., Tashiro, Y., Kuroyanagi, M., Ueno, A., Satake, M., 1989. Studies on the structures of udosaponins A, B, C, D, E and F from *Aralia cordata* Thunb. Chem. Pharm. Bull. 37, 2318-2321.
- Khan, M.R., Kihara, M., Omoloso, A.D., 2001. Antimicrobial activity of *Symplocos cochinchinensis*. Fitoterapia 72, 825-828.
- Kim, J.S., Kang, S.S., 1998. Saponins from the arial parts of *Aralia continentalis*. Nat. Prod. Sci. 4, 45-50.
- Lin, L-C., Tsai, W-J., Chou, C-J., 1996. Studies on the constituents of *Symplocos lancifolia*. Chin. Pharm. J. 48, 441-449.
- Nishibe, S., Tsukamoto, H., Hisada, S., 1984. Effects of *O*-methylation and *O*-glucosylation on carbon-13 nuclear magnetic chemical shifts of matairesinol, (+)-pinoresinol and (+)-epipinoresinol. Chem. Pharm. Bull. 32, 4653-4657.

- Penders, A., Delaude, C., 1994. Triterpenoid saponins from *Melanthera scandens*.
Phytochemistry 37, 821-825.
- Tan, N., Zhou, J., Zhao, S., 1999. Advances in structural elucidation of glucuronide oleanane-type triterpene carboxylic acid 3,28-*O*-bisdesmosides (1962-1997). Phytochemistry 52, 153-192.
- Tanaka, T., Yamasaki, K., Kohda, H., Tanaka, O., Mahato, S. B., 1980. Dihydrochalcone-glucosides as sweet principles of *Symplocos* spp. Planta Med. (Suppl.), 81-83.
- Tanaka, T., Kawamura, K., Kohda, H., Yamasaki, K., Tanaka, O., 1982. Glycosides of the leaves of *Symplocos* spp. (Symplocaceae). Chem. Pharm. Bull. 30, 2421-2423.
- Tschesche, R., Braun, T. M., Sassen, W. V., 1980. Symplocoside, a flavonol glycoside from *Symplocos uniflora*. Phytochemistry 19, 1825-1829.
- Voutquenne, L., Lavaud, C., Massiot, G., Le Men-Olivier, L., 2002. Structure-activity relationships of haemolytic saponins. Pharm. Biol. 40, 253-262.
- Yoshikawa, M., Shimada, H., Morikawa, T., Yoshizumi, S., Matsumura, N., Murakami, T., Matsuda, H., Hori, K., Yamahara, Johji, 1997. Medicinal foodstuffs. VII.⁽¹⁾ on the saponin constituents with glucose and alcohol absorption-inhibitory activity from a food garnish "Tonburi", the fruit of Japanese *Kochia scoparia* (L.) Schrad.: Structure of scoparianosides A, B, and C. Chem. Pharm. Bull. 45, 1300-1305.



	R₁	R₂	R₃
1		H	H
2			H
3		H	
4			
5		H	H
6		H	
7		H	
8			
9			
10			




Table 1
¹H and ¹³C NMR data of saponins 2, 3, 4, 5 and 11 in CD₃OD

Genin	2		3		4		5		11	
	δ_H	δ_C	δ_H	δ_C	δ_H	δ_C	δ_H	δ_C	δ_H	δ_C
3	3.22 <i>dd</i> (11.5, 4.5)	89.9	3.17 <i>dd</i> (11.7, 4.2)	89.9	3.20 <i>dd</i> (11.8, 3.7)	89.8	3.19 <i>dd</i> (11.4, 4.2)	89.8	3.16 <i>dd</i> (12, 5)	89.7
12 ax	5.27 <i>t</i> (3.7)	122.3	5.27 <i>t</i> (3.4)	122.3	5.27 <i>t</i> (3.6)	122.3	5.27 <i>t</i> (3.4)	122.3	1.6 <i>m</i>	25.7
12 eq	—	—	—	—	—	—	—	—	1.26 <i>m</i>	—
13	—	143.4	—	143.4	—	143.4	—	143.4	2.29 <i>dd</i> (11.4, 2.5)	40.8
18	2.87 <i>dd</i> (13.7, 4)	41.1	2.87 <i>dd</i> (13.5, 3.4)	41.1	2.87 <i>dd</i> (13.6, 4.5)	41.1	2.87 <i>dd</i> (13.7, 4.4)	41.1	—	137
19 ax	1.73 <i>t</i> (13.5)	45.8	1.73 <i>t</i> (13.5)	45.8	1.73 <i>t</i> (13.5)	45.8	1.73 <i>t</i> (13.5)	45.8	5.16 <i>s</i>	132.5
19 eq	1.17 <i>t</i> (13.5)	—	1.17 <i>brd</i> (13.5)	—	1.17 <i>brd</i> (13.5)	—	1.17 <i>brd</i> (13.5)	—	—	—
23	1.06 <i>s</i>	27	0.97 <i>s</i>	27	0.97 <i>s</i>	26.9	1.07 <i>s</i>	27	0.95 <i>s</i>	26.7
24	0.86 <i>s</i>	15.5	0.75 <i>s</i>	15.5	0.75 <i>s</i>	15.4	0.87 <i>s</i>	15.5	0.73 <i>s</i>	15.3
25	0.97 <i>s</i>	14.5	0.95 <i>s</i>	14.5	0.96 <i>s</i>	14.5	0.97 <i>s</i>	14.5	0.91 <i>s</i>	15.7
26	0.82 <i>s</i>	16.3	0.81 <i>s</i>	16.3	0.81 <i>s</i>	16.2	0.82 <i>s</i>	16.2	1.03 <i>s</i>	15.1
27	1.18 <i>s</i>	24.8	1.17 <i>s</i>	24.9	1.17 <i>s</i>	24.8	1.18 <i>s</i>	24.8	0.83 <i>s</i>	14.1
28	—	176.6	-	176.6	-	176.6	-	176.6	-	175.4
29	0.93 <i>s</i>	32	0.93 <i>s</i>	32	0.93 <i>s</i>	32	0.93 <i>s</i>	32	1.01 <i>s</i>	29.4
30	0.95 <i>s</i>	22.5	0.96 <i>s</i>	22.5	0.95 <i>s</i>	22.5	0.95 <i>s</i>	22.5	0.99 <i>s</i>	27.9
3- β -D-glucuronic acid										
1'	4.51 <i>d</i> (7.8)	105.2	4.63 <i>d</i> (8)	102.9	4.74 <i>d</i> (8.1)	102.6	4.44 <i>d</i> (7.8)	105.4	4.6 <i>d</i> (7.9)	102.9
2'	3.41 <i>dd</i> (9.4, 7.8)	71.9	4.83 <i>dd</i> (9.5, 8.1)	73.4	4.9 <i>t</i> (9.8)	71.7	3.31 <i>dd</i> (7.9, 9.3)	73.5	4.83 <i>t</i> (8.6)	73.4
3'	5.03 <i>t</i> (9.4)	75.1	3.71 <i>t</i> (9.4)	72.4	5.19 <i>t</i> (9.5)	73.1	3.54 <i>t</i> (9)	74.5	3.69 <i>t</i> (8.4)	72.5
4'	3.86 <i>t</i> (9.4)	77.2	3.82 <i>t</i> (9.5)	79.6	3.94 <i>brt</i> (9.6)	78	3.72 <i>t</i> (9)	79.7	3.82 <i>dd</i> (9.3, 8.4)	79.8
5'	4.03 <i>brd</i> (9.2)	73.4	4.03 <i>brd</i> (9.6)	73.4	4.05 <i>brs</i>	nd	3.98 <i>brd</i> (9.2)	73.5	nd	nd
6'	—	171.1	—	170	—	171.2	—	nd	—	170
28- β -D-glucose										
1''	5.4 <i>d</i> (8.1)	94.2	5.4 <i>d</i> (8.2)	94.2	5.4 <i>d</i> (8.1)	94.2	5.4 <i>d</i> (8.2)	94.2	5.53 <i>d</i> (8.2)	94.3
2''	3.34 <i>t</i> (8.6)	72.4	3.33 <i>m</i>	72.7	3.33 <i>dd</i> (9.1, 8.1)	72.4	3.34 <i>dd</i> (9, 8)	72.5	3.32 <i>t</i> (8.7)	72.6
3''	3.43 <i>t</i> (8.4)	76.8	3.43 <i>t</i> (9)	76.8	3.42 <i>t</i> (9.1)	76.8	3.41 <i>t</i> (9)	76.8	3.44 <i>t</i> (8.9)	77
4''	3.37 <i>m</i>	69.6	3.39 <i>t</i> (9.5)	69.6	3.37 <i>t</i> (9.1)	69.6	3.38 <i>m</i>	69.6	3.41 <i>m</i>	69.6
5''	3.37 <i>m</i>	77.3	3.37 <i>m</i>	77.2	3.36 <i>m</i>	77.2	3.37 <i>m</i>	77.2	3.40 <i>m</i>	77.3
6a''	3.7 <i>dd-like</i> (12.1, 4.5)	60.9	3.70 <i>dd</i> (12, 4.2)	60.9	3.7 <i>dd</i> (12.2, 4.6)	60.9	3.7 <i>dd-like</i> (12.1, 4.3)	60.9	3.72 <i>dd</i> (11.9, 3.9)	61
6b''	3.84 <i>dd</i> (12.1, 1.4)	—	3.83 <i>dd</i> (12, 1.4)	—	3.84 <i>dd</i> (12.2, 1.7)	—	3.84 <i>dd</i> (12.1, 1.6)	—	3.85 <i>dd</i> (11.9, 1.7)	—
4'- β -D-xylose										
1'''	4.28 <i>d</i> (7.4)	103.8	4.4 <i>d</i> (7.5)	103.4	4.31 <i>brd</i> (5.2)	104.1	4.35 <i>d</i> (6.7)	103.4	4.41 <i>brd</i> (7.5)	103.5
2'''	3.06 <i>dd</i> (9, 7.4)	73.4	3.22 <i>dd</i> (9, 7.5)	73.1	3.1 <i>brt</i> (8)	73.5	3.57 <i>dd</i> (9, 6.7)	70.9	3.23 <i>brdd</i> (9.5, 7.5)	73.2
3'''	3.27 <i>t</i> (9)	76.1	3.33 <i>t</i> 9.4	75.9	3.28 <i>dd</i> (9.5, 8.5)	76.1	3.54 <i>dd</i> (9.2, 3.3)	72.5	3.33 <i>t</i> (9.8)	76
4'''	3.45 <i>ddd</i> (10, 9, 5.3)	69.6	3.51 <i>ddd</i> (9.9, 8.9, 5.3)	69.4	3.45 <i>ddd</i> (9.9, 9, 5)	69.6	3.83 <i>brs</i>	68.1	3.52 <i>ddd</i> (11.4, 10.4, 5.3)	69.4
5ax'''	3.17 <i>dd</i> (11.5, 10)	65.6	3.26 <i>dd</i> (11.5, 10)	65.5	3.17 <i>t</i> (11)	65.7	3.62 <i>dd</i> (12.5, 1.3)	65.9	3.26 <i>dd</i> (11.4, 10.4)	65.5
5eq'''	3.83 <i>dd</i> (11.5, 5.2)	—	3.93 <i>dd</i> (5.3, 11.5)	—	3.83 <i>dd</i> (11.4, 5)	—	3.95 <i>dd</i> (12.5, 2.8)	—	3.93 <i>dd</i> (11.7, 5.3)	—
2'-acetate										
CH ₃	—	—	2.09 <i>s</i>	19.7	2.02 <i>s</i>	19.6	—	—	2.08 <i>s</i>	19.7
C=O	—	—	—	170	—	169.6	—	—	—	170
3'-acetate										
CH ₃	2.08 <i>s</i>	19.8	—	—	2.01 <i>s</i>	19.4	—	—	—	—
C=O	—	171.1	—	—	—	170.6	—	—	—	—

nd = not determined.

Table 2
¹H and ¹³C NMR data of saponins 6, 8, 9 and 10 in CD₃OD

Genin	6		8		9 ^a		10	
	δ _H	δ _C	δ _H	δ _C	δ _H	δ _C	δ _H	δ _C
3	3.14 <i>dd</i> (11.7, 4.2)	89.8	3.17 <i>dd</i> (11.7, 4.2)	90.5	3.11 <i>dd</i> (12.3, 4.3)	89.7	3.24 <i>dd</i> (12, 4.4)	91
12	5.23 <i>t</i> (3.4)	122.3	5.27 <i>t</i> (3.6)	122.3	5.38 <i>t</i> (3.5)	123.6	5.27 <i>t</i> (4.6)	122.3
13	—	143.4	—	143.4	—	143.8	—	143.4
18	2.84 <i>dd</i> (13.4, 3.8)	41.1	2.87 <i>dd</i> (13.6, 4.1)	41.1	3.6 <i>dd</i> (13.6, 4)	41.4	2.88 <i>dd</i> (13.5, 4)	41.1
23	0.94 <i>s</i>	26.9	1.06 <i>s</i>	26.8	1.09 <i>s</i>	27.5	1.07 <i>s</i>	26.8
24	0.72 <i>s</i>	15.4	0.86 <i>s</i>	15.1	1.04 <i>s</i>	16.3	0.88 <i>s</i>	15.4
25	0.93 <i>s</i>	14.5	0.97 <i>s</i>	14.6	0.74 <i>s</i>	15.1	0.98 <i>s</i>	14.5
26	0.78 <i>s</i>	16.2	0.81 <i>s</i>	16.3	1.04 <i>s</i>	17.1	0.82 <i>s</i>	16.3
27	1.14 <i>s</i>	24.8	1.18 <i>s</i>	24.9	1.23 <i>s</i>	25.8	1.18 <i>s</i>	24.9
28	—	176.6	—	176.6	—	176.2	—	176.6
29	0.9 <i>s</i>	32	0.93 <i>s</i>	32.1	0.88 <i>s</i>	32.8	0.93 <i>s</i>	32
30	0.92 <i>s</i>	22.5	0.96 <i>s</i>	22.5	0.85 <i>s</i>	23.3	0.95 <i>s</i>	22.5
3-β-D-glucuronic acid								
1'	4.61 <i>d</i> (8)	102.9	4.65 <i>d</i> (7.3)	103.8	4.85 <i>d</i> (7.5)	104.7	4.69 <i>d</i> (7.4)	103.7
2'	4.8 <i>dd</i> (9.4, 8.1)	73.3	3.71 <i>dd</i> (9, 7.3)	77.2	4.42 <i>dd</i> (9.2, 7.3)	77.2	3.86 <i>dd</i> (9.2, 7.6)	75.9
3'	3.7 <i>t</i> (9.3)	72.4	5.2 <i>t</i> (9)	74.5	5.86 <i>t</i> (9)	74.9	5.27 <i>t</i> (9.2)	74.6
4'	3.78 <i>t</i> (9.3)	79.7	3.94 <i>t</i> (9.3)	77.1	4.54 <i>m</i>	78.9	3.9 <i>t</i> (9.2)	77.2
5'	4.01 <i>brd</i> (9.2)	na	4.09 <i>d</i> (9.3)	73.6	4.55 <i>m</i>	75.7	4.08 <i>brd</i> (9)	nd
6'	—	170	—	171.1	—	170.1	—	171.1
28-β-D-glucose								
1''	5.37 <i>d</i> (8.2)	94.2	5.4 <i>d</i> (8.2)	94.2	6.31 <i>d</i> (8.1)	95.5	5.41 <i>d</i> (8.1)	94.2
2''	3.30 <i>t</i> (8.4)	72.4	3.33 <i>t</i> (8.4)	72.4	4.21 <i>t</i> (8.8)	73.8	3.34 <i>t</i> (8.4)	72.5
3''	3.40 <i>t</i> (8.8)	76.8	3.42 <i>t</i> (9.2)	76.8	4.29 <i>t</i> (8.7)	78.6	3.44 <i>t</i> (8.9)	76.8
4''	3.36 <i>t</i> (8.8)	69.6	3.38 <i>m</i>	69.7	4.36 <i>t</i> (8.8)	70.7	3.38 <i>m</i>	69.6
5''	3.33 <i>m</i>	77.2	3.37 <i>m</i>	77.2	4.03 <i>m</i>	79.1	3.37 <i>m</i>	77.2
6a''	3.67 <i>dd</i> (11.9, 4.3)	60.9	3.7 <i>dd</i> (12.2, 4.1)	60.9	4.4 <i>dd</i> (12.1, 4.3)	61.9	3.7 <i>dd</i> (11.9, 5.3)	60.9
6b''	3.8 <i>dd</i> (11.9, 1.4)	60.9	3.84 <i>dd</i> (12.2, 1.9)		4.47 <i>dd</i> (12.1, 2.3)		3.84 <i>dd</i> (11.9, 1.9)	
1'''			2'-β-D-xylose	104	2'-β-D-glucose		2'-β-D-glucose	
2'''			4.37 <i>d</i> (7.6)	104	5.13 <i>d</i> (7.6)	104.3	4.46 <i>d</i> (7.7)	102.8
3'''			3.13 <i>dd</i> (9.1, 7.4)	73.6	3.96 <i>dd</i> (8.6, 7.6)	75.1	3.14 <i>dd</i> (9.2, 7.7)	73.7
4'''			3.28 <i>t</i> (9)	76.4	4.22 <i>t</i> (9)	78.1	3.33 <i>t</i> (9.2)	76.4
5ax'''			3.44 <i>m</i>	69.6	4.12 <i>t</i> (8.9)	72	3.11 <i>t</i> (9.3)	70.7
5eq'''/6a'''			3.13 <i>t</i> (11)	65.6	4.03 <i>m</i>	78.2	3.27 <i>m</i>	77.1
6b'''			3.82 <i>dd</i> (11.4, 5.4)		4.37 <i>dd</i> (11.5, 5.5)	63.2	3.58 <i>dd</i> (12, 7.5)	62.1
					4.62 <i>dd</i> (11.5, 2.5)		3.86 <i>dd</i> (12, 1.9)	
	4'-α-L-arabinopyranose		4'-β-D-xylose		4'-β-D-xylose		4'-α-L-arabinofuranose	
1'''	4.34 <i>d</i> (6.6)	103.3	4.29 <i>d</i> (7.5)	103.8	5.03 <i>d</i> (7.5)	105.8	4.96 <i>brs</i>	109.4
2'''	3.53 <i>brt</i> (9)	70.9	3.08 <i>dd</i> (9, 7.5)	73.4	3.92 <i>dd</i> (8.3, 7.5)	74.7	3.9 <i>m</i>	81.7
3'''	3.51 <i>dd</i> (9.1, 3.4)	72.5	3.27 <i>t</i> (9)	76.1	4.07 <i>t</i> (8.8)	77.9	3.83 <i>dd</i> (5.6, 3.4)	76.8
4'''	3.79 <i>brs</i>	68	3.44 <i>m</i>	69.6	4.15 <i>ddd</i> (10.9, 8.8, 5.6)	70.6	3.9 <i>m</i>	85
5ax'''	3.58 <i>dd</i> (12.6, 1.3)	65.8	3.17 <i>dd</i> (13, 11.2)	65.4	3.67 <i>t</i> (10.8)	67.1	3.63 <i>dd</i> (12, 4.8)	61.4
5eq'''	3.91 <i>dd</i> (12.6, 2.9)	65.8	3.84 <i>dd</i> (11.4, 5.1)	65.6	4.31 <i>dd</i> (10.8, 5)		3.71 <i>dd</i> (12, 3.6)	
	2'-acetate		3'-acetate		3'-acetate		3'-acetate	
CH ₃	2.06 <i>s</i>	19.7	2.09 <i>s</i>	20.1	2.2 <i>s</i>	21.3	2.18 <i>s</i>	20.4
C=O		170		171.1		170.1		171.1

na: not assigned; signal was too weak.

^a Measured in C₅D₅N.