



## **Type XIX collagen: A new partner in the interactions between tumor cells and their microenvironment.**

Jean-Baptiste Oudart, Jean-Claude Monboisse, François-Xavier Maquart, B Brassart, S Brassart-Pasco, L. Ramont

### **► To cite this version:**

Jean-Baptiste Oudart, Jean-Claude Monboisse, François-Xavier Maquart, B Brassart, S Brassart-Pasco, et al.. Type XIX collagen: A new partner in the interactions between tumor cells and their microenvironment.. Matrix Biology, 2017, 57 (58), pp.169-177. 10.1016/j.matbio.2016.07.010 . hal-02183231

**HAL Id: hal-02183231**

**<https://hal.univ-reims.fr/hal-02183231>**

Submitted on 4 Feb 2022

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

## **Matrix Biology**

### **Mini review**

#### **Type XIX Collagen: A new partner in the interactions between tumor cells and their microenvironment.**

Jean-Baptiste Oudart<sup>a,b</sup>, Jean-Claude Monboisse<sup>a,b</sup>, François-Xavier Maquart<sup>a,b</sup>, Bertrand Brassart<sup>a</sup>, Sylvie Brassart-Pasco<sup>a</sup>, and Laurent Ramont<sup>a,b,\*</sup>

#### **Author's Affiliations:**

<sup>a</sup> Université de Reims Champagne-Ardenne, CNRS UMR 7369 (Matrice Extracellulaire et Dynamique Cellulaire, MEDyC), 51095 Reims, France.

<sup>b</sup> CHU de Reims, Laboratoire Central de Biochimie, 51092 Reims, France.

**\* Corresponding** author : Dr Laurent Ramont, Université de Reims Champagne-Ardenne, CNRS UMR 7369 (Matrice Extracellulaire et Dynamique Cellulaire, MEDyC), rue Cognacq Jay, 51095 Reims, France. Phone: 0033-326783181; Fax: 0033-326788539; E-Mail: [lrumont@chu-reims.fr](mailto:lrumont@chu-reims.fr)

#### **Keywords:**

FACIT

Collagen XIX

NC1 domain

Matrikine

Tumor invasion

Integrin

Signal transduction

## ABSTRACT

Collagen XIX is a minor collagen associated with basement membranes zone, which belongs to the family of FACITs (Fibril Associated Collagens with Interrupted Triple Helices). The FACIT family is composed of collagens IX, XII, XIV, XVI, XX, XXI, XXII and XIX, which share many highly conserved structural motifs: short NC1 domain, a thrombospondin-like N-terminal domain (TspN), and numerous cysteine residues. The main role of FACITs is to ensure integrity and stability of the extracellular matrix and its fibrillar collagen network by regulating formation and size of collagen fibrils.

Collagen XIX was discovered in a human rhabdomyosarcoma cell line. The  $\alpha 1(\text{XIX})$  chain is composed of 5 collagenous domains interrupted by 6 non-collagenous domains with a short C-terminal 19 amino-acid residues non-collagenous domain (NC1). This collagen is involved in the differentiation of muscle cells, central nervous system development and esophagus formation.

Collagen XIX is associated with the basement membrane zone like collagen XVIII and XV. Its short NC1(XIX) C-terminal domain inhibits migration and invasion of melanoma cells. It also exerts a strong anti-angiogenic effect *via* an inhibition of MMP-14 and VEGF expression. NC1(XIX) binding to  $\alpha v\beta 3$  integrin decreases protein phosphorylation of proteins involved in the FAK (Focal Adhesion Kinase) / PI3K (Phosphoinositide 3-Kinase) / Akt (protein Kinase B) / mTOR (Mammalian Target Of Rapamycin) pathway. On the other hand, NC1(XIX) induces an increase in GSK3 $\beta$  activity by decreasing its degree of phosphorylation. The inhibition of this pathway could explain the anti-tumor properties of the NC1(XIX) domain.

## **Contents**

1. Introduction
2. FACIT family
3. Collagen XIX
  - 3.1 Discovery
  - 3.2 Chromosomal localization and molecular structure
  - 3.3 Cell and tissue expression
    - 3.3.1 Expression in mouse
    - 3.3.2 Expression in human
4. Physiopathology
  - 4.1 Collagen XIX in myogenic differentiation
  - 4.2 Collagen XIX in central nervous system
  - 4.3 Other potential implications of collagen XIX
5. Collagen XIX and tumor process
  - 5.1 Collagen of the basement membrane zone
  - 5.2 Collagen XIX in breast cancer
  - 5.3 The NC1domain of collagen XIX
  - 5.4 NC1(XIX) domain and melanoma
6. Conclusion
- References

## 1. Introduction

Collagen XIX is a minor collagen associated with basement membranes zone, which belongs to the FACITs family (Fibril Associated Collagens with Interrupted Triple Helices) (Myers et al., 1993; Yoshioka et al., 1992). Collagens are the most abundant proteins in mammals, with about 30% of the total protein weight. The collagen family is composed of 28 members numbered from I to XXVIII. Each collagen is composed of 3 polypeptide chains commonly referred as  $\alpha$  chain. Currently, 42 genes coding for 46 different  $\alpha$  chains have been characterized (Ricard-Blum, 2011; Ricard-Blum and Ruggiero, 2005). The lengths of the  $\alpha$  chains vary from 662 amino acids for  $\alpha 1(X)$  to 3152 amino acids for  $\alpha 3(VI)$ .

Some collagens are homotrimers composed of 3 identical  $\alpha$  chains (types II, III, VII, VIII, X, XII, XXVIII). The others are heterotrimers composed of different  $\alpha$  chains encoded by different genes or resulting from the alternative splicing of the same gene. The main characteristic of collagen is the presence of a triple helical structure resulting of three  $\alpha$  chain coiling. The triple helix may represent more than 90% of intact molecule (as in collagen I) or less than 10% (as in collagen XII). It is composed of repeating triplets Gly - X - Aaa where X is predominantly a proline and Aaa a 4-hydroxyproline (van der Rest and Garrone, 1991). The collagen family may be divided into different subfamilies according to their supramolecular assemblies and their functions (for recent review of this classification, see (Ricard-Blum, 2011). In the current review, we will focus on knowledge about collagen XIX: its molecular structure, expression, function and its more recently described anti-tumor properties.

## 2. FACIT family

Collagen XIX belongs to the FACIT family. By definition, triple helix of FACIT collagens is interrupted by non-collagenous domains. They differ by their  $\alpha$  chain composition and length, the number of their collagenous and non-collagenous domains (2 to 10). Nevertheless, the members of this family share many highly conserved structural motifs (Ricard-Blum and Ruggiero, 2005), as a short NC1 domain of a few tens of amino acids, a thrombospondin-like N-terminal domain (TspN), the presence of numerous cysteine residues, an especially highly conserved type Cys - (Xaa)<sub>4</sub> - Cys pattern at the junction of the Col1 and NC1 domains, and two exceptions to the repetition of the Gly - X - Aaa motif in the Col2 domain (Ricard-Blum, 2011; Ricard-Blum and Ruggiero, 2005). This family is composed of the collagens IX, XII, XIV, XVI, XX, XXI, XXII and XIX. The major role of FACITs is to ensure integrity and stability of the extracellular matrix and its fibrillar collagen network.

FACITs regulate formation and size of collagen fibrils ([Ivanova and Krivchenko, 2014](#)) but also the training and control of cell organization in the extracellular matrix ([Tao et al., 2012](#)). They possess domains promoting adhesion and interaction with collagen fibrils, a rigid domain which projects outside the fibrils and multiple domains for interactions with other components of the extracellular matrix. These interactions have been demonstrated in cartilage, where collagen IX binds to fibronectin ([Parsons et al., 2011](#)) and in the skin, where collagen XII interacts with collagen I and tenascin X ([Chiquet et al., 2014](#)). Moreover, FACIT collagens can interact with cells ([Inoguchi et al., 1995](#)). Considering the restrictive and specific expression of FACITs in some tissues, their fine regulation of expression during development and the serious functional alterations caused by their mutations, they are probably involved in specific functions in different organs. For example, during the embryonic period, collagen XIV is involved in the formation of interstitial myocardial tissues ([Tao et al., 2012](#)). Collagen XII is involved in the regulation of the organization and properties of the collagen fibers in the muscle and bone ([Chiquet et al., 2014](#)).

### **3. Collagen XIX**

#### *3.1 Discovery*

Collagen XIX was discovered in 1992 by [Yoshioka et al., 1992](#), in Marion Gordon's team. It was characterized by cDNA sequencing in a human rhabdomyosarcoma cell line (CCL-136, American Type Culture Collection). Because of its isolation in the HY-67 clone, this new collagen chain was at first called  $\alpha 1(Y)$  ([Yoshioka et al., 1992](#)). In 1993, Myers et al described in turn a new collagen that they called "RH COL" because of its isolation in human Rhabdomyosarcoma cell lines (RH) ([Myers et al., 1993](#)). In fact, these two discoveries were identical and the name collagen XIX was used to replace  $\alpha 1(Y)$  and RH COL terms in 1994 ([Myers et al., 1994](#)). Then, structural analysis were conducted on collagen XIX and allowed its classification in the FACIT family. ([Inoguchi et al., 1995](#); [Khaleduzzaman et al., 1997](#))

#### *3.2 Chromosomal localization and molecular structure*

The human COL19A1 gene, encoding the  $\alpha 1(XIX)$  chain, is mapped on human chromosome 6 in the q12-q14 region. Genes encoding the  $\alpha 1$  chain of two other FACIT collagens, collagen XII (COL12A1) ([Gerecke et al., 1997](#); [Oh et al., 1992](#)) and collagen IX (COL9A1) ([Kimura et al., 1989](#)) are also located in the same region. Collagen XIX gene comprises 250 kb and 51 exons. The exons encoding the NC1, Col1, NC2, Col2, NC3, Col3, NC4 domains of the  $\alpha 1(XIX)$  chain share high homology with those encoding the NC1, Col1, NC2, Col2, NC3 domains of the  $\alpha 2(IX)$  chain ([Khaleduzzaman et al., 1997](#)). Similarly, there

are significant homologies between the exons encoding the Col5 - NC6 domains of the  $\alpha 1(\text{XIX})$  chain and the exons encoding the Col3 and NC4 domains of the  $\alpha 1(\text{IX})$  chain. These data suggest the existence of an ancestral gene encoding the different FACIT  $\alpha$  chains (Khaleduzzaman et al., 1997). COL19A1 encodes a messenger RNA of 10.4 kb, with a long 3'-untranslated region of 5.3 kb (Inoguchi et al., 1995). There is 82% sequence homology between the human and murine  $\alpha 1(\text{XIX})$  chains. The strongest homologies are located at the C-terminal side (Sumiyoshi et al., 1997).

Collagen XIX is composed of three  $\alpha 1(\text{XIX})$  chains, forming a homotrimer of 400 kDa. Each  $\alpha$  chain consists of 1142 amino acids with a molecular mass of 112 kDa. The first 23 amino acids constitute the signal peptide (Inoguchi et al., 1995). The amino acid composition is comparable to other collagens, although collagen XIX is the richest in leucine (10.7%) (Nassa et al., 2012). Each chain contains 5 collagenous domains (COL1 to COL5) interrupted by 6 non-collagenous domains (NC1 to NC6), composed of 19, 44, 23, 31, 20, 291 amino acid residues, respectively (Inoguchi et al., 1995).

From a structural point of view, collagen XIX is close to another FACIT, collagen XVI. The NC2  $\alpha 1(\text{XIX})$  and NC4  $\alpha 1(\text{XIX})$  domains are similar to the corresponding domains of collagen XVI (Myers et al., 1994). The NC6  $\alpha 1(\text{XIX})$  domain shows very strong homologies with the NC4  $\alpha 1(\text{XII})$ , NC3  $\alpha 1(\text{XII})$ , NC3  $\alpha 1(\text{XIV})$  and NC11  $\alpha 1(\text{XVI})$  domains (Khaleduzzaman et al., 1997). NC6  $\alpha 1(\text{XIX})$  domain also shows very strong homologies with two domains of others proteins, the PARP (Proline Arginine Rich Peptide) domain of collagen V and XI and the N terminal module of thrombospondin (Khaleduzzaman et al., 1997). Phylogenetic analysis shows that collagen XIX also has similarities with collagens IX and XXII, which also belong to the FACIT family (Nassa et al., 2012). The NC2  $\alpha 1(\text{XIX})$  domain has a stable helix structure and is probably responsible for the trimerization of the  $\alpha 1(\text{XIX})$  chains (Boudko et al., 2008). This domain is also involved in the formation of disulfide inter-chain bonds at the junction between Col1 and NC1 domains (Boudko et al., 2008). The  $\alpha 1(\text{XIX})$  chain also contains 14 cysteine residues, 2 at the junction of Col1 and NC1 domains, 2 others in the NC4 domain and the 10 others in the NC6 domain (thrombospondin-like domain) (Inoguchi et al., 1995). The presence of cysteine promotes the formation of inter-chain and intra-chain disulfide bonds (Myers et al., 2003; Myers et al., 1994), allowing the formation and the stabilization of the triple helix. The presence of inter-chain disulfide bonds was confirmed by rotatory shadowing electron microscopy. In addition, electron microscopy showed that collagen XIX has a rod-like structure (240nm) with a globular domain at the N-terminal end (Myers et al., 2003). This technique also showed the presence of 45% collagen XIX monomers and 55% oligomers (dimers, trimers, tetramers, pentamers, hexamers and other multimers) connected and stabilized by N-terminus inter-molecular disulfide bonds (Myers et

al., 2003). Collagen XIX presents no glycosylation site (Inoguchi et al., 1995). However, it contains a binding site for heparin in the NC6 domain (Inoguchi et al., 1995).

The use of polyclonal anti-sera against the C- or N-terminal domains of collagen XIX, under reducing conditions, in a sample of human rhabdomyosarcoma cells, allowed to highlight a Mr 165 kDa band corresponding to the  $\alpha 1$ (XIX) chain. In non-reducing conditions, a band with Mr around 230 kDa was observed, which confirms the presence of disulfide inter-chain bonds (Myers et al., 1997). Experimentally, the 165 kDa band was sensitive to the action of bacterial collagenase, which released a fragment of 34 kDa. Moreover, amino-acid sequence analysis revealed a presumed cleavage site for bacterial collagenase capable of releasing a fragment of a potential molecular weight of 37 kDa, which corresponds to the fragment experimentally obtained (Myers et al., 1997).

Despite its structural similarities with other members of the FACIT family, collagen XIX has also been classified within the multiplexin family, with collagens XV and XVIII. Indeed, it is not associated with the fibers and possesses a thrombospondin-like N-terminal domain, like collagen XIII and XV. Moreover, immunohistochemistry localized collagen XIX in the basement membrane zone of different organs, like other multiplexin (Myers et al., 1997).

### 3.3 Cell and tissue expression

#### 3.3.1 Expression in mouse

The COL19A1 gene encoding the  $\alpha 1$  chain of XIX collagen is mapped on mouse chromosome 1 (Khaleduzzaman et al., 1997). Sumiyoshi *et al* have examined tissue expression of Col19A1 gene by RT-PCR during mouse embryogenesis and in adult mice. COL19A1 murine transcripts are detectable as soon as the 11<sup>th</sup> day of gestation in almost all organs studied (except liver). In adult mice, the tissue expression of the messenger RNA is restricted to the central nervous system (cerebrum and cerebellum), testis and eye, with traces in the aorta, lung or kidney. The central nervous system is the only organ where COL19A1 gene expression gradually increases after birth. It is ten fold higher in adult mice compared to the embryonic period (Sumiyoshi et al., 1997). The locations of the transcripts are summarized in Table 1 .

#### 3.3.2 Expression in Human

During the embryonic period, human collagen XIX has a broad tissue distribution. However, it represents only a tiny proportion of tissues, for example  $10^{-6}$  % of the weight of the umbilical cord (Myers et al., 1997). In adult, its tissue distribution is restricted to specific



areas, mainly vascular, neuronal and epithelial basement membranes ([Sumiyoshi et al., 1997](#)). Some biochemical and immunohistochemical studies have studied the distribution of collagen XIX in human organs ([Myers et al., 1997](#); [Su et al., 2010](#)) ([Table 2](#)).

#### 4. Physiopathology

Functions of collagen XIX currently remain unclear. Like other FACITs, it participates in the formation and maintenance of extracellular matrix, mainly during embryonic period ([Sumiyoshi et al., 1997](#)). However, many other functions of collagen XIX are probably yet to be discovered.

##### *4.1 Collagen XIX in myogenic differentiation*

*In situ* hybridization analysis of the transcripts of the murine COL19A1 gene in the muscle shows gene expression in the myotome as soon as the 9<sup>th</sup> day of gestation. This expression is superposed with that of the regulatory gene of muscle differentiation: myf-5 (myogenic factor 5). Afterwards, muscle expression of COL19A1 gene gradually decreases ([Sumiyoshi et al., 2001](#)). In other words, collagen XIX seems to play a role in the early stages of embryonic muscle differentiation. Rhabdomyosarcoma cell lines were used to study embryogenesis of skeletal muscle and differentiation of mesenchymal cells. Some cells of these lines produce collagen XIX and muscle proteins such as myosin and  $\alpha$ -actinin. Under culture conditions, collagen XIX and muscle protein production were greatly increased in a sub-population of cells and induced myoblast differentiation. This study suggests a role of collagen XIX in the early stages of muscle differentiation ([Myers et al., 1999](#)). This essential role of collagen XIX in muscle differentiation and physiology was also confirmed by Sumiyoshi *et al* ([Sumiyoshi et al., 2004](#)).

In mouse, the double invalidation of collagen XIX gene leads to a greater mortality than 95% in the first three weeks after birth. Animals die mainly of malnutrition. The autopsy reveals the presence of a mega-esophagus. The animals present a motor dysfunction of smooth muscles associated with an alteration of the transformation of smooth muscle cells in striated muscle cells, particularly in the abdominal portion of the esophagus. In addition, mice also exhibit a hypertonicity of their cardia. These abnormalities are related to a disruption of extracellular matrix and a lack of activation of myogenic regulatory factors. Heterozygous mice are viable, fertile and with normal morphology. By consequence, collagen XIX seems to be heavily involved in the stomach sphincter functions and muscle development, mainly esophageal ([Sumiyoshi et al., 2004](#)).

More recently, a study showed overexpression of collagen XIX in muscles of patients with amyotrophic lateral sclerosis ([Shtilbans et al., 2011](#)).

#### *4.2 Collagen XIX in central nervous system*

Collagen XIX is found in specific populations of hippocampal interneurons such as the interneurons expressing neuropeptide Y, somatostatin and calbindin. In a mouse model mutated for collagen XIX gene, Su et al showed that the morphology of the neurons was normal but that the formation of inhibitory synapses was diminished at the hippocampal level and in the subiculum, two parts of the temporal lobe largely involved in memory. Therefore, the expression of collagen XIX in interneurons appears to be essential for synapse formation in hippocampal tissue ([Su et al., 2010](#)).

In 2010, Hilario et al showed that collagen XIX had a role in axon guidance in a zebrafish model. During the embryonic period, axons of the neurons of spinal cord are progressing toward their target organ. Herein, axon growth cone is guided by various molecules that determine specific relations between various structures of the developing organism. Moreover, collagen XIX acts as a relay to guide these axons. This new feature appears to be concentration-dependent, permitting a regulation of axonal migration ([Hilario et al., 2010](#)).

#### *4.3 Other potential implications of collagen XIX.*

Due to its localization in vascular basement membranes of many organs, some authors have suggested that collagen XIX may also be involved in the angiogenesis process ([Myers et al., 1997](#)).

Arya *et al* have found strong evidence for a gene on chromosome 6q that influences variation in birth weight in Mexican and American populations. The gene of collagen XIX is in this region and is consequently a candidate gene among other for its involvement in low birth weight ([Arya et al., 2006](#)).

### **5. Collagen XIX and tumor progression**

#### *5.1 Collagens of the basement membrane zone*

Collagen XIX is associated to the basement membrane zone. Basement membranes are specialized extracellular matrix, present in all vertebrates and some invertebrates. They are located at the interface between epithelia or endothelia and connective tissue, at blood vessel wall, axons of peripheral nerves, adipocytes or muscle cells ([Paulsson, 1992](#)). They are amorphous and dense structures, which play a role of cell support ([Kalluri, 2003](#)). Apart

from this architectural role, basement membranes exert many important biological functions and play key roles in many physiological and pathological situations ([Kruegel and Miosge, 2010](#)). Basement membranes dynamically interact with cells *via* adapter proteins, mainly integrins, to determine the cell polarization, and regulate many biological functions such as adhesion, differentiation, proliferation, chemotaxis or cell migration ([Aumailley and Timpl, 1986](#); [Paulsson, 1992](#)). More recently, associations between basement membrane and minor collagens such as type XVIII, XV and XIX were reported and it is better now to talk about basement membrane zone ([Yurchenco, 2011](#); [Yurchenco and O'Rear, 1994](#)). Collagen XVIII and XV are the better known. They exert anti-tumor and anti-angiogenic properties, especially through their NC1 domains (Endostatin and Restin respectively) ([O'Reilly et al., 1997](#); [Ramchandran et al., 1999](#)). We and others have previously shown that NC1 domains of collagen IV exhibit anti-tumor and / or anti-angiogenic activity ([Thevenard et al., 2006](#); [Thevenard et al., 2010](#)). For example, tumstatin, the NC1 domain of  $\alpha 3(\text{IV})$  chain, tetrastatin, the NC1 domain of  $\alpha 4(\text{IV})$  chain, canstatin, the NC1 domain of  $\alpha 2(\text{IV})$  chain and arresten, the NC1 domain of  $\alpha 3(\text{IV})$  chain, have been described as having anti-tumor and / or anti-angiogenic properties in several experimental in vitro and in vivo models ([Floquet et al., 2004](#); [Pasco et al., 2005](#)). More recently, our group demonstrated the antitumor properties of the NC1 domain of collagen XIX ([Ramont et al., 2007](#); [Toubal et al., 2010](#)).

### *5.2 Collagen XIX in breast cancer*

Amenta *et al* studied the expression of collagens XV and XIX in breast cancer. During tumor invasion, an early reduction of collagen XIX in the epithelial basement membrane zone was found. This reduction occurred in the early stages of tumor progression, even before the complete destruction of the basement membrane. Therefore, this decrease may be involved in the extracellular matrix remodeling processes that precede tumor invasion ([Amenta et al., 2003](#)). Various hypotheses have been proposed to explain this decrease, such as an increased susceptibility of collagen XIX to proteolysis or a decrease in its synthesis.

### *5.3 The NC1 domain of collagen XIX*

The partial and controlled proteolysis of macromolecules of the extracellular matrix, results in the release of peptides with biological activities, called "matrikines" ([Maquart et al., 1999](#)) or "matricryptins" ([Davis et al., 2000](#)). Some matrikines derived from the proteolysis of basement membrane zone collagens, particularly the NC1 domains of collagens IV, XV, XVIII and XIX, exert anti-tumor and/or anti-angiogenic activities ([Monboisse et al., 2014](#)). Various enzymatic systems are involved in the proteolysis of extracellular matrix components, such as matrix metalloproteinases (MMPs) or the plasminogen-plasmin system.

The NC1 (XIX) domain located at the C-terminus of the  $\alpha 1$ (XIX) chain is very short. It consists of 19 amino acid residues with the following sequence: NPEDCLYPVSHAHQRTGGN. Collagen XIX and its fragments can be detected in cell culture supernatants, in tissue extracts and in many biological fluids such as amniotic fluid, cord blood or human serum ([Oudart et al., 2013](#)).

The physiological mechanisms leading to the release of this matrikine remained unknown for a long time. Therefore, the study of collagen XIX proteolysis appeared as a key step to understand its antitumor mechanisms. Our laboratory investigated the potential cleavage of collagen XIX by different enzymes specifically involved in tumor invasion. Initially, experiments conducted with MMP-2, MMP-9 and MMP-14 showed no release of this peptide. However, the presence of an arginine in the junction area between NC1 and Col1 domains, led us to study the proteolysis of collagen XIX by plasmin. We demonstrated that plasmin was able to cleave collagen XIX. We identified by mass spectrometry the cleavage products and demonstrated the release of a peptide comprising the first 15 amino acids of the NC1 domain, preceded by a cysteine and minus the 4 amino-acids residues (TGGN). This peptide reproduce the anti-tumor activity of the complete NC1(XIX) domain. In cell culture, we demonstrated that melanoma cells transfected with an anti-tPA siRNA (one of the plasminogen activators) became unable to activate plasmin and to cleave collagen XIX for liberating the active anti-tumor peptide ([Oudart et al., 2015](#)).

#### *5.4 NC1(XIX) domain and melanoma*

Studies in our laboratory enabled us to demonstrate that NC1(XIX) significantly inhibits tumor growth in a mouse melanoma model and in a xenograft model of human melanoma in nude mice. Histological study of the tumors showed a decrease in the size and number of vessels in the tumors treated with the NC1(XIX) peptide associated with a significant decrease in VEGF expression, suggesting an anti-angiogenic effect ([Ramont et al., 2007](#)). In vitro, NC1(XIX) inhibited cell migration in artificial wound models and video-microscopy analysis showed slow and uncontrolled movements of cells. Furthermore, NC1(XIX) inhibited Matrigel® invasion by tumor cells in modified Boyden chambers. New vessel formation by endothelial cells seeded on a Matrigel® layer was also inhibited in the presence of NC1(XIX) ([Toubal et al., 2010](#)). The NC1(XIX) domain must be considered as a new anti-tumor and anti-angiogenic matrikine.

Finally, we identified for the first time the  $\alpha v\beta 3$  integrin as a receptor of NC1(XIX). Moreover, we demonstrated that NC1(XIX) inhibits the FAK (Focal Adhesion Kinase) / PI3K (Phosphoinositide 3-Kinase) / Akt (protein kinase B) / mTOR (mammalian Target Of Rapamycin) pathway, by decreasing the phosphorylation and activity of the major proteins

involved in this pathway ([Oudart et al., in press](#)). On the other hand, NC1(XIX) induced an increase of GSK3 $\beta$  activity by decreasing its degree of phosphorylation ([Oudart et al., in press](#)). Treatments targeting this central signaling pathway in the development of melanoma are promising and new molecules should be developed. NC1(XIX) appears to have the potential for the design of new anti-cancer drugs.

## 6. Conclusion

Collagen XIX is a minor collagen associated with basement membranes zone. It was isolated for the first time in a human cDNA library from rhabdomyosarcoma cells and belongs to the FACIT (Fibril Associated Collagens with Interrupted Triple Helices) family. It is located in the basement membranes of vascular, neural and epithelial tissue. The  $\alpha$ 1(XIX) chain is composed of 5 collagenous domains interrupted by 6 non-collagenous domains with a short C-terminal 19 amino-acid residues non-collagenous domain (NC1). This collagen is involved in muscle cells differentiation, central nervous system development and esophagus formation but other biological functions cannot be excluded. We have shown that the short NC1(XIX) C-terminal domain inhibits migration and invasion capacities of melanoma cells. It also exerts a strong inhibition of *in vivo* tumor growth in murine or human melanoma models. In addition to its anti-tumor properties, NC1(XIX) domain shows potent anti-angiogenic effects through an intense inhibition of MMP-14 and VEGF expression. Collagen XIX is expressed in human cell cultures, tissue extracts and various biological fluids, such as serum, amniotic fluid, cord blood and many other body fluids ([Oudart et al., 2013](#)). Finally, we identified the  $\alpha$ v $\beta$ 3 integrin as a receptor of NC1(XIX) and demonstrated that NC1(XIX) inhibits the FAK/PI3K/Akt/mTOR pathway. The discovery of natural anti-tumor and anti-angiogenic peptides such as NC1(XIX) and their development as anti-angiogenic drugs might open new therapeutic opportunities with fewer deleterious side effects compared to synthetic drugs.

Table 1: Main locations of transcripts from the COL19A1 gene during the embryonic period and in adult mice ([Sumiyoshi et al., 1997](#)).

Organ	Embryonic period	Adult
Muscle (esophagus, stomach, tongue)	+	-
Heart	+	-
Aorta	Not determined	Traces
Skin	+	-
Intestine	+	-
Central nervous system	+	++
Eyes	Not determined	+
Tail	+	-
Lung	+	Traces
Liver	-	-
Kidney	+	Traces
Testis	Not determined	+

Table 2: Main locations of collagen XIX in human adults ([Amenta et al., 2003](#); [Myers et al., 2003](#); [Myers et al., 1997](#); [Su et al., 2010](#)).

Location	Immuno-localization
Brain	Hippocampus
Skeletal muscle	Vascular basement membrane zone Peripheral nerve fiber Endomysium
Spleen	Vascular basement membrane zone Splenic sinusoids and Billroth's cords Splenic Capsule
Prostate	Vascular basement membrane zone
Kidney	Mesangium Bowman's capsule Vascular and tubular basement membrane zone
Liver	Vascular basement membrane zone Hepatic sinusoids (low level)
Colon	Vascular and epithelial, basement membrane zone
Placenta	Vascular basement membrane zone
Umbilical cord	Vascular basement membrane zone
Skin	Vascular, smooth muscle and neural basement membrane zone Dermo-epidermal basement membrane zone

## Figure captions

**Fig. 1.** Molecular organization of the  $\alpha 1(\text{XIX})$  chain. Collagen XIX belongs to the FACIT (Fibril Associated Collagens with Interrupted Triple Helices) family. Collagen XIX is composed of three  $\alpha 1(\text{XIX})$  chains, forming a homotrimer of 400 kDa. Each  $\alpha$  chain consists of 1142 amino acids with a molecular mass of 112 kDa. Each chain contains 5 collagenous domains (Col1 to Col5) interrupted by 6 noncollagenous domains (NC1 to NC6), composed of 19, 44, 23, 31, 20, 291 amino acid residues, respectively. NC6  $\alpha 1(\text{XIX})$  domain also shares very strong homologies with the N-terminal module of thrombospondin. The NC2  $\alpha 1(\text{XIX})$  domain has a stable helix structure and is probably responsible for the trimerization of the  $\alpha 1(\text{XIX})$  chains. The  $\alpha 1(\text{XIX})$  chain also presents 14 cysteine residues. The presence of cysteine promotes the formation of inter-chain and intra-chain disulfide bonds, allowing the formation and the stabilization of the triple helix. The NC1(XIX) domain located at the C-terminus of the  $\alpha 1(\text{XIX})$  chain is very short. Plasmin was able to cleave collagen XIX and release a peptide comprising the first 15 amino acids of the anti-tumor NC1 domain.

**Fig. 2.** NC1(XIX) binding to  $\alpha v\beta 3$  integrin receptor interfere with the FAK (Focal Adhesion Kinase) / PI3K (Phospholinositide 3-Kinase) / Akt (protein Kinase B) / mTOR (Mammalian Target Of Rapamycin) pathway by decreasing the phosphorylation and activity of the major proteins involved in this pathway. On the other hand, NC1(XIX) induces an increase in GSK3 $\beta$  activity by decreasing its phosphorylation level. The inhibition of this pathway could explain anti-tumor properties of the NC1(XIX) domain.




## References

- Amenta, P.S., Hadad, S., Lee, M.T., Barnard, N., Li, D., Myers, J.C., 2003. Loss of types XV and XIX collagen precedes basement membrane invasion in ductal carcinoma of the female breast. *J Pathol* 199, 298-308.
- Arya, R., Demerath, E., Jenkinson, C.P., Goring, H.H., Puppala, S., Farook, V., Fowler, S., Schneider, J., Granato, R., Resendez, R.G., Dyer, T.D., Cole, S.A., Almasy, L., Comuzzie, A.G., Siervogel, R.M., Bradshaw, B., DeFronzo, R.A., MacCluer, J., Stern, M.P., Towne, B., Blangero, J., Duggirala, R., 2006. A quantitative trait locus (QTL) on chromosome 6q influences birth weight in two independent family studies. *Hum Mol Genet* 15, 1569-1579.
- Aumailley, M., Timpl, R., 1986. Attachment of cells to basement membrane collagen type IV. *J Cell Biol* 103, 1569-1575.
- Boudko, S.P., Engel, J., Bachinger, H.P., 2008. Trimerization and triple helix stabilization of the collagen XIX NC2 domain. *J Biol Chem* 283, 34345-34351.
- Chiquet, M., Birk, D.E., Bonnemant, C.G., Koch, M., 2014. Collagen XII: Protecting bone and muscle integrity by organizing collagen fibrils. *Int J Biochem Cell Biol* 53, 51-54.
- Davis, G.E., Bayless, K.J., Davis, M.J., Meininger, G.A., 2000. Regulation of tissue injury responses by the exposure of matricryptic sites within extracellular matrix molecules. *Am J Pathol* 156, 1489-1498.
- Floquet, N., Pasco, S., Ramont, L., Derreumaux, P., Laronze, J.Y., Nuzillard, J.M., Maquart, F.X., Alix, A.J., Monboisse, J.C., 2004. The antitumor properties of the alpha3(IV)-(185-203) peptide from the NC1 domain of type IV collagen (tumstatin) are conformation-dependent. *J Biol Chem* 279, 2091-2100.
- Gerecke, D.R., Olson, P.F., Koch, M., Knoll, J.H., Taylor, R., Hudson, D.L., Champlaud, M.F., Olsen, B.R., Burgeson, R.E., 1997. Complete primary structure of two splice variants of collagen XII, and assignment of alpha 1(XII) collagen (COL12A1), alpha 1(IX) collagen (COL9A1), and alpha 1(XIX) collagen (COL19A1) to human chromosome 6q12-q13. *Genomics* 41, 236-242.
- Hilario, J.D., Wang, C., Beattie, C.E., 2010. Collagen XIXa1 is crucial for motor axon navigation at intermediate targets. *Development* 137, 4261-4269.
- Inoguchi, K., Yoshioka, H., Khaleduzzaman, M., Ninomiya, Y., 1995. The mRNA for alpha 1(XIX) collagen chain, a new member of FACITs, contains a long unusual 3' untranslated region and displays many unique splicing variants. *J Biochem* 117, 137-146.
- Ivanova, V.P., Krivchenko, A.I., 2014. [A current viewpoint on structure and evolution of collagens. II. The fibril-associated collagens with interrupted triple helices]. *Zh Evol Biokhim Fiziol* 50, 245-254.
- Kalluri, R., 2003. Basement membranes: structure, assembly and role in tumour angiogenesis. *Nat Rev Cancer* 3, 422-433.
- Khaleduzzaman, M., Sumiyoshi, H., Ueki, Y., Inoguchi, K., Ninomiya, Y., Yoshioka, H., 1997. Structure of the human type XIX collagen (COL19A1) gene, which suggests it has arisen from an ancestor gene of the FACIT family. *Genomics* 45, 304-312.
- Kimura, T., Mattei, M.G., Stevens, J.W., Goldring, M.B., Ninomiya, Y., Olsen, B.R., 1989. Molecular cloning of rat and human type IX collagen cDNA and localization of the alpha 1(IX) gene on the human chromosome 6. *Eur J Biochem* 179, 71-78.
- Kruegel, J., Miosge, N., 2010. Basement membrane components are key players in specialized extracellular matrices. *Cell Mol Life Sci* 67, 2879-2895.
- Maquart, F.X., Simeon, A., Pasco, S., Monboisse, J.C., 1999. Regulation of cell activity by the extracellular matrix: the concept of matrikines. *J Soc Biol* 193, 423-428.
- Monboisse, J.C., Oudart, J.B., Ramont, L., Brassart-Pasco, S., Maquart, F.X., 2014. Matrikines from basement membrane collagens: a new anti-cancer strategy. *Biochim Biophys Acta* 1840, 2589-2598.
- Myers, J.C., Li, D., Amenta, P.S., Clark, C.C., Nagaswami, C., Weisel, J.W., 2003. Type XIX collagen purified from human umbilical cord is characterized by multiple sharp kinks delineating collagenous subdomains and by intermolecular aggregates via globular, disulfide-linked, and heparin-binding amino termini. *J Biol Chem* 278, 32047-32057.

- Myers, J.C., Li, D., Bageris, A., Abraham, V., Dion, A.S., Amenta, P.S., 1997. Biochemical and immunohistochemical characterization of human type XIX defines a novel class of basement membrane zone collagens. *Am J Pathol* 151, 1729-1740.
- Myers, J.C., Li, D., Rubinstein, N.A., Clark, C.C., 1999. Up-regulation of type XIX collagen in rhabdomyosarcoma cells accompanies myogenic differentiation. *Exp Cell Res* 253, 587-598.
- Myers, J.C., Sun, M.J., D'Ippolito, J.A., Jabs, E.W., Neilson, E.G., Dion, A.S., 1993. Human cDNA clones transcribed from an unusually high-molecular-weight RNA encode a new collagen chain. *Gene* 123, 211-217.
- Myers, J.C., Yang, H., D'Ippolito, J.A., Presente, A., Miller, M.K., Dion, A.S., 1994. The triple-helical region of human type XIX collagen consists of multiple collagenous subdomains and exhibits limited sequence homology to alpha 1(XVI). *J Biol Chem* 269, 18549-18557.
- Nassa, M., Anand, P., Jain, A., Chhabra, A., Jaiswal, A., Malhotra, U., Rani, V., 2012. Analysis of human collagen sequences. *Bioinformation* 8, 26-33.
- O'Reilly, M.S., Boehm, T., Shing, Y., Fukai, N., Vasios, G., Lane, W.S., Flynn, E., Birkhead, J.R., Olsen, B.R., Folkman, J., 1997. Endostatin: an endogenous inhibitor of angiogenesis and tumor growth. *Cell* 88, 277-285.
- Oh, S.P., Taylor, R.W., Gerecke, D.R., Rochelle, J.M., Seldin, M.F., Olsen, B.R., 1992. The mouse alpha 1(XII) and human alpha 1(XII)-like collagen genes are localized on mouse chromosome 9 and human chromosome 6. *Genomics* 14, 225-231.
- Oudart, J.B., Brassart-Pasco, S., Luczka, E., Dupont-Deshorgue, A., Bellon, G., Boudko, S.P., Bachinger, H.P., Monboisse, J.C., Maquart, F.X., Ramont, L., 2013. Analytical methods for measuring collagen XIX in human cell cultures, tissue extracts, and biological fluids. *Anal Biochem* 437, 111-117.
- Oudart, J.B., Brassart-Pasco, S., Vautrin, A., Sellier, C., Machado, C., Dupont-Deshorgue, A., Brassart, B., Baud, S., Dauchez, M., Monboisse, J.C., Harakat, D., Maquart, F.X., Ramont, L., 2015. Plasmin releases the anti-tumor peptide from the NC1 domain of collagen XIX. *Oncotarget* 6, 3656-3668.
- Parsons, P., Gilbert, S.J., Vaughan-Thomas, A., Sorrell, D.A., Notman, R., Bishop, M., Hayes, A.J., Mason, D.J., Duance, V.C., 2011. Type IX collagen interacts with fibronectin providing an important molecular bridge in articular cartilage. *J Biol Chem* 286, 34986-34997.
- Pasco, S., Brassart, B., Ramont, L., Maquart, F.X., Monboisse, J.C., 2005. Control of melanoma cell invasion by type IV collagen. *Cancer Detect Prev* 29, 260-266.
- Paulsson, M., 1992. Basement membrane proteins: structure, assembly, and cellular interactions. *Crit Rev Biochem Mol Biol* 27, 93-127.
- Ramchandran, R., Dhanabal, M., Volk, R., Waterman, M.J., Segal, M., Lu, H., Knebelmann, B., Sukhatme, V.P., 1999. Antiangiogenic activity of restin, NC10 domain of human collagen XV: comparison to endostatin. *Biochem Biophys Res Commun* 255, 735-739.
- Ramont, L., Brassart-Pasco, S., Thevenard, J., Deshorgue, A., Venteo, L., Laronze, J.Y., Pluot, M., Monboisse, J.C., Maquart, F.X., 2007. The NC1 domain of type XIX collagen inhibits in vivo melanoma growth. *Mol Cancer Ther* 6, 506-514.
- Ricard-Blum, S., 2011. The collagen family. *Cold Spring Harb Perspect Biol* 3, a004978.
- Ricard-Blum, S., Ruggiero, F., 2005. The collagen superfamily: from the extracellular matrix to the cell membrane. *Pathol Biol (Paris)* 53, 430-442.
- Shtilbans, A., Choi, S.G., Fowkes, M.E., Khitrov, G., Shahbazi, M., Ting, J., Zhang, W., Sun, Y., Sealfon, S.C., Lange, D.J., 2011. Differential gene expression in patients with amyotrophic lateral sclerosis. *Amyotroph Lateral Scler* 12, 250-256.
- Su, J., Gorse, K., Ramirez, F., Fox, M.A., 2010. Collagen XIX is expressed by interneurons and contributes to the formation of hippocampal synapses. *J Comp Neurol* 518, 229-253.
- Sumiyoshi, H., Inoguchi, K., Khaleduzzaman, M., Ninomiya, Y., Yoshioka, H., 1997. Ubiquitous expression of the alpha1(XIX) collagen gene (Col19a1) during mouse embryogenesis becomes restricted to a few tissues in the adult organism. *J Biol Chem* 272, 17104-17111.

- Sumiyoshi, H., Laub, F., Yoshioka, H., Ramirez, F., 2001. Embryonic expression of type XIX collagen is transient and confined to muscle cells. *Dev Dyn* 220, 155-162.
- Sumiyoshi, H., Mor, N., Lee, S.Y., Doty, S., Henderson, S., Tanaka, S., Yoshioka, H., Rattan, S., Ramirez, F., 2004. Esophageal muscle physiology and morphogenesis require assembly of a collagen XIX-rich basement membrane zone. *J Cell Biol* 166, 591-600.
- Tao, G., Levay, A.K., Peacock, J.D., Huk, D.J., Both, S.N., Purcell, N.H., Pinto, J.R., Galantowicz, M.L., Koch, M., Lucchesi, P.A., Birk, D.E., Lincoln, J., 2012. Collagen XIV is important for growth and structural integrity of the myocardium. *J Mol Cell Cardiol* 53, 626-638.
- Thevenard, J., Floquet, N., Ramont, L., Prost, E., Nuzillard, J.M., Dauchez, M., Yezid, H., Alix, A.J., Maquart, F.X., Monboisse, J.C., Brassart-Pasco, S., 2006. Structural and antitumor properties of the YSNSG cyclopeptide derived from tumstatin. *Chem Biol* 13, 1307-1315.
- Thevenard, J., Ramont, L., Devy, J., Brassart, B., Dupont-Deshorgue, A., Floquet, N., Schneider, L., Ouchani, F., Terryn, C., Maquart, F.X., Monboisse, J.C., Brassart-Pasco, S., 2010. The YSNSG cyclopeptide derived from tumstatin inhibits tumor angiogenesis by down-regulating endothelial cell migration. *Int J Cancer* 126, 1055-1066.
- Toubal, A., Ramont, L., Terryn, C., Brassart-Pasco, S., Patigny, D., Sapi, J., Monboisse, J.C., Maquart, F.X., 2010. The NC1 domain of type XIX collagen inhibits melanoma cell migration. *Eur J Dermatol* 20, 712-718.
- van der Rest, M., Garrone, R., 1991. Collagen family of proteins. *FASEB J* 5, 2814-2823.
- Yoshioka, H., Zhang, H., Ramirez, F., Mattei, M.G., Moradi-Ameli, M., van der Rest, M., Gordon, M.K., 1992. Synteny between the loci for a novel FACIT-like collagen locus (D6S228E) and alpha 1 (IX) collagen (COL9A1) on 6q12-q14 in humans. *Genomics* 13, 884-886.
- Yurchenco, P.D., 2011. Basement membranes: cell scaffoldings and signaling platforms. *Cold Spring Harb Perspect Biol* 3.
- Yurchenco, P.D., O'Rear, J.J., 1994. Basement membrane assembly. *Methods Enzymol* 245, 489-518.



**Figure 1**




Figure 2