

**Glucosinolates of *Lepidium graminifolium* L.
(Brassicaceae) from Croatia**

Sabine Montaut, Sharayah Read, Ivica Blažević, Jean-Marc Nuzillard,
Dominique Harakat, Patrick Rollin

► **To cite this version:**

Sabine Montaut, Sharayah Read, Ivica Blažević, Jean-Marc Nuzillard, Dominique Harakat, et al..
Glucosinolates of *Lepidium graminifolium* L. (Brassicaceae) from Croatia. *Natural Product Research*,
Taylor & Francis, In press, 10.1080/14786419.2019.1633643 . hal-02307209

HAL Id: hal-02307209

<https://hal.univ-reims.fr/hal-02307209>

Submitted on 5 Nov 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Glucosinolates of *Lepidium graminifolium* L. (Brassicaceae) from Croatia

Sabine Montaut^{a,*}, Sharayah Read^a, Ivica Blažević^b, Jean-Marc Nuzillard^c,
Dominique Harakat^c, Patrick Rollin^d

^a *Department of Chemistry and Biochemistry, Biomolecular Sciences Programme,
Laurentian University, 935 Ramsey Lake Road, Sudbury, ON P3E 2C6 Canada*

^b *Department of Organic Chemistry, Faculty of Chemistry and Technology, University
of Split, Ruđera Boškovića 35, 21000 Split, Croatia*

^c *Institut de Chimie Moléculaire de Reims, CNRS UMR 7312, Moulin de la Housse,
CPCBAI Bâtiment 18, BP 1039, 51687 Reims Cedex 2, France*

^d *Université d'Orléans et CNRS, ICOA, UMR 7311, BP 6759, F-45067 Orléans, France*

* Corresponding author: Email: smontaut@laurentian.ca

ORCID ID for Sabine Montaut: <https://orcid.org/0000-0002-8223-4809>

Glucosinolates of *Lepidium graminifolium* L. (Brassicaceae) from Croatia

The glucosinolate (GSL) profiles (inflorescence, stem, root, and fruit) of the wild-growing plant *Lepidium graminifolium* L. (Brassicaceae) from Croatia was established by LC-MS analysis. During this investigation, we confirmed the presence of benzyl- (**1**), 3-methoxybenzyl- (**2**), 4-hydroxybenzyl- (**4**), 4-methoxyindol-3-ylmethyl- (**7**) GSLs and reported for the first time in the plant the presence of (2*R*)-hydroxybut-3-enyl- (**11**), (2*S*)-hydroxybut-3-enyl- (**12**), but-3-enyl- (**13**), and 2-phenylethyl- (**14**) GSLs. Finally, 3-hydroxybenzyl GSL (**3**) was isolated for the first time from *L. graminifolium* inflorescence and characterized by spectroscopic data interpretation.

Keywords: *Lepidium graminifolium*; Brassicaceae; glucosinolate; glucolepigramin; LC-MS; NMR.

1. Introduction

Lepidium graminifolium L. (grassleaf pepperweed) is a wild and annual plant (Brassicaceae) of the Mediterranean region and Eastern Europe, used as a medicinal, food, or ornamental plant (Agelet et al. 2000; Savo et al. 2011; Guarrera & Savo 2016).

The fatty acid content of the seed oil of *L. graminifolium* and flavonoids in the aerial parts of the plant have been reported previously (Fursa et al. 1981; Miller et al. 1965; Nilsson et al. 1998). Using paper chromatography (PC), benzyl isothiocyanate (ITC) was detected in root and aerial part extracts of a wild *L. graminifolium* (origin France), suggesting that benzyl glucosinolate (GSL) (**1**) (Figure 1) was present in those plant organs (Delaveau, 1958). 3-Methoxybenzyl- (**2**) and 3-hydroxybenzyl- (**3**) GSLs were identified by PC, IR spectroscopy, and mass spectrometry of the hydrolysis products from seeds of plants cultivated in Denmark (Friis & Kjær 1963). PC of a seed extract detected two more GSLs – one of which was supposed to be 4-hydroxybenzyl GSL (**4**) (Danielak & Borkowski 1969). 3-Methylsulfanylpropyl ITC, benzyl ITC, and phenylacetone nitrile were identified by GC-MS in 8-week-old seedlings grown in a glasshouse, indicating the presence of 3-methylsulfanylpropyl GSL (**5**) and **1** (Cole 1976). Indol-3-ylmethyl- (**6**), 4-methoxyindol-3-ylmethyl- (**7**), and 1-methoxyindol-3-

ylmethyl- (**8**) GSLs were detected by HPLC analysis of 10-15-day-old seedlings (Bäuerle et al. 1986). Finally, a GC analysis of the hydrolysis products of the compounds extracted from the seeds (origin Spain) showed the presence of 3-hydroxybenzyl-, methoxybenzyl-, 3,4-dimethoxybenzyl-, 3,4,5-trimethoxybenzyl ITCs and thiocyanate ion, suggesting that the seeds contained **2-4**, 3,4-dimethoxybenzyl- (**9**), and 3,4,5-trimethoxybenzyl- (**10**) GSLs (Figure 1) (Daxenbichler et al. 1991).

As part of our continued interest in the chemistry of the *Lepidium* sp. (Montaut et al. 2017), the aim of this study was to identify the GSLs present in inflorescence, stem, root, and fruit of *L. graminifolium* wild-growing in Croatia by HPLC-ESI-MS coupled with a photodiode-array detector. Finally, the major GSL in the fruit, 3-hydroxybenzyl GSL (**3**), was isolated and characterized using spectroscopic techniques.

2. Results and discussion

The inflorescence, stem, root, and fruit of *L. graminifolium* were harvested on Rab Island (Croatia). The plant parts were extracted and analyzed by HPLC-ESI-MS for intact GSLs (Tutin et al. 1968-1980; Zrybko et al. 1997). The t_R , mass, and UV data of the products were compared with those of standards from our HPLC-ESI-MS library including the authenticated isolated GSLs **2** and **7** (Baird et al. 1988; Montaut et al. 2010) and the commercial standards **1**, **4**, (2*R*)-hydroxybut-3-enyl- (**11**), (2*S*)-hydroxybut-3-enyl- (**12**), but-3-enyl- (**13**), and 2-phenylethyl- (**14**) GSLs (see supplementary material section). In the inflorescence (Figure S1a), the minor compounds at t_R 7.4 min, t_R 9.1 min, and t_R 23.4 min were determined to be **11** (Figures 1 and S1, Tables S1 and S2), **12**, and **1**, respectively. The other minor compound at t_R 25.6 min was found to be **2**. GSLs **2**, **11**, and **12** were also present in the stem (Figure S1b), root (Figure S2a), and fruit (Figure S2b) while **1** was also identified in the root and fruit. In addition, **7** (t_R 29.1 min) was identified in the stem. GSLs **13** (t_R 18.2 min) and **14** (t_R 26.9 min) were found in the root. Finally, a minor compound at t_R 17.9 min observed in the fruit and stem was found to be **4**. The compound at t_R 20.3 min found in the inflorescence, stem, root, and fruit extracts (Figures S1 and S2) did not match any GSLs in our library. Therefore, it was isolated (see supplementary material section) and its structure was elucidated using UV, NMR (^1H , ^{13}C , HMBC, COSY, and HSQC), and HRMS data. The data reported in the supplementary material section suggest that the peak at t_R 20.3 min corresponds to 3-hydroxybenzyl GSL (**3**) (Figure 1).

The presence of **3** in *L. graminifolium* seeds was previously proposed thanks to the structural elucidation of its hydrolysis product (Friis & Kjær 1963). It was shown in our study that **3** is not only produced in the seed but in all parts of the plant. Like other *Lepidium* species, *L. graminifolium* appears to be a source of arylaliphatic GSLs (Montaut et al. 2017); in fact, **1** and **2** are the major GSLs in the root while **2** and **3** are the major GSLs in the fruit. In previous investigations, **2** was only reported in the seed of *L. graminifolium* (Friis & Kjær 1963; Daxenbichler et al. 1991); our study additionally showed that **2** is found in all plant parts. Furthermore, **4** was only detected in the seeds in a previous study (Danielak & Borkowski, 1969), whereas it was also found by us to be a minor GSL in both stem and fruit. While **7** was previously detected in 10-15-day-old seedlings of *L. graminifolium* (Bäuerle et al. 1986), we could only detect it in the stem. Contrary to previous reports, we neither detected any **5**, **6**, **9** nor **10** in *L. graminifolium*. The differences in GSL profiles may be due to genetic, geographical and ecological impacts, analytical methods, or the fact that we have analyzed a wild *L. graminifolium*, since some previous investigations were carried out on cultivated plants or samples.

3. Conclusion

We report for the first time in *L. graminifolium* harvested in Croatia the presence of four known GSLs (**11-14**). Furthermore, **3** was isolated and characterized by spectroscopic data interpretation for the first time from the inflorescence. Finally, we have confirmed by LC-MS analyses the presence of four previously reported GSLs (**1**, **2**, **4**, and **7**).

Supplementary material

Experimental details related to this paper are available online, alongside Tables S1-S2 and Figures S1-S2.

Disclosure statement

No potential conflict of interest was reported by the authors.

Funding

Funding from 1) Natural Sciences and Engineering Research Council of Canada (RTI grant 315095-05), Canadian Foundation for Innovation (Leaders Opportunity Fund) – Ontario Research Fund (grant 11666), and Laurentian University (Career and Employment Centre) is acknowledged by SM; 2) CNRS, Conseil Régional Champagne Ardenne, Conseil Général de la Marne, Ministry of Higher Education and Research (MESR), and EU-programme FEDER to the P1AneT CPER project is acknowledged by JMN, and 3) the Croatian Science Foundation (grant IP-2016-06-1316) is acknowledged by IB.

References

- Agelet A, Bonet MÀ, Vallès J. 2000. Homegardens and their role as a main source of medicinal plants in mountain regions of Catalonia (Iberian peninsula). *Econ Bot.* 54: 295-309.
- Baird WM, Zennie TM, Ferin M, Chae Y-H, Hatchell J, Cassady JM. 1988. Glucolimnanthin, a plant glucosinolate, increases the metabolism and DNA binding of benzo[a]pyrene in hamster embryo cell cultures. *Carcinogenesis* 9: 657-660.
- Bäuerle R, Wagner H, Schraudolf H. 1986. Distribution of 4-methoxy-3-indolylmethyl-glucosinolate (4-methoxy-glucobrassicin) in Brassicaceae. *Experientia* 42: 86.
- Cole RA., 1976. Isothiocyanates, nitriles and thiocyanates as products of autolysis of glucosinolates in Cruciferae. *Phytochemistry* 15: 759-762.
- Danielak R, Borkowski B. 1969. Biologically active compounds in seeds of crucifers. Part III. Chromatographical search for glucosynolates. *Dissert Pharm Pharmacol.* 21: 563-575.
- Daxenbichler ME, Spencer GF, Carlson DG, Rose GB, Brinker AM, Powell RG. 1991. Glucosinolate composition of seeds from 297 species of wild plants. *Phytochemistry.* 30(8): 2623-2638.
- Delaveau P. 1958. Multiplicité des hétérosides à sévenol chez les Crucifères; leur relation avec la physiologie et la taxinomie (2e note). *Bull Soc Bot France* 105: 224-226.

- Friis P, Kjær A. 1963. Glucolepigramin, a new thioglucoside, present in *Lepidium graminifolium* L. Acta Chem Scand. 17: 1515-1520.
- Fursa NS, Chaika EA, Belyaeva LE. 1981. Flavonoids of *Lepidium* L. and *Cardaria* Desv. species in the southern Ukraine. Rastitel'nye Resursy 17: 91-94.
- Guarrera PM, Savo V. 2016. Wild food plants used in traditional vegetable mixtures in Italy. J Ethnopharmacol. 185: 202-234.
- Miller RW, Earle FR, Wolff IA. 1965. Search for new industrial oils. XIII. Oils from 102 species of Cruciferae. J Am Oil Chem Soc. 42: 817-821.
- Montaut S, Bleeker RS, Jacques C. 2010. Phytochemical constituents of *Cardamine diphylla*. Can J Chem. 88: 50-55.
- Montaut S, Benson HJ, Kay M, Guido BS, Mahboob SS, Chénier J, Gasparetto J-L, Joly HA. 2017. Probing the free-radical scavenging activity of the extract, the major glucosinolate and isothiocyanate of *Eruca sativa* Mill. and *Lepidium densiflorum* Schrad. seeds. J Food Compos Anal. 61: 52-58.
- Nilsson P, Johansson S-Å, Merker A. 1998. Variation in seed oil composition of species from the genera *Barbarea* and *Lepidium*. Acta Agric Scand Sect B, Soil and Plant Sci. 48: 159-164.
- Savo V, Caneva G, Guarrera PM, Reedy D. 2011. Folk phytotherapy of the Amalfi Coast (Campania, Southern Italy). J Ethnopharmacol. 135: 376-392.
- Tutin TG, Heywood VH, Burges NA, Moore DM, Valentine DH, Walter SM, Webb DA. (1968-1980). *Flora Europaea*. Cambridge, U.K.: Cambridge University Press.
- Zrybko CL, Fukuda EK, Rosen RT. 1997. Determination of glucosinolates in domestic and wild mustard by high-performance liquid chromatography with confirmation by electrospray mass spectrometry and photodiode-array detection. J Chromatogr A. 767: 43-52.

Figure caption

Figure 1. Structures of glucosinolates **1-14** from *Lepidium graminifolium*