



HAL
open science

Relationship between gender and survival in a real-life cohort of patients with COPD

Maéva Zysman, Pierre-Régis Burgel, Isabelle Court-Fortune, Graziella Brinchault-Rabin, Pascale Nesme-Meyer, Pascale Surpas, Gaetan Deslée, Thierry Perez, Olivier Le Rouzic, Gilles Jebrak, et al.

► **To cite this version:**

Maéva Zysman, Pierre-Régis Burgel, Isabelle Court-Fortune, Graziella Brinchault-Rabin, Pascale Nesme-Meyer, et al.. Relationship between gender and survival in a real-life cohort of patients with COPD. *Respiratory Research*, 2019, 20 (1), pp.191. 10.1186/s12931-019-1154-3 . hal-02448813

HAL Id: hal-02448813

<https://hal.univ-reims.fr/hal-02448813>

Submitted on 3 Jun 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

LETTER TO THE EDITOR

Open Access

Relationship between gender and survival in a real-life cohort of patients with COPD



Maeva Zysman^{1,2,14*} , Pierre-Régis Burgel³, Isabelle Court-Fortune⁴, Graziella Brinchault-Rabin⁵, Pascale Nesme-Meyer⁶, Pascale Surpas⁷, Gaetan Deslée⁸, Thierry Perez⁹, Olivier Le Rouzic⁹, Gilles Jebrak¹⁰, Pascal Chanez¹¹, Jean-Louis Paillasseur¹², Denis Caillaud¹³, Nicolas Roche³ and on behalf of the Initiatives BPCO scientific committee and investigators

Abstract

Background: Although COPD affects both men and women, its prevalence is increasing more rapidly in women. Disease outcomes appear different among women with more frequent dyspnea and anxiety or depression but whether this translates into a different prognosis remains to be determined. Our aim was to assess whether the greater clinical impact of COPD in women was associated with differences in 3-year mortality rates.

Methods: In the French Initiatives BPCO real-world cohort, 177 women were matched up to 458 men on age (within 5-year intervals) and FEV₁ (within 5% predicted intervals). 3-year mortality rate and survival were analyzed. Univariate and multivariate logistic regression analyses were performed.

Results: For a given age and level of airflow obstruction, women with COPD had more severe dyspnea, lower BMI, and were more likely to exhibit anxiety. Nevertheless, three-year mortality rate was comparable among men and women, respectively 11.2 and 10.8%. In a multivariate model, the only factors significantly associated with mortality were dyspnea and malnutrition but not gender.

Conclusion: Although women with COPD experience higher levels of dyspnea and anxiety than men at comparable levels of age and FEV₁, these differences do not translate into variations in 3-year mortality rates.

Trial registration: 04–479.

Keywords: Chronic obstructive pulmonary disease, Survival, Gender differences

Background

Influence of gender on COPD expression and outcomes is an area of sustained interest [1, 2]. Although COPD affects both men and women, its prevalence is increasing more rapidly in women, particularly in younger women [1]. Women are more likely to be misdiagnosed [3], whereas there is increasing evidence suggesting gender-related differences in COPD risk. For example, female smokers are at greater risk of airflow obstruction than male smokers [4]. Disease progression and outcomes appear different among women and men with COPD [5, 6]. Younger women with COPD have a greater likelihood of more severe dyspnea and airflow limitation, and

exhibit a higher risk of exacerbations [7, 8]. In COPD populations, several longitudinal studies showed an association between higher levels of symptoms and poorer prognosis [9]. Of interest, studies have found discrepant results regarding the relationship between gender and survival [10].

As shown in other studies [10, 11], previous analysis of the Initiatives BPCO cohort found that women suffer from higher levels of dyspnea and anxiety even after matching on age and FEV₁ [12]. Whether these gender-related differences in symptoms translate into differences in survival remains unknown. Our aim was to assess whether the greater clinical impact of COPD in women was associated with differences in 3-year mortality rates.

* Correspondence: maeva55@club-internet.fr

¹Pulmonary Department, Nancy, France

²Inserm U955, team 04, IMRB, Créteil, France

Full list of author information is available at the end of the article



Methods

As previously described, Initiatives BPCO is a rolling cohort of patients with COPD followed at French University Hospitals [13]. The primary aim of the cohort was to study COPD phenotypes, as previously reported [13]. The following data are collected as part of routine practice at inclusion: demographic and anthropometric characteristics, occupational exposures, smoking history, chronic bronchitis, exacerbation frequency, dyspnea assessed by mMRC dyspnea scale, health status, physician diagnosed comorbidities (asthma, rhinitis, cardiovascular diseases, obesity, diabetes, mechanical limitation, psychological status), medications and post-bronchodilator spirometry (FEV₁, FVC).

Men and women were matched up to 3:1 on age (within 5-year intervals) and FEV₁ (within 5% predicted intervals) leading to a small loss of sample size. Three-year mortality

rate and survival were analyzed using logistic regression and Kaplan-Meier analysis with log-rank test, respectively. Univariate comparisons between matched men and women were performed by chi2 and t-test. To identify which risk factors play a critical role as determinants of mortality in the studied population, we performed a multivariate stepwise logistic regression analysis with the following tested covariates: cumulative smoking, chronic bronchitis, mMRC grade, FEV₁% predicted, exacerbation history during the year prior to inclusion, allergic rhinitis, associated asthma, nutritional status, hypertension, ischemic heart disease, left heart failure, diabetes, sleep apnea syndrome and age. Data are provided as median [Q1 – Q3] or n (%), as appropriate.

The study was approved by the Ethics Committee of Versailles (France), trial registration #04–479, and all subjects provided informed written consent.

Table 1 Characteristics of the studied population and univariate comparisons between age- and FEV₁-matched (3:1 ratio) men (*n* = 458) and women (*n* = 177)

Variables	Women	Men	p
N	177	458	
Age (years)	62 [56–70]	63 [57–71]	0.4171
BMI	23.2 [20.2–27.1]	25.5 [22.1–29.0]	< 0.0001
Malnutrition (BMI < 18 kg/m ²)	25 (14.1%)	31 (6.8%)	0.003
Smoking (Pack-years)	38.0 [24.8–57.0]	41 [27–56]	0.4080
FEV ₁ (%)	54 [37–69]	53 [36–67]	0.4257
Number of moderate to severe exacerbations in the previous year	1 [0–3]	1 [0–2]	0.1209
mMRC	2 [1–3]	1 [1–2]	0.0014
mMRC ≥ 2	100 (56.5%)	212 (46.3%)	0.021
BOD index	3 [1–4]	2 [1–4]	0.0831
SGRQ	46 [32–57] ^a	43 [28–59] ^a	0.5776
Asthma history	28 (15.8%)	52 (11.4%)	0.128
Rhinitis	29 (16.4%)	42 (9.2%)	0.010
Chronic bronchitis	118 (66.7%)	302 (65.9%)	0.862
Hypertension	59 (33.3%)	159 (34.7%)	0.742
Left heart failure	16 (9%)	54 (11.8%)	0.321
Ischemic heart disease	11 (6.2%)	79 (17.2%)	0.0001
Diabetes mellitus	16 (9%)	58 (12.7%)	0.202
Obstructive sleep apnea	4 (2.3%)	40 (8.7%)	0.004
HAD total score	15 [11–21] ^b	12 [8–17] ^b	< 0.0001
Anxiety HAD A ≥ 10	65 (44.5%) ^b	92 (27.6%) ^b	0.0001
Depression HAD D ≥ 10	34 (23.6%) ^b	59 (17.7%) ^b	0.136
3-year mortality	19 (10.8%)	51 (11.2%)	0.896
Age at death	72 [66–79]	68 [63–76]	0.1691

BMI body mass index, HAD hospital anxiety and depression scale, mMRC modified medical respiratory council, SGRQ Saint George's Respiratory Questionnaire.

Data are provided as median [Q1 – Q3] or n (%), as appropriate

^aMissing data for SGRQ, *n* = 28 in women, *n* = 116 in men

^bMissing data for HAD scores: *n* = 33 in women, *n* = 131 in men

Results

Among 954 patients (226 women) with COPD included at the time of the analyses, 177 women were matched to 458 men. Unmatched (non-included) women did not differ from matched (included) ones except for age and FEV₁, which were the matching criteria (*data not shown*). Median values of age and percent predicted (pp) FEV₁ were 63 years and 53%, respectively. Women had lower body mass index (BMI), higher mMRC dyspnea grade, resulting in a higher BOD (BMI, airflow obstruction, dyspnea) index, and a greater proportion of anxiety (defined by a hospital anxiety-depression-A subscore ≥ 10). Rhinitis was more frequent in women, while coronary heart disease and obstructive sleep apnea syndrome were less frequent in women (Table 1). Three-year mortality rates were 11.2% in men and 10.8% in women with no significant difference (OR for men vs. women 0.9; 95% confidence interval [0.5–1.7]). Age at death was 68 years in men and 72 years in women with no significant difference. Survival was also comparable (Log-rank $p = 0.9724$, Fig. 1). In multivariate analysis, mortality was independently associated with only malnutrition ($p = 0.02$) and mMRC ($p = 0.03$), with cumulative smoking being retained in the model although of borderline significance ($p = 0.06$). Conversely, gender was not retained ($p = 0.68$).

Discussion

Several studies have been performed to assess gender-related differences in COPD expression and many found more severe manifestations of the disease in women [6, 10]. Some studies suggest that women with chronic bronchitis

have significantly worse survival [14] whereas others have demonstrated that survival does not vary among men and women in smaller cohorts [15]. In a previous analysis of the Initiatives BPCO cohort, for a given age and level of airflow obstruction, women with COPD had higher BOD (BMI, airflow obstruction, dyspnea) scores due to greater dyspnea and lower BMI, suggesting the possibility of worse prognosis in women. However, the present data showed no difference in survival between men and women matched for age and ppFEV₁, both in univariate analyses. Furthermore, even after multivariate analyses confirming a link between worse prognosis, malnutrition and breathlessness, in accordance to BODE, gender was neither validated. Similar findings were reported in the TORCH study, in which the risk of death was similar among men and women once analyses were adjusted for differences in baseline confounders [10]. Previous data comparing 265 women and 272 men with COPD matched using BODE score have shown that all-cause mortality was higher in males than females. However, women and men were not matched on age and women were significantly younger (63 vs 67 years, $p < 0.001$) [16]. Even if this age difference is small, it can have an influence on mortality. To exclude this bias, men and women were matched on age in our study.

One limitation of this study is the relatively short-term survival analysis and the absence of available data regarding specific causes of mortality, which prevents from analyzing whether some specific mortality rates differ between men and women. Our results suggest that differences between men and women in prognostic scores (here, the BOD score) and burden of symptoms (exacerbation number and




Fig. 1 Survival according to gender, Kaplan Meier analysis, women in blue, men in red

dyspnea) do not translate into higher mortality in women. Finally, as in other studies [2, 6, 12], male gender was associated with a more frequent history of cardiovascular disease (here, ischemic heart disease), maybe explaining why the risk of mortality is similar among men and women despite women exhibiting more symptoms.

Conclusion

In the present study, COPD expression differed between men and women, with women experiencing more dyspnea and anxiety, and less diagnosed coronary heart disease and sleep apnea. These differences did not translate into significant differences in 3-year mortality rates and survival.

Abbreviations

BMI: Body mass index; COPD: Chronic obstructive pulmonary disease; FEV1: Forced Expiratory volume in one second; GOLD: Global initiative for obstructive lung disease; HAD: Hospital anxiety and depression scale; mMRC: Modified Medical Research Council; OSAS: Obstructive sleep apnea syndrome

Acknowledgements

We thank the initiatives BPCO study group P.R.Burgel (Paris), G. Deslee (Reims), P. Surpas (Charnay), O. Le Rouzic (Lille), T.Perez (Lille), N. Roche (Paris), G. Brinchault-Rabin (Rennes), D. Caillaud (Clermont-Ferrand), P. Chanez (Marseille), I. Court-Fortune (Saint-Etienne), R. Escamilla (Toulouse), G. Jebrak (Paris), P.Nesme-Meyer (Lyon), M. Zysman (Nancy), C. Pinet (Toulon) and Brigitte Risse (ARAIRLOR, Association régionale d'aide aux insuffisants respiratoires de Lorraine, Nancy).

Authors' contributions

Every author made substantial contributions to conception and design, acquisition of data, and analysis and interpretation of data; has been involved in drafting the manuscript or revising it critically for important intellectual content; has given final approval of the version to be published.

Funding

The Initiatives BPCO cohort is supported by Boehringer Ingelheim France since its creation and has been funded by Pfizer until 2015, in the form of unrestricted grant. The sponsors fund the database, statistical analyses, meetings and submission fees when required. They do not participate in decisions regarding collected data, analyses, article writing and submission. For all these aspects, the Initiatives BPCO group works in total independence.

An abstract describing the results presented here was presented at ERS International Congress 2018 Paris/France, and published in the European Respiratory Journal.

Availability of data and materials

The datasets used and analyzed during the current study are available from the corresponding author on reasonable request.

Ethics approval and consent to participate

Study protocol has been approved by the Ethics Committee of Versailles Saint Quentin University, authorization number 04–479, for protection of human beings involved in biomedical research. The study has also been approved by CCTIRS (Comité Consultatif sur le Traitement de l'Information en matière de Recherche dans le domaine de la Santé), on the 6th January, 2005 (04–479). All patients provided written consent.

Consent for publication

All patients provided written consent. All authors provided consent to publication.

Competing interests

MZ reports grants and personal fees from Boehringer Ingelheim, personal fees from Novartis, personal fees from Chiesi, and personal fees from GSK outside the submitted work.

PRB reports personal fees from Aptalis, personal fees from Astra-Zeneca, grants and personal fees from Boehringer Ingelheim, personal fees from Chiesi, personal fees from GSK, personal fees from Novartis, personal fees from Pfizer, personal fees from Vertex, personal fees from Zambon, outside the submitted work.

ICF reports grants and personal fees from Boehringer Ingelheim, grants and personal fees from Novartis, outside the submitted work.

GBR reports grants and personal fees from Boehringer Ingelheim, grants and personal fees from Novartis, personal fees and non-financial support from Chiesi, outside the submitted work.

PNM reports grants and personal fees from Boehringer Ingelheim, grants and personal fees from NOVARTIS, personal fees and non-financial support from Chiesi, outside the submitted work.

PS reports grants and personal fees from Boehringer Ingelheim outside the submitted work.

GD reports personal fees from Novartis, personal fees from Astra Zeneca, personal fees from BTG/PneumRx, personal fees from Chiesi, personal fees from Boehringer Ingelheim, outside the submitted work.

TP reports personal fees from Boehringer Ingelheim, personal fees from Novartis, personal fees from GSK, personal fees from Chiesi, personal fees from Pierre Fabre, outside the submitted work.

OLR reports grants and personal fees personal fees and non-financial support from Astra Zeneca, Boehringer Ingelheim, Chiesi, Lilly and Novartis; non-financial support from Glaxo Smith Kline, MundiPharma, Pfizer, Teva, Santelys Association, Vertex and Vitalaire, all outside the submitted work.

GJ reports personal fees from Boehringer Ingelheim, personal fees from GSK, personal fees from Novartis, personal fees from Menarini, personal fees from Astra Zeneca, personal fees from Chiesi, outside the submitted work.

PC reports personal fees from Boehringer Ingelheim, personal fees from GSK, personal fees from ALK, personal fees from Novartis, personal fees from Teva, personal fees from Astra Zeneca, personal fees from Chiesi, personal fees from Sanofi, personal fees and non-financial support from SNCF, outside the submitted work.

JLP reports grants from iBPCO association, during the conduct of the study. DC has nothing to disclose.

NR reports grants and personal fees from Boehringer Ingelheim, grants and personal fees from Novartis, personal fees from Teva, personal fees from GSK, personal fees from AstraZeneca, personal fees from Chiesi, personal fees from Mundipharma, personal fees from Cipla, grants and personal fees from Pfizer, personal fees from Sanofi, personal fees from Sandoz, personal fees from 3 M, personal fees from Zambon, outside the submitted work.

Author details

¹Pulmonary Department, Nancy, France. ²Inserm U955, team 04, IMRB, Créteil, France. ³Respiratory and Intensive Care Medicine Department, Cochin Hospital, AP-HP and Paris Descartes University (EA 2511), Sorbonne Paris Cité, Paris, France. ⁴Service de Pneumologie, CHU Saint Etienne, Saint Etienne, France. ⁵Service de Pneumologie, Hôpital Pontchaillou, Rennes, France. ⁶Service de Pneumologie, Hôpital de la Croix-Rousse, Lyon, France. ⁷Centre médical de Bayère, 30, route du Vieux-Château, 69380 Charnay, France. ⁸Pulmonary Department, Maison Blanche University Hospital, INSERM U01250, Reims, France. ⁹Univ.Lille, CNRS, Inserm, CHU Lille, Institut Pasteur de Lille, U1019 - UMR 8204 - CIIL - Center for Infection and Immunity of Lille, F-59000 Lille, France. ¹⁰Service de Pneumologie, Hôpital Bichat, AP-HP, Paris, France. ¹¹Département des Maladies Respiratoires, AP-HM, Université de la Méditerranée, Marseille, France. ¹²EFFI-STAT, Paris, France. ¹³Service de Pneumologie, Hôpital Gabriel Montpied, CHU Clermont-Ferrand, Clermont-Ferrand, France. ¹⁴Maéva Zysman 8 rue du général sarraill, 94000 Creteil, France.

Received: 16 February 2019 Accepted: 5 August 2019

Published online: 22 August 2019

References

- Jenkins CR, Chapman KR, Donohue JF, Roche N, Tsiligianni I, Han MK. Improving the management of COPD in women. *Chest*. 2017;151:686–96.

2. Han MK, Arteaga-Solis E, Blenis J, Bourjeily G, Clegg DJ, DeMeo D, et al. Female sex and gender in lung/sleep health and disease. Increased understanding of basic biological, pathophysiological, and behavioral mechanisms leading to better health for female patients with lung disease. *Am J Respir Crit Care Med*. 2018;198:850–8.
3. Chapman KR, Tashkin DP, Pye DJ. Gender bias in the diagnosis of COPD. *Chest*. 2001;119:1691–5.
4. Amaral AFS, Strachan DP, Burney PGJ, Jarvis DL. Female smokers are at greater risk of airflow obstruction than male smokers. UK biobank. *Am J Respir Crit Care Med*. 2017;195:1226–35.
5. Naberan K, Azpeitia A, Cantoni J, Miravittles M. Impairment of quality of life in women with chronic obstructive pulmonary disease. *Respir Med*. 2012;106:367–73.
6. Raheison C, Tillie-Leblond I, Prudhomme A, Taillé C, Biron E, Nocent-Ejnaini C, et al. Clinical characteristics and quality of life in women with COPD: an observational study. *BMC Womens Health*. 2014;14:31.
7. DeMeo DL, Ramagopalan S, Kavati A, Vegesna A, Han MK, Yadao A, et al. COPD Gene investigators. Women manifest more severe COPD symptoms across the life course. *Int J Chron Obstruct Pulmon Dis*. 2018;13:3021–9.
8. Aryal S, Diaz-Guzman E, Mannino DM. Influence of sex on chronic obstructive pulmonary disease risk and treatment outcomes. *Int J Chron Obstruct Pulmon Dis*. 2014;9:1145–1154.12.
9. Nishimura K, Izumi T, Tsukino M, Oga T, on Behalf of the Kansai COPD Registry and Research Group in Japan. Dyspnea is a better predictor of 5-year survival than airway obstruction in patients with COPD. *Chest*. 2002;121:1434–40.
10. Celli B, Vestbo J, Jenkins CR, Jones PW, Ferguson GT, Calverley PM, et al. Investigators of the TORCH Study. Sex differences in mortality and clinical expressions of patients with chronic obstructive pulmonary disease. The TORCH experience. *Am J Respir Crit Care Med*. 2011;183:317–22.
11. Di Marco F, Verga M, Reggente M, Maria Casanova F, Santus P, Blasi F, et al. Anxiety and depression in COPD patients: the roles of gender and disease severity. *Respir Med*. 2006;100:1767–74.
12. Roche N, Deslée G, Caillaud D, Brinchault G, Court-Fortune I, Nesme-Meyer P, Surpas P, Escamilla R, Perez T, Chanez P, Pinet C, Jebrak G, Paillasseur JL, Burgel PR, INITIATIVES BPCO Scientific Committee. Impact of gender on COPD expression in a real-life cohort. *Respir Res*. 2014;15:20.
13. Burgel PR, Paillasseur JL, Caillaud D, Tillie-Leblond I, Chanez P, Escamilla R, Court-Fortune I, Perez T, Carré P, Roche N, Initiatives BPCO Scientific Committee. Clinical COPD phenotypes: a novel approach using principal component and cluster analyses. *Eur Respir J*. 2010;36:531–9.
14. Lahousse L, Seys LJM, Joos GF, Franco OH, Stricker BH, Brusselle GG. Epidemiology and impact of chronic bronchitis in chronic obstructive pulmonary disease. *Eur Respir J*. 2017;50:1602470.
15. Prudente R, Franco EAT, Mesquita CB, Ferrari R, de Godoy I, Tanni SE. Predictors of mortality in patients with COPD after 9 years. *Int J Chron Obstruct Pulmon Dis*. 2018;17(13):3389–98.
16. de Torres JP, Cote CG, López MV, Casanova C, Díaz O, Marin JM, et al. Sex differences in mortality in patients with COPD. *Eur Respir J*. 2009;33:528–35.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Ready to submit your research? Choose BMC and benefit from:

- fast, convenient online submission
- thorough peer review by experienced researchers in your field
- rapid publication on acceptance
- support for research data, including large and complex data types
- gold Open Access which fosters wider collaboration and increased citations
- maximum visibility for your research: over 100M website views per year

At BMC, research is always in progress.

Learn more biomedcentral.com/submissions

