



RHAMNOLIPID SOLUTIONS WITCH ANTIFUNGAL AND ELICITING PROPERTIES TO PROTECT RAPESEED AGAINST LEPTOSPHERA MACULANS

Monnier M, Marion Cordier, Dahi A, Sarrazin Catherine, Penaud A,
Christophe Clement, Sylvain Cordelier, Stephan Dorey, Rippa S

► To cite this version:

Monnier M, Marion Cordier, Dahi A, Sarrazin Catherine, Penaud A, et al.. RHAMNOLIPID SOLUTIONS WITCH ANTIFUNGAL AND ELICITING PROPERTIES TO PROTECT RAPESEED AGAINST LEPTOSPHERA MACULANS. PHOMA PlantBioProtech, 2019, AGADIR, Morocco. hal-03123879

HAL Id: hal-03123879

<https://hal.univ-reims.fr/hal-03123879>

Submitted on 28 Jan 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Rhamnolipid solutions with antifungal and eliciting properties to protect rapeseed against *Leptosphaeria maculans*.

N. MONNIER², M. CORDIER¹, A. DAHI²,, C. SARAZIN³, A. PENAUD⁴, C. CLEMENT¹, S. DOREY¹, S. CORDELIER¹, S. RIPPA²


¹Université de Technologie de Compiègne, Laboratoire de Génie Enzymatique et Cellulaire, UMR CNRS 7025, Compiègne, SFR Condorcet – FR CNRS 3417, France

²Université de Reims Champagne-Ardenne, Laboratoire Résistance Induite et Bio Protection des plantes - EA 4707, SFR Condorcet – FR CNRS 3417, Reims, France

³Université de Picardie Jules Vernes, Laboratoire de Génie Enzymatique et Cellulaire, UMR CNRS 7025, Amiens, France


⁴Terres Inovia, Département de Génétique et Protection des Cultures, Thiverval-Grignon, France

The rapeseed crop has to cope with fungal diseases that impact significantly the yield. In the present context of reducing fungicides used in agriculture, it is essential to develop innovative and sustainable products to protect rapeseed crops. Currently, farmers do not have cost-effective and eco-friendly products to replace phytopharmaceuticals. Natural glycolipids, especially rhamnolipids from bacterial origin, were shown to have an elicitor effect (stimulation of defenses) in Arabidopsis, rapeseed and grapevine and antimicrobial properties on different pathogenic microorganisms^{1,2,3,4}. The present project aims to develop and optimize the properties of these compounds against the fungal pathogen *Leptosphaeria maculans*, the causal agent of blackleg disease on rapeseed.





In vitro experiments

Growth inhibition of *L. maculans* mycelium



Both R90 and 90L rhamnolipid solutions induce a significant growth inhibition of the fungus


Callose deposits on rapeseed



Callose deposits were observed on Basalti rapeseed cotyledons sprayed with both rhamnolipid solutions

In vivo experiments in controlled conditions


Variability of cultivar resistance



Lesion areas observed on 29-day-old rapeseed cotyledons. 12-day-old cotyledons are inoculated with *L. maculans* at 10⁶ spores/mL. Error bars represent the standard errors. (A) Lesion areas on untreated cotyledons (B) Percentage of lesion areas on cotyledons treated 2 days before inoculation with R90 at 0.1 g / L. Percentage is calculated from the untreated control of the corresponding cultivar set at 100%.

L. maculans sensitivity is cultivar-dependent. The variability is found when the plants are treated with the R90 solution before inoculation, illustrating a cultivar effect.


Protective effect of rhamnolipids



Lesion areas on 27-day-old cotyledons of Basalti cultivar, untreated (control) or treated with R90 (A) or 90L (B) solutions at different concentrations. Treatment is realized 2 days before inoculation and 12-day-old cotyledons are inoculated with *L. maculans* at 10⁶ spores/mL. Error bars represent the standard errors.

The preventive application of both rhamnolipid solutions triggers a plant protection with concentration-dependent intensity. Even at low concentration (0.025 g/L) the solutions are still very effective.

Effect of treatment timing on protection



Lesion areas on detached leaves from 35-day-old Basalti cultivar, untreated (control) or treated with R90 0.1 g/L. Treatment 2 days before inoculation; treatment and inoculation at the same time (T. + inoc); treatment 2 days after inoculation; treatment 7 days after inoculation. Twelve-day-old cotyledons are inoculated with *L. maculans* at 10⁶ spores/mL. Error bars represent standard errors.

The R90 rhamnolipid solution treatment triggers a protection of rapeseed cotyledons against *L. maculans* under controlled conditions when the treatment occurs 2 days before inoculation (preventive), in co-inoculation (direct antimycelium effect) and 7 days post-inoculation (curative effect).

Conclusions



Both rhamnolipid solutions show antifungal activity, eliciting properties and good adherence to rapeseed leaf. These properties lead to an effective protection against *L. maculans*, including concentrations where the antifungal activity is not significant. These results suggest a major role of the eliciting properties of these compounds in the rapeseed protection. Rhamnolipids therefore appear to be potentially cost-effective biocontrol products to fight fungal diseases of rapeseed.

References : ¹ Varnier *et al.*, *Plant. Cell Environ.*, 2009; ² Vatsa *et al.* *Int J Mol Sci.* 2010; ³ Sanchez *et al.*, *Plant Physiol.*, 2012; ⁴ Monnier *et al.*, *Front. Plant Sci.*, 2018

This work was carried out in partnership with SAS PIVERT, within the framework of the Institute for Energy Transition (ITE) P.I.V.E.R.T. (www.institut-pivert.com) selected as "Investissement d'Avenir". This work was funded by « Investissements d'Avenir » program by the French government under the reference ANR-001-01