



HAL
open science

Monanthosin, a new monoterpene from the leaves of *Monanthotaxis littoralis* (Annonaceae) with the antimicrobial and antioxidant activities of chemical constituents

Joseph Arnaud Dongmo Nguetse, Argan Kelly Nkwenti Wonkam, Jean-De-Dieu Tamokou, Steve Endeguele Ekom, Dominique Harakat, Laurence Voutquenne-Nazabadioko, David Ngnokam

► To cite this version:

Joseph Arnaud Dongmo Nguetse, Argan Kelly Nkwenti Wonkam, Jean-De-Dieu Tamokou, Steve Endeguele Ekom, Dominique Harakat, et al.. Monanthosin, a new monoterpene from the leaves of *Monanthotaxis littoralis* (Annonaceae) with the antimicrobial and antioxidant activities of chemical constituents. IOSR Journal of Applied Chemistry, 2020, 13 (12), pp.01-09. hal-03340605

HAL Id: hal-03340605

<https://hal.univ-reims.fr/hal-03340605v1>

Submitted on 16 Sep 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Monanthosin, a new monoterpene from *Monanthotaxis littoralis* (Annonaceae) with the antimicrobial and antioxidant activities of chemical constituents

Arnaud Joseph Nguetse Dongmo^a, Kelly Argan Wonkam Nkwenti^b, Jean-de-Dieu Tamokou^c, Steve Endeguele Ekom^c, Dominique Harakat^d, Laurence Voutquenne-Nazabadioko^e and David Ngnokam^{a,*},

^aResearch Unit of Applied and Environmental Chemistry, Department of Chemistry, Faculty of Science, University of Dschang, P.O. Box 67. Dschang Cameroon

^bDepartment of Organic Chemistry, Faculty of Science, University of Yaounde 1, P.O. Box 812 Yaounde Cameroon

^cResearch Unit of Microbiology and Antimicrobial Substances, Department of Biochemistry, Faculty of Science, University of Dschang, P.O. Box 67. Dschang Cameroon

^dService Commun d'Analyses, Institut de Chimie Moléculaire de Reims (ICMR), CNRS UMR 7312, Bat. 18 B.P. 1039, 51687 Reims Cedex 2, France

^eGroupe Isolement et Structure, Institut de Chimie Moléculaire de Reims (ICMR), CNRS UMR 7312, Bat. 18 B.P. 1039, 51687 Reims Cedex 2, France

*dngnokam@gmail.com

Abstract:

A new monoterpene, monanthosin (1), together with eight known compounds (chrysin (2), quercitrin (3), astilbin (4), heptulose (5), allantoin (6), heptitol (7), *cis-N-p*-coumaroyl tyramin (8) and *trans-N-p*-coumaroyl tyramin (9)) were isolated from the leaves of *Monanthotaxis littoralis* (Annonaceae). Structures were assigned by direct interpretation of their spectral data, mainly HR-TOFESIMS, 1-D NMR (¹H and ¹³C) and 2-D NMR (¹H-¹H COSY, HSQC, HMBC and NOESY) and by comparison with reported values. The MeOH, EtOAc and *n*-BuOH extracts as well as compounds **1**, **2**, **4** and **8** exhibited variable antimicrobial and antioxidant activities. The ethyl acetate extract and compound **4** were the most active samples among extracts and compounds, respectively. The ethyl acetate extract and antibiotics (vancomycin and fluconazole) demonstrated synergistic effect against *Escherichia coli*, *Pseudomonas aeruginosa*, *Candida albicans*, *Candida tropicalis* and *Cryptococcus neoformans* and additive effect against *Staphylococcus aureus*.

Keywords: *Monanthotaxis littoralis*; leaves; Annonaceae; Monanthosin; Antimicrobial, Antioxidant.

1. Introduction

The **Annonaceae family is characterized by the presence of** flavonoids, isoquinoline alkaloids and acetogenins (Biba et al., 2014; Chang et al.; 1998). Some members of *Monanthes* species contain oxygenated cyclohexane epoxide derivatives (Liang et al., 1998; Mulholland et al., 2000; Makhuvele et al., 2018), polyoxygenated cyclohexene (Starks et al., 2012), caryophyllene and caryophyllene-oxide (Parmena et al., 2012), **phenols** and triterpene (Ibrahim et al., 2016). *Monanthes littoralis* (Bagsh. & Baker F.) Verdc., is a persistent shrub with oblong elliptic leaves and solitary flowers («Botanique de *Popowia littoralis*» at [http://www.Plantes-botanique.org/espece-Popowia littoralis.](http://www.Plantes-botanique.org/espece-Popowia_littoralis.), February 20, 2015). It is a vascular plant widely distributed in some tropical African countries such as Cameroon, Central African Republic, Congo and Uganda (Chepkirui et al., 2014). Previous studies on this species reported the presence of flavonoids essential oils, (Chepkirui et al., 2014), roseoside and its related compounds (Yamamoto and Ito, 2005). In our continuous search for potentially interesting novel and bioactive secondary metabolites from Cameroonian medicinal plants (Foning et al., 2016; Joubouhi et al., 2015), we have examined the MeOH extract of the leaves of *M. littoralis*. In the present paper we report, the isolation and structural elucidation of a new monoterpene, monanthosin 1 together with eight known compounds (Fig. 1); the result of the antimicrobial and antioxidant activities of extracts and some of isolated compounds from *M. littoralis* (Tables 2–5) was also presented.

2. Methods and materials

2.1. General

^1H and ^{13}C -NMR spectra were performed in deuterated methanol on a Bruker AVANCE III 600 spectrometer equipped with a cryoprobe (^1H at 600 MHz and ^{13}C at 150 MHz). 2D NMR experiments were recorded by means of standard Bruker microprograms (Xwin-NMR version 2.1 software TopSpin 3.2). All chemical shifts (δ) are given in ppm with reference to tetramethylsilane (TMS) as internal standard and the coupling constants (J) are in Hz. TOF-ESIMS and HR-TOF-ESIMS spectra were recorded using a Micromass Q-TOF micro instrument (Manchester, UK) equipped with an electrospray source. The samples were introduced by direct infusion in a solution of MeOH at a rate of $5 \mu\text{L min}^{-1}$. The optical rotations were measured on a Bellingham & Stanley ADP 220 polarimeter (Bellingham + Stanley Ltd., United Kingdom). Column chromatography was run on Merck silica gel 60 (70–230 mesh) and gel permeation on Sephadex LH-20 while TLC was carried out on silica gel

GF₂₅₄ pre-coated plates with detection accomplished by spraying with 10% H₂SO₄ followed by heating at 90°C, or by visual inspection under UV lamp at 254 and 365 nm.

2.2. Plant material

The leaves of *M. littoralis* (Bagsh. & Baker F.) Verdc were collected in Dschang (5°27'0" N, 10°04'00" E, Menoua Division, Western Region of Cameroon), in January 2016. Authentication was done by Mr. Fulbert TADJOUTEU, a Botanist of the Cameroon National Herbarium, Yaounde, where the voucher specimen (N° 35048/HNC) has been deposited.

2.3. Extraction and isolation

The air-dried plant material (3.0 Kg) was powdered and extracted at room temperature with methanol (18 L, 72 h). Evaporation of solvent under reduced pressure yielded 620.9 g of crude extract (ML). Part of this extract (613.9 g) was extracted with ethyl acetate and *n*-butanol to give 206.9 g and 70.2 g dry fractions, respectively. The *n*-butanol extract (MLB) was fractionated by silica gel column chromatography (CC), eluting with gradient solvent system of EtOAc/MeOH (00-60% MeOH) to give seven fractions (MLB1-MLB7). Fraction MLB4 (3.57 g) was subjected to a silica gel CC and eluted with EtOAc/MeOH/H₂O (90/5/2, v/v/v) to provide three sub-fractions (MLB4.1-MLB4.3). Sub-fraction MLB4.3 (0.37 g) was submitted to silica gel CC, eluting with EtOAc/MeOH (98/2, v/v) to yield compound **1** (14.5 mg). Fraction MLB5 (10.5 g) was subjected to silica gel CC and eluted with EtOAc/MeOH (85/15, v/v) to give compounds **5** (19.4 mg) and **6** (20.5 g). The ethyl acetate extract (MLE) was fractionated by silica gel CC, eluting with gradient solvent system of hexane/EtOAc (15-100% EtOAc) to give nine fractions MLE1-MLE9. Fraction MLE2 (9.6 g) was loaded to a silica gel CC and eluted with hexane/EtOAc (85/15, v/v) to give compound **2** (15 mg). Fraction MLE5 (14.5 g) was purified by silica gel CC, eluting with hexane/EtOAc (60/40, v/v) to give compounds **8** (4.5 mg) and **9** (15 mg). Fraction MLE7 (8.0 g) was subjected to sephadex LH-20 and eluted with MeOH to give compound **4** (4.5 mg). Fraction MLE8 (15.3 g) was subjected to silica gel CC, eluted with EtOAc to provide six sub-fractions (MLE8.1-MLE8.6). Sub-fraction MLE8.6 (1.8 g) was separated by silica gel CC and eluted with EtOAc/MeOH (95/5, v/v) to give compound **3** (4.0 mg). Fraction MLE9 (8.5 g) was submitted by silica gel CC, eluted with hexane/EtOAc (85/15, v/v) to provide eighth sub-fractions (MLE9.1-MLE9.8). Sub-fraction MLE9.8 (2.45 g) was separated by silica gel CC and eluted with EtOAc/MeOH/ H₂O (90/5/5, v/v/v) to give six sub-fractions (MLE9.8a-MLE9.8f). Subfraction MLE9.8c (0.35 g) was subjected to silica gel CC and eluted with EtOAc/MeOH (98/2, v/v) to give compound **7** (17.0 mg).

2.4. Antimicrobial assay

2.4.1. Microorganisms

The studied microorganisms were one Gram-positive bacteria (*Staphylococcus aureus* ATCC 25923), two Gram-negative bacteria (*Escherichia coli* S2 (1) and *Pseudomonas aeruginosa* PA01) and three strains of yeasts (*Candida tropicalis* PK233, *Candida albicans* ATCC10231 and *Cryptococcus neoformans* H99) taken from our laboratory collection. The bacterial and fungal species were grown at 37 °C and maintained on nutrient agar (NA, Conda, Madrid, Spain) and Sabouraud Dextrose Agar (SDA, Conda) slants respectively.

2.4.2. Determination of minimum inhibitory concentration (MIC) and minimum microbicidal concentration (MMC)

MIC values were determined by a broth micro-dilution method as described earlier (Navickiene and Lopes, 2001) with slight modifications. Each test sample was dissolved in dimethylsulfoxide (DMSO) and the solution was then added to Mueller Hinton Broth (MHB) for bacteria or Sabouraud Dextrose Broth (SDB) for yeasts to give a final concentration of 8192 µg/mL. This was serially diluted twofold to obtain a concentration range of 0.125–4096 µg/mL. Then, 100 µL of each concentration were added in each well (96-well microplate) containing 95 µL of MHB or SDB and 5 µL of inoculum for final concentrations varying from 0.0625–2048 µg/mL. The inoculum was standardized at 2.5×10^5 cells/mL for yeasts and 10^6 CFU/mL for bacteria using a JENWAY 6105 UV/Vis spectrophotometer. The final concentration of DMSO in each well was <1% [preliminary analyses with 1% (v/v) DMSO did not inhibit the growth of the test organisms]. The negative control well consisted of 195 µL of MHB or SDB and 5 µL of the standard inoculum. The cultured micro plates were covered; then, the contents of each well were mixed thoroughly using a plate shaker (Flow Laboratory, Germany) and incubated at 35 °C for 24 h (bacteria) and 48 h (yeasts) under shaking. The assay was repeated three times. The MIC values of samples were determined by adding 50 µL of a 0.2 mg/mL *p*-iodonitrotetrazolium violet solution followed by incubation at 35 °C for 30 min. Viable microorganisms reduced the yellow dye to a pink color. MIC values were defined as the lowest sample concentrations that prevented this change in color indicating a complete inhibition of microbial growth. For the determination of MMC values, a portion of liquid (5 µL) from each well that showed no growth of microorganism was plated on Mueller Hinton Agar or SDA and incubated at 35 °C for 24 h (for bacteria) or 35 °C for 48 h (for yeasts). The lowest concentrations that yielded no growth after this subculturing were taken as the MMC values. Vancomycin (Sigma-Aldrich, Steinheim, Germany) and

fluconazole (Merck, Darmstadt, Germany) were used as positive controls for bacteria and yeasts, respectively.

2.4.3. Combined effect of the ethyl acetate extract and antibiotics.

The antimicrobial effects of a combination of the ethyl acetate extract of *Monanthotaxis littoralis* (EtOAcML), which exhibited the highest antimicrobial activity, and antibiotics were assessed by the checkerboard test as previously described (Climo et al., 1999). The antimicrobial combinations assayed included EtOAcML with antibiotics, vancomycin and fluconazole. Serial dilutions of three different antimicrobial agents were mixed in Mueller-Hinton broth. After 24-48 h of incubation at 37 °C, the MICs were determined as described above. The fractional inhibitory concentration (FIC) index was calculated according to the equation: FIC index = FICA + FICB = MIC of drug A in combination / MIC of drug A alone + MIC of drug B in combination / MIC of drug B alone. The FIC indices are the sum of the FICs of each of the drugs, which in turn is defined as the MIC of each drug when it is used in combination divided by the MIC of the drug when it is used alone. The interaction was defined as synergistic if the FIC index was less than or equal to 0.5, additive if the FIC index was greater than 0.5 and less than or equal 1.0, indifferent if the FIC index was greater than 1.0 and less than or equal to 2.0, and antagonistic if the FIC index was greater than 2.0. All the experiments were performed in triplicate.

2.5. Antioxidant assay

2.5.1. DPPH free radical scavenging assay

The free radical scavenging activity of extracts as well as most of their isolated compounds was performed according to (Brand-Williams et al., 1995) with slight modifications. Briefly, different concentrations (10 to 2000 µg/mL) of extracts/compounds and vitamin C (positive control) were thoroughly mixed with 3 mL of methanolic DPPH solution (20 mg/L) in test-tubes and the resulting solution was kept standing for 30 minutes at room temperature before the optical density (OD) was measured at 517 nm. The percentage radical scavenging activity was calculated from the following formula: % scavenging [DPPH] = $[(A_0 - A_1)/A_0] \times 100$ (Ikuo et al., 1991) Where A_0 was the absorbance of the negative control (methanolic DPPH solution) and A_1 was the absorbance in the presence of the samples. EC_{50} value was determined from the graph obtained using standard vitamin C by using the “ $y = mx + c$ ” formula from the slope of the graph. All the analyses were carried out in triplicate.

2.5.2. Gallic acid equivalent antioxidant capacity (GEAC) assay

The GEAC test was done as previously described by Rice-Evans and Miller with slight modifications (Rice-Evans and Miller, 1994). In a quartz cuvette, to 950 μL acetate buffer (pH = 5.0, 100 mM), the following were added: 20 μL laccase (1 mM stock solution), 20 μL test sample, 10 μL ABTS (2,2'-azinobis(3-ethylbenzothiazoline-6-sulfonic acid) (74 mM stock solution). The laccase were purified from *Sclerotinia sclerotiorum* according to the protocol described by (Mot et al., 2012). The sample concentrations in the assay mixture were 800, 400, 200, 100, 10 $\mu\text{g}/\text{mL}$ for the extracts and 200, 100, 50, 25, 12.5 $\mu\text{g}/\text{mL}$ for the isolated compounds. The content of the generated ABTS \bullet^+ radical was measured at 420 nm after 240 s reaction time and was converted to gallic acid equivalents by the use of a calibration curve (Pearson's correlation coefficient: $r = 0.996$) constructed with 0, 4, 10, 14, 28, 56, 84 μM gallic acid standards rather than Trolox. Experiments were done in triplicate.

2.6. Statistical analysis

Data were analyzed by one-way analysis of variance followed by Waller-Duncan Post Hoc test. The experimental results were expressed as the mean \pm Standard Deviation (SD). Differences between groups were considered significant when $p < 0.05$. All analyses were performed using the Statistical Package for Social Sciences (SPSS, version 12.0) software.

3. Results and discussion

3.1. Phytochemical analysis

The MeOH extract of the *M. littoralis* leaves was separated into two extracts by liquid/liquid chromatography, to give EtOAc soluble part and *n*-BuOH soluble part. Compounds **1**, **5** and **6** were then obtained by purification of the *n*-BuOH extract, while compounds **2-4**, and **7-9** were obtained by fractionation and purification of the EtOAc extract.

Compound **1** was obtained as a yellowish gum; $[\alpha]_{\text{D}}^{25} + 26$ (c 0.25, MeOH). ^1H NMR (600 MHz, CD_3OD) and ^{13}C NMR (150 MHz, CD_3OD): see Table 1. HR-ESI-MS (negative ion mode): m/z 253.1077 $[\text{M}-\text{H}]^-$ (Calcd for $\text{C}_{13}\text{H}_{18}\text{O}_5$, 253.1076), 507.2228 $[2\text{M}-\text{H}]^-$, 153.0915 $[\text{M}-\text{H}-\text{C}_4\text{H}_5\text{O}_3]^-$. The molecular formula $\text{C}_{13}\text{H}_{18}\text{O}_5$ indicates five degrees of unsaturation. The IR spectrum indicates the presence of hydroxyl and carbonyl groups, and carbon-carbon double bonds (see Experimental).

The ^{13}C NMR spectrum combined with the HSQC spectroscopic analysis displays 13 carbon: two carbonyl at δ_{C} 199.7 (C-3) and 177.7 (C-10), four olefinic carbons at δ_{C} 166.2 (C-5), 130.6 (C-7), 131.2 (C-8) and 125.7 (C-4), one sp^3 methine bearing oxygen at δ_{C} 73.7 (C-9), one methylene at δ_{C} 49.3 (C-2), three methyls at δ_{C} 22.0 (C-11), 23.1 (C-12) and 18.3

(C-13), and two quaternary aliphatic carbon at δ_C 41.2 (C-1) and δ_C 78.7 (C-6). The downfield shift observed for C-6, indicated that they are substituted by hydroxyl group.

The ^1H NMR confirms the presence of three singlet methyls of which one is vinylic (δ 1.93, 3H), and the two others were attached to the same sp^3 quaternary carbon (δ 1.06, 6H). The vinyl protons appear at δ 5.89 (1H, s), and 5.97 (2H, m). Signals observed as doublet at δ 2.17 and 2.56 were attributable to one methylenic proton. In the COSY spectrum, the two vinyl protons were correlated each other's and with the sp^3 methine proton bearing and oxygen at δ 4.66 (1H, brs).

Positions of methyl groups were deduced from its HMBC spectrum on which correlation between the protons at δ_H 1.06 (6H, H-11 and H-12) with carbons at δ_C 41.2 (C-1), 49.3 (C-2) and 78.7 (C-6), and proton at δ_H 1.93 (3H, H-13) with carbons at δ_C 78.7 (C-6), 125.7 (C-4) and 166.2 (C-5) were observed. The location of the carbonyl at C-3 position (δ_C 199.7) was deduced from its HMBC correlations with the methylenic protons H-2, and the vinylic proton H-4, thus indicating a conjugated ketone group. These correlations allowed us to build, in addition with the ^1H - ^1H COSY spectrum (Fig. 2), the carbon skeleton of the molecule.

Comparison of ^1H and ^{13}C data of monanthosin **1** with those of vomifoliol (Otsuka et al., 1995) and cucumegastigmane I (Kai et al., 2007) indicates that compound **1**, possess an intact fragment of cucumegastigmane I (Table 1). The difference was in the presence of a carbonyl group (δ_C 177.7) instead of an oxymethylene group in cucumegastigmane I. Thus, suggesting that compound **1** is the carbonyl derivative of cucumegastigmane I. This is confirmed in the MS spectrum giving an ion fragment at m/z 153.0915 ($\text{C}_9\text{H}_{13}\text{O}_2$) corresponding to the loss of the lateral chain in $\text{C}_4\text{H}_5\text{O}_3$ ($\text{C}_2\text{H}_2\text{-CHOH-COOH}$).

NOESY experiment gave no conclusive information on the absolute configurations around the C-6 and C-9 carbons. Only the correlations between H-4 and H-7, H-13 and H-11 were observed on this spectrum. Usually, cucumegastigmane and its derivatives have (6*S*, 9*R*)-configuration. The oxidation of C-10 carbon to carboxylic acid function in compound **1** would therefore reverse the absolute configuration around the C-9 carbon. This would allow us to suggest that the absolute configurations around the two stereocenters would be (6*S*, 9*S*). Horeau's method (Horeau, 1962; Schoofs and Horeau, 1977) was applied to **1** in order to confirm the configuration at C-9. A mixture of **1** with an excess of 2-phenylbutyric anhydride and DMAP in chloroform showed an immediate evolution of the optical rotation in the (-) sense, thus including the preferential esterification by the (+) antipode of the acid. Silica gel column chromatography coupled with an optical rotation detector (Chiral detector: Knauer France, reference: 1000) allowed the isolation of levorotatory 2-phenylbutyric acid.

According to the Horeau's method, when (-)-(*R*)-2-phenylbutyric acid accumulates in the mixture (i.e. when the (+)-(*S*)-acid is the preferential esterifying acid), the C-9 secondary hydroxyl has the (*S*) configuration. On the basis of aforementioned information, the structure of **1** was elucidated as (6*S*,7*E*,9*S*)-6,9-dihydroxy-3-oxo-4,7-megastigmadien-10-oic acid named monanthosin.

Structures of compounds **2**, **3**, **4**, **5**, **6**, **7**, **8** and **9** (Figure 1) were determined by means of spectroscopic data and by comparative analysis of their spectral data with those reported in the literature as known chrysin (**2**) (Larit et al., 2012), quercitrin (**3**) (Aderogba et al., 2013), astilbin (**4**) (Jusoh et al., 2013), heptulose (**5**) (Jaseja and Perlin, 1990), allantoin (**6**) (Sripathi et al., 2011), heptitol (**7**) (Lewis and Angyal, 1989), *cis-N-p*-coumaroyl tyramine (**8**) (Navickiene and Lopes, 2001) and *trans-N-p*-coumaroyl tyramine (**9**) isolated as a mixture of **8** and **9** (Ikuo et al., 1991; Nyaa et al., 2009).

3.2. Antimicrobial activity

In the present work, the extracts as well as four compounds isolated from the leaves of *M. littoralis* were tested for their antimicrobial activities against three bacterial (*Staphylococcus aureus* ATCC 25923, *Escherichia coli* S2 (1) and *Pseudomonas aeruginosa* PA01) and three fungal strains (*Candida albicans* ATCC10231, *Candida tropicalis* PK233 and *Cryptococcus neoformans* H99) using broth microdilution method (Nyaa et al., 2009) (Table 2). The MIC results indicated that the MeOH, *n*-BuOH and EtOAc extracts inhibited the growth of all tested bacterial species. The most active extract was the EtOAc extract (MIC = 64-256 µg/mL). Compound **1** displayed weak antibacterial activity (MIC = 128-256 µg/mL) and no antifungal activity (MIC = > 256 µg/mL). Compounds **2**, **4** and **8** inhibited the growth of all tested bacterial and fungal strains. Compound **4** (MIC = 8-16 µg/mL) was the most active with lowest MIC value of 8 µg/mL on *Staphylococcus aureus*, *Candida albicans*, *Candida tropicalis* and *Cryptococcus neoformans*, highlighting some medicinal potential for this compound. As shown in Table 2, vancomycin and fluconazole used as standard drugs were more potent than the tested samples against yeasts, Gram-positive and Gram-negative bacteria with the exception against *E. coli* and *P. aeruginosa* where the antibacterial activity of compound **4** was equal to or higher than that of vancomycin. The antibacterial activities of chrysin and astilbin are highest compared to those of the early reports (Wu et al., 2013; Moulari et al., 2006; Kuspradini et al., 2009). Indeed, the relative antibacterial activity (MIC₅₀ = 36.72 µg/ml) was recorded for chrysin against *E. coli* ATCC25922 (Wu et al., 2013) whereas astilbin had MIC values of 225 µg/ml against *Streptococcus sobrinus* (Kuspradini et al., 2009). The minimal inhibitory quantity (MIQs) of astilbin ranged from 50 to 100 µg against bacterial strains representative of skin microflora (Moulari et al., 2006).

The antimicrobial activity of phenolic conjugate coumaroyl tyramine can be explained by the fact that it has been suggested to have two possible roles in plant defence, as direct antimicrobial agents and in cell-wall reinforcement (Keller et al., 1996; Newman et al., 2001). However the antifungal activities of chrysin, astilbin and *cis-N-p*-coumaroyl tyramin are reported here for the first time.

The microbicidal activities of extracts and isolated compounds against susceptible strains were analysed by the minimum microbicidal concentration (MMC) assay and summarized as MMC/MIC ratios in Table 2. Indeed, an antimicrobial agent is considered microbicidal if the MMC is not more than fourfold higher than the MIC, i.e. $MMC/MIC \leq 4$ (Djouossi et al., 2015). The MeOH, EtOAc and *n*-BuOH extracts as well as compounds **2**, **4** and **8** were shown to be microbicidal ($MMC/MIC \leq 2$) against the susceptible microorganisms whereas compound **1** displayed the bacteriostatic/fungistatic character ($MMC/MIC > 4$) against all the tested microorganisms.

The combination of the ethyl acetate extract (EtOAcML), which exhibited the highest antimicrobial activity, and antibiotics (vancomycin and fluconazole) were assessed by the checkerboard test as previously described (Climo et al., 1999). EtOAcML and antibiotics demonstrated synergistic effect against *E. coli*, *P. aeruginosa*, *C. albicans*, *C. tropicalis* and *C. neoformans* and additive effect against *S. aureus* (Table 3).

3.3. DPPH radical scavenging activity

In this study, free radical scavenging capacities were measured using DPPH radical and ABTS radical cation. The results are expressed as gallic acid equivalent antioxidant capacity of tested samples (Table 4) and as equivalent concentrations of test samples scavenging 50% of DPPH radical (Table 4). In all, the DPPH and ABTS scavenging activities in this study indicated that the MeOH, EtOAc and *n*-BuOH extracts were potent antioxidants. On order to identify compounds responsible to this activity, the antioxidant properties of the flavonoids **2**, **4** and the *cis-N-p*-coumaroyl tyramine (**8**) as well as the new compound **1** were measured. No antioxidant activity was observed with compound **1** (results not shown), while compounds **2**, **4** and **8** were potent antioxidants. Compound **4** ($EC_{50} = 3.68 \mu\text{g/mL}$; $GEAC = 96.71 \mu\text{g/mL}$) exerted the greatest activity whereas compound **8** ($EC_{50} = 58.44 \mu\text{g/mL}$; $GEAC = 51.27 \mu\text{g/mL}$) displayed the lowest antioxidant activity in both assays ($p < 0.05$); suggesting that the

ability of these compounds to scavenge DPPH could also reflect their ability to inhibit the formation of ABTS⁺. However, their antioxidant activities are lesser than that of vitamin C (EC₅₀ = 1.96 µg/mL).

The antioxidant properties of chrysin and astilbin are in agreement to those of the literature (Pushpavalli et al., 2010; Vijayalakshmi et al., 2011). Indeed, Vijayalakshmi et al. (2011) demonstrated significant antioxidant activity of astilbin with IC₅₀ values of 7.50, 21.50 and 24.10 µg/ml against DPPH, nitric oxide and lipid peroxide radicals, respectively. A study conducted by Pushpavalli et al. (2010) showed that the treatment of d-galactosamine-intoxication rats with chrysin (25, 50 and 100mg/kg body weight) increased the activities of free-radical scavenging (enzymes superoxide dismutase, catalase and glutathione peroxidase) and the levels of non-enzymatic antioxidants (reduced glutathione, vitamin C and vitamin E); suggesting that chrysin acts as antioxidant agent. Chemical properties of chrysin, due to lack of oxygenation on B and C-ring are linked with various pharmacological properties that varies from antioxidant to anticancer properties (Habtemariam, 1997). Though, differences in the structure of flavones have been revealed to persuade the antioxidant property. Astilbin, which has 3',4' hydroxylation demonstrated more antioxidant activity than chrysin. This finding foresees potential applications of astilbin as an antioxidant. The presence of 3',4' hydroxylation, a double bond between carbons 2 and 3, and the presence of a carbonyl group on carbon 4 have been demonstrated to be crucial to generate antioxidant activity (Harris et al., 2006).

Acknowledgements

The authors are grateful to the University of Dschang for financing some consumables used in this work, to the “Service Commun d'Analyses” and “Groupe Isolement et Structure” of the “Institut de Chimie Moléculaire de Reims” for the spectroscopic and spectrometric analysis on the ESIMS and NMR equipment of the PIANet Platform. The EU-programme FEDER to the PIANet CPER project is gratefully acknowledged.

References

«Botanique de *Popowia littoralis*» available at [http://www.Plantes-botanique.org/espece-Popowia littoralis](http://www.Plantes-botanique.org/espece-Popowia_littoralis). (February 20, 2015).

- Aderogba, M.A., Ndhkala, A.R., Rengasamy, K.R.R., Staden, J.V., 2013. Antimicrobial and selected *in vitro* enzyme inhibitory effects of leaf extracts, flavonols and indole alkaloids isolated from *Croton menyharthii*. *Molecules* 18, 12633–12644.
- Biba, V.S., Amily, A., Sangeetha, S., Remani, P., 2014. Anticancer, antioxidant and antimicrobial activity of Annonaceae family. *World J. Pharm. Pharm. Sci.* 3, 1595–1604.
- Brand-Williams, W., Cuvelier, M., Berset, C., 1995. Use of a free radical method to evaluate antioxidant activity. *LWT-Food Sci. Technol.* 28, 25-30.
- Chang, F.R., Wei, J.L., Teng, C.M., 1998. Antiplatelets aggregation constituents from *Annona purpurea*. *J. Nat. Prod.* 61, 1457–1461.
- Chepkirui, C., Josphat, C.M., Isabel, N.W., Jesca, N., 2014. Antifungal activity of flavonoids isolated from *Monanthotaxis littoralis* against micotoxigenic fungi from maize. *Am. J. Chem. Appl.* 1, 54–60.
- Climo, M.W., Patron, R.L., Archer, G.L., 1999. Combinations of vancomycin and betalactams are synergistic against Staphylococci with reduced susceptibilities to vancomycin. *Antimicrob. Agents Chemother.* 43, 1747–1753.
- Djouossi, M. G., Tamokou, J. D. D., Ngnokam, D., Kuate, J-R., Tapondjou L. A., Harakat, D., Voutquenne-Nazabadioko, L., (2015). Antimicrobial and antioxidant flavonoids from the leaves of *Oncoba spinosa* Forssk. (Salicaceae). *BMC Complementary and Alternative Medicine* 15,134
- Foning, T.P.L., Mabou, F.D., Ngnokam, D., Harakat, D., Voutquenne-Nazabadioko, L., 2016. Owariensisone: a new iridolactone from the whole plant of *Brillantaisia owariensis* P. Beauv. *Nat. Prod. Res.* 30, 1611–1616.
- Habtemariam, S., 1997. Flavonoids as inhibitors or enhancers of the cytotoxicity of tumor necrosis factor- α in L-929 tumor cells. *J. Nat. Prod.* 60, 775–778.
- Harris, G.K., Qian, Y., Leonard, S.S., 2006. Luteolin and chrysin differentially inhibit cyclooxygenase-2 expression and scavenge reactive oxygen species but similarly inhibit prostaglandin-E-2 formation in RAW 264.7 cells. *J. Nutr.* 136, 1517-1521.

- Horeau, A., 1962. Détermination des configurations absolues par dédoublement partiel-II précisions et compléments. *Tetrahedron Lett.* 21, 965–969.
- Ibrahim, M., Rasoanaivo, L.H., Paltinean, R., Vlase, L., Raharisololalao, A., Crisan, G., 2016. Phenolic compounds and triterpene identified from *Monanthotaxis pilosa* Baillon. *Int. J. Chem. Stud.* 4, 22–27.
- Ikuo, M., Tamihide, M., Hiroe, K., Kazuhito, W., Hidetoshi, Y., 1991. Analysis and pharmacotoxicity of feruloyltyramine as a new constituent and *p*-coumaroyltyramine in *Cannabis sativa* L. *Pharmacolo. Biochem. Beha.* 40, 465–469.
- Jaseja, M., Perlin, A.S., 1990. Two-Dimensional NMR spectral study of the tautomeric equilibria of D-fructose and related compounds. *Magn. Reson. Chem.* 28, 283–289.
- Joubouhi, C., Mabou, F.D., Foning, T.P.L., Ngnokam, D., Harakat, D., Voutquenne-Nazabadioko, L., 2015. Five new iridoïd dimers from the fruits of *Canthium subcordatum* DC (*syn. Psydrax subcordata* DC). *Phytochem. Lett.* 13, 348–354.
- Jusoh, S., Zakaria, Z., Din, B.L., 2013. Isolation of astilbin from leaves of *Cratogeomys arborescens*. *Malaysian J. Anal. Sci.* 3, 430–435.
- Kai, H., Baba, M., Okuyama, T., 2007. Two new megastigmanes from the leaves of *Cucumis sativus*. *Chem. Pharm. Bull.* 55(1), 133–136.
- Keller, H., Hohlfeld, H., Wray, V., Hahlbrock, K., Scheel, D. and Strack, D. (1996) Changes in the accumulation of soluble and cell wall-bound phenolics in elicitor-treated cell suspension cultures and fungus-infected leaves of *Solanum tuberosum*. *Phytochem.* 42, 389–396.
- Kuspradini H, Mitsunaga T, Ohashi H. 2009. Antimicrobial activity against *Streptococcus sobrinus* and glucosyltransferase inhibitory activity of taxifolin and some flavanone rhamnosides from kempas (*Koompassia malaccensis*) extracts. *J Wood Sci* 55:308–313
- Larit, F., Benyahia, S., Benayache, S., Benayache, F., Léon, F., Brouard, I., Bermijo, J., 2012. Flavonoids from *Calycotome spinosa* (L.). *Lamk. Int. J. Med. Arom. Plants* 2, 34–37.

- Lewis, D., Angyal, S.J., 1989. ^1H -NMR spectra and conformations of three heptitols in deuterium oxide. *J. Chem. Soc., Perkin Trans. 2*. 0, 1763–1765.
- Liang, G.Y., Gray, A.I., Thomas, D.W., Waterman, P.G., 1988. Polyoxygenated cyclohexane epoxide derivatives from stem bark of *Monanthes bichananii*. *Phytochemistry* 27, 3857–3860.
- Makhuvele, R., Foubert, K., Apers, S., Pieters, L., Verschaeve, L., Elgorashi, E., 2018. Antimutagenic constituents from *Monanthes caffra* (Sond.) Verdc. *J. Pharm. Pharmacol.* 70, 976–984.
- Mot, A.C., Pârvu, M., Damian, G., Irimie, F.D., Darula, Z., Medzihradsky, K.F., Brem, B., Silaghi-Dumitrescu, R., 2012. A “yellow” laccase with “blue” spectroscopic features, from *Sclerotinia sclerotiorum*. *Process Biochem.* 47, 968-975.
- Moulari B, Pellequer Y, Lboutounne H, Girard C, Chaumont JP, Millet J, Muyard F. 2006. Isolation and in vitro antibacterial activity of astilbin, the bioactive flavanone from the leaves of *Harungana madagascariensis* Lam. ex Poir. (Hypericaceae). *J Ethnopharmacol* 106: 72–278.
- Mulholland, D., Naidoo, N., Hutchings, A., Lavaud, C., Massiot, G., 2000. Crotepoxide, a cyclohexane diepoxide from *Monanthes caffra*. *Biochem. Syst. Ecol.* 28, 595–597.
- Navickiene, H.M.D., Lopes, L.M.X., 2001. Alkamides and phenethyl derivatives from *Aristolochia gehrtii*. *J. Braz. Chem. Soc.* 12(4), 467–472.
- Newman, M.-A., von Roepenack-Lahaye, E., Parr, A., Daniels, M.J. and Dow, J.M. (2001) Induction of hydroxycinnamoyl-tyramine conjugates in pepper by *Xanthomonas campestris*, a plant defence response activated by hrp gene-dependent and hrp gene-independent mechanisms. *Mol. Plant-Microbe Interact.* 14, 785–792.
- Nyaa, T.B.L., Tapondjou, A.L., Barboni, L., Tamokou, J.D., Kuate, J.R., Tane, P., Park, H., 2009. NMR assignment and antimicrobial/antioxidant activities of 1-hydroxyeuscaphic acid from the seeds of *Butyrospermum parkii*. *Nat. Prod. Sci.* 15, 76–82.

- Otsuka, H., Yao, M., Kamada, K., Takeda, Y., 1995. Alangionosides G-M: Glycosides of megastigmane derivatives from the leaves of *Alangium premnifolium*. Chem. Pharm. Bull. 43, 754–759.
- Parmena, D.S., Mgina, C.A., Joseph, C.C., 2012. Composition of non volatile oils and antimicrobial activities of extracts from *Mononhotaxis discolor* and an undescribed *Uvariondendron* species. Tanz. J. Sci. 38, 221–231.
- Pushpavalli, G., Kalaiarasi, P., Veeramani, C., Pugalendi, K.V., 2010. Effect of chrysin on hepatoprotective and antioxidant status in D-galactosamine-induced hepatitis in rats. Eur. J. Pharmacol. 631(1-3), 36–41.
- Rice-Evans, C., Miller, N.J., 1994. Total antioxidant status in plasma and body fluids. Methods Enzymol. 234, 279-293.
- Schoofs, A., Horeau, A., 1977. Nouvelle méthode générale de détermination de la pureté énantiomérique et de la configuration des alcools secondaires chiraux. Tetrahedron Lett. 37, 3259–3262.
- Sripathi, K.S., Gopal, P., Lalitha, P., 2011. Allantoin from the leaves of *Pisonia grandis* R.Br. Int. J. Pharm. Life Sci. 2, 815–817.
- Starks, C.M., Williams, R.B., Rice, S.M., Norman, V.L., Lawrence, J.A., Goering, M.G., O’Neil-Johnson, M., Hu, J-F., Eldridge, G.R., 2012. Polyoxygenated cyclohexene derivatives from *Monanthotaxis congoensis*. Phytochemistry 74, 185–189.
- Vijayalakshmi, A., Ravichandiran, V., Jayakumari, S., Velraj, M., 2011. Antioxidant and anticonvulsant activity of astilbin, a flavonoid glycoside from the rhizome of *Smilax china* Linn. J. Pharm. Res. 4(3), 561–563.
- Wu, T., He, M., Zang, X., Zhou, Y., Qiu, T., Pan, S., & Xu, X. 2013. A structure–activity relationship study of flavonoids as inhibitors of *E. coli* by membrane interaction effect. *Biochimica et Biophysica Acta (BBA) - Biomembranes*, 1828(11), 2751–2756.
- Yamamo, Y., Ito, M., 2005. Synthesis of optically active vomifoliol and roseoside stereoisomers. Chem. Pharm. Bull. 53, 541–546.

Table 1: ^1H and ^{13}C NMR data of compounds **1**

Pos.	δ_{C} ^{a,b}	δ_{H} ^{a,c} (mult.; <i>J</i> in Hz)	COSY	HMBC
1	41.2	-		
2	49.3	2.17 (d, 16.5, H-2 α) 2.56 (d, 16.5, H-2 β)	H-2 β , 4, 13 H-2 α , 4, 13	C-1, 3, 11, 12
3	199.7	-		H-2, 11, 12
4	125.7	5.89 (s)	H-11	C-2, 6, 13
5	166.2	-		
6	78.7	-		
7	130.6	5.97 (m)	H-8	C-8, 9
8	131.2	5.97 (m)	H-7,9	C-7, 9
9	73.7	4.66 (br s)	H-8	
10	70.2	3.66 (m)		
11	22.9	1.06 (s)		C-1, 2, 3, 6, 12
12	21.8	1.06 (s)		C-1, 2, 3, 6, 11
13	18.3	1.93 (s)		C-4, 5, 6
14	53.7	3.58 (s)		C-10

^aRecorded in CD₃OD, ^b 150MHz, ^c600 MHz

Table 2. Antimicrobial activity (MIC and MMC in $\mu\text{g/mL}$) of extracts and, isolated compounds from *M. littoralis* and reference antimicrobial drugs.

Extracts/ Compounds	Inhibition parameters	<i>E. coli</i>	<i>P. aeruginosa</i>	<i>S. aureus</i>	<i>C. tropicalis</i>	<i>C. albicans</i>	<i>C. neoformans</i>
MeOH extract	MIC	256	512	256	512	512	256
	MMC	512	512	512	1024	1024	512
	MMC/MIC	2	2	2	2	2	2
EtOAc extract	MIC	64	64	64	256	256	256
	MMC	64	64	64	512	512	256
	MMC/MIC	1	1	1	2	2	1
<i>n</i> -BuOH extract	MIC	128	128	128	512	512	512
	MMC	128	128	256	1024	512	512
	MMC/MIC	1	1	2	2	1	1
1	MIC	256	256	128	>256	>256	>256
	MMC	>256	>256	>256	/	/	/
	MMC/MIC	/	/	/	/	/	/
2	MIC	64	32	32	64	64	32
	MMC	128	64	64	64	64	32
	MMC/MIC	2	2	2	1	1	1
4	MIC	16	16	8	8	8	8
	MMC	16	16	8	16	16	8
	MMC/MIC	1	1	1	2	2	1
8	MIC	64	32	16	64	32	16
	MMC	64	32	16	64	32	16
	MMC/MIC	1	1	1	1	1	1
Ref*	MIC	32	16	0.5	0.5	1	2
	MMC	32	16	0.5	0.5	1	2
	MMC/MIC	1	1	1	1	1	1

/: not determined; MIC: Minimum Inhibitory Concentration; MMC Minimum Microbicidal Concentration; *: fluconazole for yeasts and vancomycin for bacteria; compounds **1**, **5** and **6** were not active at concentrations up to 256 $\mu\text{g/mL}$; compounds **3**, **7** and **9** were not tested.

Table 3. Checkerboard assay of the ethyl acetate extract of *M. littoralis* (EtOAcML), fluconazole and vancomycin against pathogenic strains.

Strains	Agent	MIC ($\mu\text{g/mL}$)		FIC	FICI	Outcome
		Alone	Combination			
<i>E. coli</i>	EtOAcML	64	16	0.25	0.375	Synergistic
	Vancomycin	32	4	0.125		
<i>P. aeruginosa</i>	EtOAcML	64	8	0.125	0.25	Synergistic
	Vancomycin	16	2	0.125		
<i>S. aureus</i>	EtOAcML	64	32	0.50	0.75	Additive
	Vancomycin	0.5	0.125	0.25		
<i>C. tropicalis</i>	EtOAcML	256	32	0.125	0.25	Synergistic
	Fluconazole	0.50	0.062	0.125		
<i>C. albicans</i>	EtOAcML	256	64	0.25	0.375	Synergistic
	Fluconazole	1	0.125	0.125		
<i>C. neoformans</i>	EtOAcML	256	8	0.0312	0.0937	Synergistic
	Fluconazole	2	0.125	0.0625		

MIC: Minimum Inhibitory Concentration; FIC : fractional inhibitory concentration; FICI: fractional inhibitory concentration index

Table 4. antioxidant activities of extracts and some of isolated compounds from *M. littoralis* leaves

Extracts/compounds	DPPH free radical scavenging activity (EC ₅₀)	Gallic acid equivalent antioxidant capacity (GEAC)
MeOH extract	65.18 ± 0.29 ^a	62.02 ± 0.54 ^a
<i>n</i> -BuOH extract	69.07 ± 1.22 ^b	43.17 ± 0.56 ^b
EtOAc extract	74.21 ± 0.63 ^c	48.09 ± 0.36 ^c
2	27.41 ± 0.79 ^d	80.53 ± 0.82 ^d
4	3.68 ± 0.32 ^e	96.71 ± 0.41 ^e
8	58.44 ± 1.62 ^f	51.27 ± 0.72 ^f
Vitamin C	1.96 ± 0.14 ^g	NA

EC₅₀: Equivalent concentrations of test samples scavenging 50% of DPPH radical. Data represent the mean ± SD of three independent experiments carried out in triplicate. In the same column, values affected by different superscript letters (a-f) are significantly different according to one way ANOVA and Waller Duncan test; p < 0.05.




Fig. 1. Structures of the isolated compounds (1-9) from *M. littoralis*.




Fig. 2. The key HMBC and COSY correlation of 1.