



**HAL**  
open science

## Infection of Human Dental Pulp Stromal Cells by *Streptococcus mutans*: Shedding Light on Bacteria Pathogenicity and Pulp Inflammation

Elodie Maisonneuve, Julie Chevrier, Marie Dubus, Jennifer Varin, Johan Sergheraert, Sophie C Gangloff, Fany Reffuveille, Cédric Mauprivez, Halima Kerdjoudj

### ► To cite this version:

Elodie Maisonneuve, Julie Chevrier, Marie Dubus, Jennifer Varin, Johan Sergheraert, et al.. Infection of Human Dental Pulp Stromal Cells by *Streptococcus mutans*: Shedding Light on Bacteria Pathogenicity and Pulp Inflammation. *Frontiers in Cell and Developmental Biology*, 2020, 8, pp.785. 10.3389/fcell.2020.00785 . hal-03406275

**HAL Id: hal-03406275**

**<https://hal.univ-reims.fr/hal-03406275v1>**

Submitted on 27 Oct 2021

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License



# Infection of Human Dental Pulp Stromal Cells by *Streptococcus mutans*: Shedding Light on Bacteria Pathogenicity and Pulp Inflammation

Elodie Maisonneuve<sup>1</sup>, Julie Chevrier<sup>1</sup>, Marie Dubus<sup>1,2</sup>, Jennifer Varin<sup>1</sup>, Johan Sergheraert<sup>1,2,3</sup>, Sophie C. Gangloff<sup>1,4</sup>, Fany Reffuveille<sup>1,4</sup>, Cédric Mauprivez<sup>1,2,3</sup> and Halima Kerdjoudj<sup>1,2\*</sup>

<sup>1</sup> Université de Reims Champagne Ardenne, EA 4691, Biomatériaux et Inflammation en Site Osseux (BIOS), Reims, France, <sup>2</sup> Université de Reims Champagne Ardenne, UFR d'Odontologie, Reims, France, <sup>3</sup> Pôle Médecine bucco-dentaire, Hôpital Maison Blanche, Centre Hospitalier Universitaire de Reims, Reims, France, <sup>4</sup> Université de Reims Champagne Ardenne, UFR de Pharmacie, Reims, France

## OPEN ACCESS

### Edited by:

Vicente Herranz-Pérez,  
University of Valencia, Spain

### Reviewed by:

Jangho Kim,  
Chonnam National University,  
South Korea  
Liliana Burlibasa,  
University of Bucharest, Romania

### \*Correspondence:

Halima Kerdjoudj  
halima.kerdjoudj@univ-reims.fr

### Specialty section:

This article was submitted to  
Stem Cell Research,  
a section of the journal  
Frontiers in Cell and Developmental  
Biology

Received: 11 June 2020

Accepted: 27 July 2020

Published: 31 August 2020

### Citation:


Maisonneuve E, Chevrier J, Dubus M, Varin J, Sergheraert J, Gangloff SC, Reffuveille F, Mauprivez C and Kerdjoudj H (2020) Infection of Human Dental Pulp Stromal Cells by *Streptococcus mutans*: Shedding Light on Bacteria Pathogenicity and Pulp Inflammation. *Front. Cell Dev. Biol.* 8:785. doi: 10.3389/fcell.2020.00785

Cariogenic *Streptococcus mutans* (*S. mutans*) is implicated in the dental pulp necrosis but also in cardiovascular tissue infections. Herein, the purpose was to elucidate how human dental pulp derived stromal cells (DPSCs) react toward a direct interaction with *S. mutans*. DPSCs were challenged with *S. mutans*. Following 3 h of interaction, DPSCs were able to internalize *S. mutans* (rate < 1%), and F-actin fibers played a significant role in this process. *S. mutans* persisted in the DPSCs for 48 h without causing a cytotoxic effect. *S. mutans* was, however, able to get out of the DPSCs cytoplasm and to proliferate in the extracellular environment. Yet, we noticed several adaptive responses of bacteria to the extracellular environment such as a modification of the kinetic growth, the increase in biofilm formation on type I collagen and polyester fabrics, as well as a tolerance toward amoxicillin. In response to infection, DPSCs adopted a proinflammatory profile by increasing the secretion of IL-8, IL-1 $\beta$ , and TNF- $\alpha$ , strengthening the establishment of the dental pulp inflammation. Overall, these findings showed a direct impact of *S. mutans* on DPSCs, providing new insights into the potential role of *S. mutans* in infective diseases.

**Keywords:** dental pulp derived stromal cells, inflammation, internalization, *S. mutans*, pathogenicity

## INTRODUCTION

Among tooth colonizers, *Streptococcus mutans* (*S. mutans*) is considered as the primary etiologic agent of dental caries, an infectious disease that affects 60–90% of the population worldwide (Pitts et al., 2017). It is not unusual for *S. mutans* to seek refuge in dentin caries and to gain access to the bloodstream during dental procedures, causing opportunistic systemic infections (Nakano et al., 2006; Carinci et al., 2018). Therefore, bacteremia, through the adhesion to endocardium and collagenous matrix, is responsible amongst others of infective endocarditis and peripheral arterial disease (Abranches et al., 2009; Carinci et al., 2018; Conrads and About, 2018). Dental pulp is a vascularized and innervated connective tissue housed in mineralized dentine. Gronthos et al. isolated and characterized stromal cells from dental pulp tissue (DPSCs) (Gronthos et al., 2009).



DPSCs meet minimal core identity for mesenchymal stem cells (MSCs) immunophenotype as defined by International Society for Cellular Therapy (Dominici et al., 2006). As MSCs, DPSCs have pharmaceutical effects that are predominantly mediated by paracrine and contact factors arising from intrinsic MSCs physiological processes (Caplan and Correa, 2011; Wang et al., 2014; Zhou et al., 2020). DPSCs are able to elaborate defense mechanisms against infections with the capacity to participate in cellular defenses and modulate the overall response against microorganisms. Despite of these features, the dental pulp often succumbs to an overwhelming infection. This raises questions about the effect of *S. mutans* on impairing DPSCs defense mechanisms against infections and the underlying systemic infections.

The expression of the immunomodulatory factors in dental derived stromal cells such as DPSCs and dental follicles stem cells is influenced by bacterial products and Toll like receptor (TLR) agonists. Although the expression of the TLR family members in the DPSCs is well documented, their contribution in the immunomodulation is not yet well understood (Andruxhov et al., 2019). It was shown that TLR-3 agonist increases the secretion of TGF- $\beta$  and IL-6 by both DPSCs and dental follicles stem cells, while TLR-4 agonist increases TGF- $\beta$  production by the dental follicles stem cells, but not by DPSCs (Tomic et al., 2010). Stimulation of DPSCs for 48 h with *E. coli*-derived lipopolysaccharide enhances the expression of indolamine-2,3-dioxygenase-1 (Lee et al., 2016), but also increases the expression of pro-inflammatory mediators such as IL-8, IL-1 $\alpha$ , and TNF- $\alpha$  in dose dependent manner (Bindal et al., 2018). Despite this knowledge, functional interaction studies of DPSCs and oral pathogenic Gram-positive *S. mutans* and the subsequent inflammatory commitment are lacking.

The focus of our study was to investigate the interactions of human DPSCs with *S. mutans*. We hypothesized that DPSCs were able to internalize *S. mutans*, inducing behavioral changes in bacteria. In response to infection, DPSCs might adopt a proinflammatory profile, contributing to the establishment of the inflammation. These data support the concept that DPSCs serve as a protective niche for *S. mutans* in case of bacterial incursion into the dental pulp. In response, both DPSCs and *S. mutans* adapt their behavior in favor of the establishment of a dental pulp inflammation and bacterial infection.

## MATERIALS AND METHODS

### DPSCs, Fibroblasts, and *S. mutans* Cultures

Healthy pulp tissues were isolated from non-fully erupted third molars. Following the enzymatic digestion in collagenase and trypsin mixture, the extracted DPSCs were cultured in  $\alpha$ -MEM supplemented with 10% decomplexed fetal bovine serum (FBS), 1% Penicillin/Streptomycin/Amphotericin-B and 1% Glutamax<sup>®</sup> and maintained in a humidified atmosphere of 5% CO<sub>2</sub> at 37°C. Reaching sub-confluence, DPSCs were seeded at density of  $3 \times 10^3$  cell/cm<sup>2</sup> until the third passage. Human gingival derived fibroblasts (HGFs) were isolated from gingival fragments, obtained during teeth removal. HGFs were seeded at density of  $5 \times 10^3$  cells/cm<sup>2</sup> in DMEM supplemented with Glutamax<sup>®</sup> supplemented with 10% FBS and 1% Penicillin/Streptomycin, and maintained in a humidified atmosphere of 5% CO<sub>2</sub> at 37°C with a medium change every two days. Reaching sub-confluence, HGFs were amplified at density of  $10^4$  cell/cm<sup>2</sup> until the fourth passage. *S. mutans* Clarke-ATCC<sup>®</sup> 25175 was grown on Columbia agars with 5% sheep blood under anaerobic conditions using GenBag Anaer. For all experiments, *S. mutans* were grown for 24 h in Schaedler broth under aerobic conditions at 37°C.

### Eukaryote Cells/*S. mutans* Interaction

DPSCs and HGFs were seeded in 24-well plates at  $10^4$  cells/cm<sup>2</sup> and cultured in their corresponding medium. After 72 h of culture, cells were washed with DPBS and cultured, overnight, with antibiotic free culture medium. The next day, cells were washed twice with DPBS and 1 mL of antibiotic free culture medium was added. Following enumeration, DPSCs or HGFs were exposed to live *S. mutans* with a multiplicity of infection (MOI) of 30 bacteria: 1 cell. After 3 h of interaction, cells were washed with DPBS and incubated for 1 h with culture medium supplemented by 100  $\mu$ g/mL of amoxicillin (Panpharma) following antibiotic protection assay procedure. The count of intracellular and extracellular bacteria was determined. At the same time, infected DPSCs were maintained in culture for 24 and 48 h in 1 mL of antibiotic free culture medium. After that, conditioned media (CM) were divided: some parts were filtered and conserved at -80°C, and the rest were plated on blood agar to enumerate the extracellular bacteria.

### Cytochalasin-D Assay

DPSCs were seeded in 24-well plates at  $10^4$  cells/cm<sup>2</sup> and cultured for 72 h. Cells were washed with DPBS and cultured, overnight, in antibiotic free culture medium supplemented with 0.1  $\mu$ g/mL cytochalasin D (Sigma-Aldrich). Cytochalasin treated cells were incubated with *S. mutans* with a MOI of 30: 1. Following the antibiotic protection assay, the rate of the intracellular bacteria was determined.

### Intracellular Bacteria Quantification

Following 3 h of cell/bacteria interaction and 24 and 48 h culture post-infection ( $\pm$  cytochalasin D treatment), amoxicillin

protection assay was performed and cells were lysed with 0.1% Triton X-100 for 5 min. Cell lysates were plated on blood agar and cultured for 24 h under anaerobic conditions, at 37°C. The rate of viable intracellular *S. mutans* (i-*S. mutans*) was determined after bacterial count on agar plate.

### Extracellular Bacteria Quantification

Following 3 h of cell/bacteria interaction and 24 and 48 h culture post-infection ( $\pm$  cytochalasin D treatment), the harvested conditioned media were plated on blood agar and cultured for 24 h under anaerobic conditions, at 37°C. The rate of viable extracellular *S. mutans* (e-*S. mutans*) was determined after bacterial count on agar plate.

### Growth Monitoring

The bacteria growth was monitored for 72 h. Each isolated colony was added in 6 mL of Schaedler broth. The optical density (OD) of bacteria suspension was measured at 600 nm in a spectrophotometer cuve. Curves were achieved by calculating  $\ln(N/N_0)$  ( $N$  = OD at the analyzed time;  $N_0$  = starting OD) (Zwietering et al., 1990).

### Minimum Inhibitory Concentration (MIC) and Minimum Bactericidal Concentration (MBC)

Amoxicillin (100–0.10  $\mu\text{g/mL}$  range) and vancomycin (25–0.10  $\mu\text{g/mL}$  range) in Schaedler-broth were used. Bacteria ( $10^6$  bacteria/mL) were added and incubated at 37°C for 24 h. MIC values were determined as the lowest concentrations of antibiotic having an inhibitory effect on bacterial growth. After 24 h of incubation, the lowest concentration of the subculture (10  $\mu\text{L}$ ) showing no growth on blood agar was considered as the MBC.

### S. mutans Adhesion and Biofilm Formation on Type I Collagen (COL-I)

24-well plates were coated with 100  $\mu\text{g/mL}$  of COL-I and incubated with 500  $\mu\text{L}$  of bacteria suspension ( $10^6$  bacteria/mL) at 37°C. After 6 h of culture, adhered bacteria were exposed to ultrasound (40 kHz) for 5 min and a volume of 100  $\mu\text{L}$  from serial dilutions was plated on blood agars. After 24 h of culture the plates were gently washed and 2 mL of 0.2% crystal violet was added to determine the quantity of attached bacteria. After 20 min of incubation, unfixated crystal violet was removed and absolute ethanol was added. OD was measured at 595 nm. Controls were carried out without bacteria.

### S. mutans Adhesion to Polyester Fabrics and Scanning Electron Microscopy

The polyester braided fabrics, based on multifilament yarn, are typically used in cardiovascular applications (provided by CRITT-TJFU). Fabrics were incubated for 6 h with bacteria suspension ( $10^6$  bacteria/mL) at 37°C. Attached bacteria were determined as described above. Samples were fixed in 2.5% glutaraldehyde for 1 h, dehydrated in graded ethanol solutions for 10 min and desiccated in a drop of hexamethyldisilazane. After

air-drying at room temperature, samples were sputtered with a thin gold-palladium film. Samples were observed using Scanning Electron Microscope (SEM-JEOL-JSM-7900F).

### Lactate Dehydrogenase (LDH) Cytotoxicity Assay and DNA Quantification

LDH activity was evaluated in conditioned media of dental pulp derived stromal cells (DPSCs-CM) with the cytotoxicity detection kit following the manufacturer's instructions. OD was measured at 492 and 700 nm. DNA extraction was performed on DPSCs, 48 h post-infection using MasterPure<sup>TM</sup> DNA Purification Kit in accordance with the manufacturer's protocol. Extracted DNA was assessed by measuring the absorbance at 260 and 280 nm.

### Fluorescence Labeling

Before challenging, *S. mutans* were labeled using 4',6-diaminido-2-phenylindole (DAPI) and incubated for 10 min at 37°C. After 3 h of interaction, DPSCs were fixed with 4% paraformaldehyde at 37°C for 15 min and permeabilized with 0.5% Triton X-100 for 10 min. DPSCs were then incubated for 30 min at room temperature with Alexa<sup>®</sup> Fluor-488-conjugated-Phalloidin<sup>®</sup>. Stained cells were mounted and imaged by confocal microscopy (LSM 710, NLO, Zeiss, Germany).

### Cytokine Release by ELISA

DPSCs-CM were analyzed. Secreted levels of IL-8, TNF- $\alpha$ , and IL-1 $\beta$  were assessed using ELISA MAX<sup>TM</sup> Deluxe Set Human IL-8 and ELISA MAX<sup>TM</sup> Deluxe Set Human TNF- $\alpha$  and DuoSet<sup>®</sup> human IL-1 $\beta$ , respectively, according to the manufacturer's instructions. OD were measured at 450 nm using a microplate reader.

### Statistical Analysis

All statistical analyses were performed using GraphPad Prism 6 software. DPSCs experiments were performed with six-independent healthy donors. HGFs experiments were performed with three-independent healthy donors. Microbiological assays were performed at least three times in triplicates. Histograms represent mean  $\pm$  SEM and statistical analyses were performed using Mann & Whitney test. For each test, a value of  $p < 0.05$  was accepted as statistically significant.


## RESULTS AND DISCUSSION

Following a deep carious lesion, the dental pulp tissue establishes efficient strategies to hinder or even arrest the carious lesion progression and the bacterial incursion into the pulp (Conrads and About, 2018). In this study, DPSCs did not exert a direct antibacterial effect on *S. mutans*; instead, bacteria seem to have great ability to thrive under DPSCs environmental conditions (Supplementary Figure SI-1; LaRock and Nizet, 2015). *S. mutans* was found to adhere to the eukaryote cell membranes but without being invasive (Engels-Deutsch et al., 2003; Abranches et al., 2009; Berlutti et al., 2010; Kaufman and Skrtic, 2016). In

following, we investigated if *S. mutans* adhere to DPSCs and if the DPSCs might serve as a protective niche for *S. mutans* in case of bacterial incursion into the dental pulp. The rate of viable *S. mutans* showed that, following 3 h of contact, bacteria were able to invade DPSCs without affecting the cell integrity (**Figure 1**). To clarify the relationship between bacterial attachment and bacterial invasion, colony forming unit (CFU) counts were performed following extensive rinse with PBS or following amoxicillin-based antibiotic protection assay (**Figure 1A**). *S. mutans* is sensitive to amoxicillin and the addition of this antibiotic is expected to kill the extracellular bacteria but not the intracellular bacteria (Scaglione et al., 1993). The rate of the viable bacteria was < 1%, with very close CFU number with or without antibiotic protection assay (**Supplementary Figure SI-2**). These results suggest a neglecting number of adhered bacteria to DPSCs or a weak adhesion that was broken during washing procedure. Further quantification of bacterial adhesion forces to DPSCs should be conducted by using atomic force microscopy. Interestingly, HGFs were not able to prone to *S. mutans* adhesion and invasion. These results are in contradiction with (Berlutti et al., 2010) who demonstrated that 0.3% of *S. mutans* (ATCC® 25175) are able to adhere and invade gingival fibroblast cell line. To confirm the invasion ability of *S. mutans* into DPSCs, DAPI labeled bacteria were


used. DPSCs cytoskeleton was labeled and infected cells were observed by confocal microscopy. Infected DPSCs showed the presence of polymerized F-actin, signature of the preservation of the integrity of the adhered cells. The cross-sectional analysis of stacked confocal images revealed the presence of bacteria setting between F-actin, confirming cytoplasm invasion (**Figure 1B**). The plasma membrane of eukaryotic cells is a dynamic structure that coordinates the entry of macromolecules and cells by endocytic pathways (Gruenberg and van der Goot, 2006). To examine the role of F-actin and microtubules in *S. mutans* entry, DPSCs were treated with cytochalasin-D (i.e., known to cause microfilament depolymerization) for 24 h prior bacteria challenging. Compared to untreated DPSCs, the results showed that cytochalasin-D inhibited *S. mutans* uptake by ~ 82%, suggesting that F-actin is critical for *S. mutans* internalization (**Figure 1A**). To summarize, *S. mutans* are able to invade DPSCs. Despite a not detectable adhesion of *S. mutans* to DPSCs plasma membrane, F-actin fibers are required for *S. mutans* internalization.

Some pathogens could persist and proliferate in eukaryotic cells without affecting their integrity (Pizarro-Cerdá and Cossart, 2006). The intracellular persistence and proliferation of *S. mutans* was investigated by treating infected DPSCs for 1 h with amoxicillin and then leaving the DPSCs in antibiotic-free media for 24 and 48 h. *S. mutans* were able to survive in the



DPSCs over the study time (Figure 2A). The CFU of viable intracellular (i)-*S. mutans* was similar to those detected after 3 h of interaction. Interestingly, viable extracellular bacteria (e)-*S. mutans* were detected after 24 and 48 h of internalization. The recovery rates of e-*S. mutans* after 24 and 48 h were about 700% and 10,500% of the internalized *S. mutans*, respectively. Taken together, these results suggest that *S. mutans* are able to get out of cytoplasm and to proliferate in the extracellular environment. The most likely hypotheses are the death of infected DPSCs causing the bacteria release, or the externalization of *S. mutans* by infected DPSCs. It has been reported that lipoteichoic acid has a cytotoxic and apoptotic effect on DPSCs (Wang et al., 2001), while other reported an increase in the size of microtissue iMDP-3 and proliferation of the fibroblast following challenging with *S. mutans in vitro* (Kaufman and Skrtic, 2016; Nomura et al., 2016). Forty eight hours post-infection, lactate dehydrogenase (LDH) accumulation

in DPSCs-CM was measured and DNA of adhered DPSCs was quantified. LDH is a stable cytoplasmic enzyme present in most cells. During damage to cell cytoplasmic membranes, LDH is released from cells and into the surrounding cell-culture supernatant. Thus, the quantitation of LDH in cell-culture supernatant is one method by which investigators analyze cell-death levels. Compared to uninfected DPSCs, no significant differences in LDH accumulation and in DNA content were noticed (Figures 2B,C), suggesting a negligible DPSCs lysis following *S. mutans* internalization but also the absence of DPSCs proliferation. Thus, the absence of cytotoxic effect is in accordance with (Kaufman and Skrtic, 2016; Nomura et al., 2016) results. We examined whether cytoskeleton compounds are implicated in the bacteria release. By culturing infected DPSCs for 24 h in the presence of cytochalasin-D, viable e-*S. mutans* were detected, suggesting that cytoskeleton does not play a role in bacteria externalization (Figure 2D). However, the rates of both




**TABLE 1** | The Minimal Inhibitory Concentration (MIC) and Minimal Bactericidal Concentration (MBC) in  $\mu\text{g}/\text{mL}$  of amoxicillin and vancomycin to the intracellular (i-*S. mutans*), the extracellular (e-*S. mutans*), and the bacteria control (*S. mutans*) ( $n = 3$ ).

	Amoxicillin			Vancomycin		
	MIC	MBC	MBC: MIC ratio	MIC	MBC	MBC: MIC ratio
<i>S. mutans</i>	<0.10	0.10	< 1	0.78	12.5	16
i- <i>S. mutans</i>	0.10	0.78	8	0.39	0.39	1
e- <i>S. mutans</i>	0.10	6.25	64	0.39	3.12	8



i-*S. mutans* and e-*S. mutans* were lower than control (DPSCs in the absence of cytochalasin-D).

The pathogenicity and the virulence of both i-*S. mutans* and e-*S. mutans* isolated after 48 h post-infection was investigated and compared to the non-internalized bacteria. First, the growth curves were performed by transferring the harvested bacteria from the blood agar into fresh growth medium. In contrast to i-*S. mutans* and bacteria control that exhibited a monophasic growth pattern, e-*S. mutans* showed a biphasic growth pattern. Compared to bacteria control, i-*S. mutans* and e-*S. mutans* showed a longer lag phase (respectively, 16 and 24 h versus 6 h; **Figure 3**), indicating a delay in the bacteria growth following the contact with DPSCs. This could correspond to a period of adjustment that is needed for the establishment of the regulation of metabolic genes responsible for bacteria proliferation (LaRock and Nizet, 2015). Carbohydrates are common carbon and energy sources for all Streptococci. With certain pairs of sugar, a monophasic growth pattern was observed, whereas, others gave a biphasic growth pattern (i.e., diauxic growth) (Janda and Kuramitsu, 1978; Narang and Pilyugin, 2007). The diauxic lag reflects the time required for bacteria to build up the enzymes to sufficiently high levels, for the consumption of preferentially carbohydrate sources. Thus, the prolonged contact of *S. mutans* with DPSCs might affect enzyme synthesis and/or activity required for the carbohydrate catabolism.



Amoxicillin and vancomycin are effective antimicrobials against *S. mutans* and are commonly used in the antibiotic prophylaxis and/or in the antibiotic therapy of infective endocarditis (Habib et al., 2015; Limeres Posse et al., 2016). In 2019, a clinical study estimated the amoxicillin resistance was associated with higher mortality in a large cohort of patients with streptococci-related infective endocarditis (Pilmis et al., 2019). We investigated the susceptibility of e-*S. mutans*, i-*S. mutans* and bacterial control to amoxicillin and vancomycin, through the determination of MBC and MIC (**Table 1**). Amoxicillin showed a MBC: MIC ratio < 1 for *S. mutans* control, meanwhile, i-*S. mutans* and e-*S. mutans* exhibited an increase in MBC: MIC ratio (8 and 64, respectively). Vancomycin showed a MBC: MIC  $\sim 16$  for *S. mutans* and a decrease in MBC: MIC for both i-*S. mutans* and e-*S. mutans* (1 and 8, respectively). When the MBC is close to the MIC, the antibiotic is bactericidal; if MBC: MIC ratio is between 4 and 16, the antibiotic is bacteriostatic; and finally, if MBC: MIC ratio is > 32, we speak of “tolerance” (Brauner et al., 2016; Dafale et al., 2016). We can thus conclude that vancomycin had a bactericidal effect on i-*S. mutans* and bacteriostatic effect on e-*S. mutans*. Amoxicillin had a bacteriostatic effect on i-*S. mutans* while e-*S. mutans* seemed tolerant. This feature seems transitory and lost following the *in vitro* bacteria expansion. Antibiotic tolerance is associated with the failure of antibiotic treatment and the relapse of many bacterial infections (Brauner et al., 2016). The most likely hypothesis is the environmental changes and the metabolic adaptations of bacteria along with the prolonged lag phase of e-*S. mutans* (Brauner et al., 2016). We also cannot exclude the possibility of alteration or over expression of penicillin-binding-proteins (PBPs) in contact with DPSCs. Although the primary mode of action of vancomycin and amoxicillin is the inhibition of cell wall synthesis, their targets are different. Vancomycin binds very tightly to cytoplasmic peptides while amoxicillin binds PBPs (Watanakunakorn, 1984; Macheboeuf et al., 2006). Thus, the slower growth rate and a possible alteration of cell-wall could make e-*S. mutans* less susceptible to amoxicillin.

Bacteria can adhere and form biofilms (Lemos et al., 2019). The pathogenesis of *S. mutans* is associated with their adhesion to the extracellular matrix such as collagen and their ability to form biofilms on solid surfaces (i.e., valve implants). We thus analyzed the adhesion of i-*S. mutans* and e-*S. mutans* on either COL-I coating or polyester fabrics (i.e., raw material for valve implant and cardiovascular devices) (**Figure 4**). Compared to the control, i-*S. mutans* and e-*S. mutans* showed lower adhesion capabilities to COL-I (**Figure 4A**). Based on biofilm staining, the crystal violet assay highlighted that e-*S. mutans* increased significantly the formation of biofilm on COL-I versus i-*S. mutans* and control (**Figure 4B**). Bacterial biofilm is a community of adhered bacteria to a surface, which are embedded in a self-produced polymeric substance. Crystal violet is a basic protein dye that stains viable bacteria, dead bacteria, and polymeric substances. The low number of the adhered e-*S. mutans* on COL-I, and the high retention of the crystal violet dye by e-*S. mutans* suggested that these bacteria develop a higher capability to form a matrix versus i-*S. mutans* and control. On polyester fabrics, both i-*S. mutans* and e-*S. mutans* showed an increase



in adhesion capability compared to control (Figure 4C). Due to high non-specific fixation of crystal violet to the polyester fabrics, the biofilm was not determined. However, SEM pictures seem to indicate a matrix deposition on the polyester yarn by *e-S. mutans* (Figure 4D). Taken together, these results suggest that DPSCs interaction affects the surface motif adhesions of *S. mutans* strains.

During a carious disease, the host aims to fight the infection, through its immune-inflammatory response, and to restore the tooth structure *via* the DPSCs commitment. Healthy dental

pulp contains resident immune cells such as leukocytes that are able to detect the invaded pathogens and to release plethora of chemoattractive mediators (Farges et al., 2015). Studying the interaction of DPSCs with various components of both innate and adaptive immune systems showed that DPSCs and activated immune cells reciprocally regulate each other. However, DPSCs might adopt either an immunosuppressive or immunostimulatory phenotype depending on the level of inflammation (Andrukhov et al., 2019). The above results established that DPSCs survive to *S. mutans* infection. These



data prompted us to explore the presence of pro-inflammatory mediators in the DPSCs-CM. While *S. mutans* had a moderate effect on IL-1 $\beta$  secretion, we observed a significant increase in both IL-8 and TNF- $\alpha$  production following infection (Figure 5). Temporal and contextual profiles of IL-8, TNF- $\alpha$ , and IL-1 $\beta$  might exert negative effects on tissue repair leading to inflammation and pulp necrosis (Zanini et al., 2017). We can conclude that in contact with *S. mutans*, DPSCs get a pro-inflammatory profile, that strengthen the establishment of the dental pulp inflammation.

## CONCLUSION

This study provides new insights into the interaction between *S. mutans* and DPSCs. DPSCs internalize *S. mutans*, providing a protective niche during dental pulp infection. Internalized *S. mutans* were not able to proliferate into the cytoplasm, but were able to be externalized in the extracellular environment where they proliferated. A prolonged contact with DPSCs metabolites seems to induce adaptative profile and bacteria pathogenicity such as an increasing biofilm formation and an establishment of a tolerance toward amoxicillin. Although the cytotoxic effect of *S. mutans* toward DPSCs was not observed, these findings showed that *S. mutans* affect DPSCs paracrine activities by promoting the secretion of pro-inflammatory mediators. *S. mutans* has a high genetic variation and is classified into four serotypes (Bedoya-Correa et al., 2019). The aforementioned findings used one strain of *S. mutans* (ATCC 25175 strain). Therefore, in perspective, the use of clinical strains of *S. mutans* with various serotypes should be performed to confirm these results. Furthermore, to decipher the relationship between DPSCs and *S. mutans* pathogenicity, a genomic and proteomic analyses should be performed. Finally, the cross-talk between infected DPSCs and immune cells could bring additional insights about the inflammatory profile of DPSCs.

## REFERENCES

- Abranches, J., Zeng, L., Bélanger, M., Rodrigues, P. H., Simpson-Haidaris, P. J., Akin, D., et al. (2009). Invasion of human coronary artery endothelial cells by *Streptococcus mutans* OMZ175. *Oral Microbiol. Immunol.* 24, 141–145. doi: 10.1111/j.1399-302X.2008.00487.x
- Andrukhov, O., Behm, C., Blufstein, A., and Rausch-Fan, X. (2019). Immunomodulatory properties of dental tissue-derived mesenchymal stem cells: implication in disease and tissue regeneration. *World J. Stem Cells* 11, 604–617. doi: 10.4252/wjsc.v11.i9.604
- Bedoya-Correa, C. M., Rincón Rodríguez, R. J., and Parada-Sanchez, M. T. (2019). Genomic and phenotypic diversity of *Streptococcus mutans*. *J. Oral Biosci.* 61, 22–31. doi: 10.1016/j.job.2018.11.001
- Berlutti, F., Catizone, A., Ricci, G., Frioni, A., Natalizi, T., Valenti, P., et al. (2010). *Streptococcus mutans* and *Streptococcus sobrinus* are able to adhere and invade human gingival fibroblast cell line. *Int. J. Immunopathol. Pharmacol.* 23, 1253–1260. doi: 10.1177/039463201002300430
- Bindal, P., Ramasamy, T. S., Kasim, N. H. A., Gnanasegaran, N., and Chai, W. L. (2018). Immune responses of human dental pulp stem cells in lipopolysaccharide-induced microenvironment. *Cell Biol. Int.* 42, 832–840. doi: 10.1002/cbin.10938

## DATA AVAILABILITY STATEMENT

All datasets generated for this study are included in the article/Supplementary Material, further inquiries can be directed to the corresponding author.

## AUTHOR CONTRIBUTIONS

EM, JC, MD, JV, and JS contributed to conception, design, data analysis and interpretation, and drafted and critically revised the manuscript. FR, SG, CM, and HK contributed to data acquisition and interpretation and critically revised the manuscript. All authors gave final approval and agreed to be accountable for all aspects of the work.

## FUNDING

This work was supported by the Union des Blessés de la Face et de la Tête and Gueules Cassées Foundation (BioReg-program-58-2018). JC was supported by Institute Carnot MICA.

## ACKNOWLEDGMENTS

We are very grateful to the staff of the Core PICT, URCA (Dr. C. Terryn) and to CRITT-TJFU (Y. Khalsi) for providing polyester braided fabrics.

## SUPPLEMENTARY MATERIAL

The Supplementary Material for this article can be found online at: <https://www.frontiersin.org/articles/10.3389/fcell.2020.00785/full#supplementary-material>

- Brauner, A., Fridman, O., Gefen, O., and Balaban, N. Q. (2016). Distinguishing between resistance, tolerance and persistence to antibiotic treatment. *Nat. Rev. Microbiol.* 14, 320–330. doi: 10.1038/nrmicro.2016.34
- Caplan, A. I., and Correa, D. (2011). The MSC: an injury drugstore. *Cell Stem Cell* 9, 11–15. doi: 10.1016/j.stem.2011.06.008
- Carinci, F., Martinelli, M., Contaldo, M., Santoro, R., Pezzetti, F., Lauritano, D., et al. (2018). Focus on periodontal disease and development of *Endocarditis*. *J. Biol. Regul. Homeost. Agents* 32, 143–147.
- Conrads, G., and About, I. (2018). Pathophysiology of dental caries. *Monogr. Oral Sci.* 27, 1–10. doi: 10.1159/000487826
- Dafale, N. A., Semwal, U. P., Rajput, R. K., and Singh, G. N. (2016). Selection of appropriate analytical tools to determine the potency and bioactivity of antibiotics and antibiotic resistance. *J. Pharm. Anal.* 6, 207–213. doi: 10.1016/j.jpha.2016.05.006
- Dominici, M., Le Blanc, K., Mueller, I., Slaper-Cortenbach, I., Marini, F., Krause, D., et al. (2006). Minimal criteria for defining multipotent mesenchymal stromal cells: the international society for cellular therapy position statement. *Cytotherapy* 8, 315–317. doi: 10.1080/14653240600855905
- Engels-Deutsch, M., Pini, A., Yamashita, Y., Shibata, Y., Haikel, Y., Schöller-Guinard, M., et al. (2003). Insertional inactivation of Pac and RmlB genes reduces the release of tumor necrosis factor alpha, Interleukin-6, and Interleukin-8 Induced by *Streptococcus mutans* in monocytic, dental pulp, and

- periodontal ligament cells. *Infect. Immun.* 71, 5169–5177. doi: 10.1128/iai.71.9.5169-5177.2003
- Farges, J. C., Alliot-Licht, B., Renard, E., Ducret, M., Gaudin, A., Smith, A. J., et al. (2015). Dental Pulp defence and repair mechanisms in dental caries. *Med. Inflamm.* 2015:16.
- Gronthos, S., Mankani, M., Brahim, J., Robey, P. G., and Shi, S. (2009). Postnatal human dental pulp stem cells (DPSCs) in Vitro and in Vivo. *Proc. Natl. Acad. Sci. U.S.A.* 97, 13625–13630. doi: 10.1073/pnas.240309797
- Gruenberg, J., and van der Goot, F. G. (2006). Mechanisms of pathogen entry through the endosomal compartments. *Nat. Rev. Mol. Cell Biol.* 7, 495–504. doi: 10.1038/nrm1959
- Habib, G., Lancellotti, P., Antunes, M. J., Bongiorno, M. G., Casalta, J. P., Del Zotti, F., et al. (2015). 2015 ESC guidelines for the management of infective endocarditis: the task force for the management of infective endocarditis of the european society of cardiology (ESC) endorsed by: european association for cardio-thoracic surgery (EACTS), the european association of nuclear medicine (EANM). *Eur. Heart J.* 36, 3075–3128. doi: 10.1093/eurheartj/ehv319
- Janda, W. M., and Kuramitsu, H. K. (1978). Production of extracellular and cell-associated glucosyltransferase activity by *Streptococcus mutans* during growth on various carbon sources. *Infect. Immun.* 19, 116–122. doi: 10.1128/iai.19.1.116-122.1978
- Kaufman, G., and Skrtic, D. (2016). Structural and recovery mechanisms of 3D dental pulp cell microtissues challenged with *Streptococcus mutans* in extracellular matrix environment. *J. Med. Microbiol.* 65, 1332–1340. doi: 10.1099/jmm.0.000353
- LaRock, C. N., and Nizet, V. (2015). Cationic antimicrobial peptide resistance mechanisms of *Streptococcal* pathogens. *Biochim. Biophys. Acta* 1848, 3047–3054. doi: 10.1016/j.bbmem.2015.02.010
- Lee, S., Zhang, Q. Z., Karabucak, B., and Le, A. D. (2016). DPSCs from inflamed pulp modulate macrophage function via the TNF- $\alpha$ /IDO Axis. *J. Dent. Res.* 95, 1274–1281. doi: 10.1177/0022034516657817
- Lemos, J., Palmer, S., Zeng, L., Wen, Z., Kajfasz, J., Freires, L., et al. (2019). The biology of *Streptococcus mutans*. *Microbiol Spectr.* 7:1. doi: 10.1128/microbiolspec.GPP3-0051-2018
- Limeres Posse, J., Álvarez Fernández, M., Fernández Feijoo, J., Medina Henríquez, J., Lockhart, P. B., Chu, V. H., et al. (2016). Intravenous amoxicillin/clavulanate for the prevention of bacteraemia following dental procedures: a randomized clinical trial. *J. Antimicrob. Chemother.* 71, 2022–2030. doi: 10.1093/jac/dkw081
- Macheboeuf, P., Contreras-Martel, C., Job, V., Dideberg, O., and Dessen, A. (2006). Penicillin binding proteins: key players in bacterial cell cycle and drug resistance processes. *FEMS Microbiol. Rev.* 30, 673–691. doi: 10.1111/j.1574-6976.2006.00024.x
- Nakano, K., Inaba, H., Nomura, R., Nemoto, H., Takeda, M., Yoshioka, H., et al. (2006). Detection of cariogenic *Streptococcus mutans* in extirpated heart valve and atheromatous plaque specimens. *J. Clin. Microbiol.* 44, 3313–3317. doi: 10.1128/JCM.00377-376
- Narang, A., and Pilyugin, S. S. (2007). Bacterial gene regulation in diauxic and non-diauxic growth. *J. Theor. Biol.* 244, 326–348. doi: 10.1016/j.jtbi.2006.08.007
- Nomura, R., Ogaya, Y., and Nakano, K. (2016). Contribution of the collagen-binding proteins of *Streptococcus mutans* to bacterial colonization of inflamed dental pulp. *PLoS One* 11:e0159613. doi: 10.1371/journal.pone.0159613
- Pilmis, B., Lourtet-Hascoët, J., Barraud, O., Piau, C., Isnarde, C., Hery-Arnaud, G., et al. (2019). Be careful about MICs to amoxicillin for patients with *Streptococci*-related infective endocarditis. *Int. J. Antimicrob. Agents* 53, 850–854. doi: 10.1016/j.ijantimicag.2019.03.002
- Pitts, N. B., Zero, D. T., Marsh, P. D., Ekstrand, K., Weintraub, J. A., Ramos-Gomez, F., et al. (2017). Dental caries. *Nat. Rev. Dis. Primers* 3:17030. doi: 10.1038/nrdp.2017.30
- Pizarro-Cerdá, J., and Cossart, P. (2006). Bacterial adhesion and entry into host cells. *Cell* 124, 715–727. doi: 10.1016/j.cell.2006.02.012
- Scaglione, F., Demartini, G., Dugnani, S., and Frascini, F. (1993). A New model examining intracellular and extracellular activity of amoxicillin, azithromycin, and clarithromycin in infected cells. *Chemotherapy* 39, 416–423. doi: 10.1159/000238987
- Tomic, S., Djokic, J., Vasilijic, S., Vučević, D., Todorovic, V., Supic, G., et al. (2010). Immunomodulatory properties of mesenchymal stem cells derived from dental pulp and dental follicle are susceptible to activation by toll-like receptor agonists. *Stem Cells Dev.* 20, 695–708. doi: 10.1089/scd.2010.0145
- Wang, P. L., Shirasu, S., Daito, M., and Ohura, K. (2001). *Streptococcus mutans* lipoteichoic acid-induced apoptosis in cultured dental pulp cells from human deciduous teeth. *Biochem. Biophys. Res. Commun.* 281, 957–961. doi: 10.1006/bbrc.2001.4451
- Wang, Y., Chen, X., Cao, W., and Shi, Y. (2014). Plasticity of mesenchymal stem cells in immunomodulation: pathological and therapeutic implications. *Nat. Immunol.* 15, 1009–1016. doi: 10.1038/ni.3002
- Watanakunakorn, C. (1984). Mode of action and in-vitro activity of vancomycin. *J. Antimicrob. Chemother.* 14, 7–18. doi: 10.1093/jac/14.suppl\_d.7
- Zanini, M., Meyer, E., and Simon, S. (2017). Pulp inflammation diagnosis from clinical to inflammatory mediators: a systematic review. *J. Endod.* 43, 1033–1051. doi: 10.1016/j.joen.2017.02.009
- Zhou, L., Liu, W., Wu, Y., Sun, W., Dörfer, C. E., and Fawzy El-Sayed, K. M. (2020). Oral mesenchymal stem/progenitor cells: the immunomodulatory masters. *Stem Cells Int.* 2020, 16. doi: 10.1155/2020/1327405
- Zwietering, M. H., Jongenburger, I., Rombouts, F. M., and van't Riet, K. (1990). Modeling of the bacterial growth curve. *Appl. Environ. Microbiol.* 56, 1875–1881.

**Conflict of Interest:** The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2020 Maisonneuve, Chevrier, Dubus, Varin, Sergheraert, Gangloff, Reffuville, Mauprivez and Kerdjoudj. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.