



HAL
open science

Anti-Type VII Collagen Antibodies Are Identified in a Subpopulation of Bullous Pemphigoid Patients With Relapse

Delphine Giusti, Grégory Gatouillat, Sébastien Le Jan, Julie Plée, Philippe Bernard, Frank Antonicelli, Bach-Nga Pham

► **To cite this version:**

Delphine Giusti, Grégory Gatouillat, Sébastien Le Jan, Julie Plée, Philippe Bernard, et al.. Anti-Type VII Collagen Antibodies Are Identified in a Subpopulation of Bullous Pemphigoid Patients With Relapse. *Frontiers in Immunology*, 2018, 9, pp.570. 10.3389/fimmu.2018.00570 . hal-03424958

HAL Id: hal-03424958

<https://hal.univ-reims.fr/hal-03424958>

Submitted on 10 Nov 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution| 4.0 International License



Anti-Type VII Collagen Antibodies Are Identified in a Subpopulation of Bullous Pemphigoid Patients With Relapse

Delphine Giusti^{1,2†}, Grégory Gatouillat^{1,2†}, Sébastien Le Jan¹, Julie Plée^{1,3}, Philippe Bernard^{1,3†}, Frank Antonicelli^{1,4*†} and Bach-Nga Pham^{1,2†}

¹Laboratory of Dermatology, Faculty of Medicine, University of Reims Champagne-Ardenne, Reims, France, ²Laboratory of Immunology, Reims University Hospital, University of Reims Champagne-Ardenne, Reims, France, ³Department of Dermatology, Reims University Hospital, University of Reims Champagne-Ardenne, Reims, France, ⁴Department of Biological Sciences, Immunology, Faculty of Odontology, University of Reims Champagne-Ardenne, Reims, France

OPEN ACCESS

Edited by:

Ralf J. Ludwig,
University of Lübeck, Germany

Reviewed by:

Takashi Hashimoto,
Osaka City University, Japan
Jean Kanitakis,
Hospices Civils de Lyon, France
Hiroshi Koga,
Kurume University, Japan

*Correspondence:

Frank Antonicelli
frank.antonicelli@univ-reims.fr

[†]These authors have contributed
equally to this work.

Specialty section:

This article was submitted to
Immunological Tolerance
and Regulation,
a section of the journal
Frontiers in Immunology

Received: 12 January 2018

Accepted: 06 March 2018

Published: 21 March 2018

Citation:

Giusti D, Gatouillat G, Le Jan S, Plée J, Bernard P, Antonicelli F and Pham B-N (2018) Anti-Type VII Collagen Antibodies Are Identified in a Subpopulation of Bullous Pemphigoid Patients With Relapse. *Front. Immunol.* 9:570. doi: 10.3389/fimmu.2018.00570

Bullous pemphigoid (BP) is an autoimmune bullous skin disease characterized by anti-BP180 and anti-BP230 autoantibodies (AABs). Mucous membrane involvement is an uncommon clinical feature of BP which may evoke epidermolysis bullosa acquisita, another skin autoimmune disease characterized by anti-type VII collagen AABs. We therefore evaluated the presence of anti-type VII collagen AABs in the serum of BP patients with and without mucosal lesions at time of diagnosis and under therapy. Anti-BP180, anti-BP230, and anti-type VII collagen AABs were measured by ELISA in the serum of unselected patients fulfilling clinical and histo/immunopathological BP criteria at baseline ($n = 71$) and at time of relapse ($n = 24$). At baseline, anti-type VII collagen AABs were detected in 2 out of 24 patients with BP presenting with mucosal involvement, but not in patients without mucosal lesions ($n = 47$). At the time of relapse, 10 out of 24 BP patients either displayed a significant induction or increase of concentrations of anti-type VII collagen AABs ($P < 0.01$), independently of mucosal involvement. Those 10 relapsing BP patients were also characterized by a sustained high concentration of anti-BP180 AAb, whereas the serum anti-BP230 AAb concentrations did not vary in BP patients with relapse according to the presence of anti-type VII collagen AABs. Thus, our study showed that anti-type VII collagen along with anti-BP180 AABs detection stratified BP patients at time of relapse, illustrating a still dysregulated immune response that could reflect a potential epitope spreading mechanism in those BP patients.

Keywords: anti-type VII collagen antibodies, bullous pemphigoid, epitope spreading, mucous membrane involvement, relapse

INTRODUCTION

Bullous pemphigoid (BP) is the most common subepidermal autoimmune blistering skin disease that preferentially affects the elderly with various clinical manifestations. Clinically, BP patients typically present at diagnosis vesicles and tense clear blisters, which mainly occur on erythematous skin, together with erythematous or urticarial papules and plaques (1–4). BP is immunologically characterized by tissue-bound and circulating autoantibodies (AABs) directed against either the

BP antigen 180 (BP180) or the BP antigen 230 (BP230) or even both, which are components of hemidesmosomes involved in the dermal–epidermal cohesion (2, 3, 5–9). Most of BP patients (82–94%) display in their serum AAb that bind to the NC16A domain of the transmembrane protein BP180 (3, 10–16). AAbs against other antigenic sites of BP180 are associated with the severity and the phenotype of BP. Depending on the detection assay used, the presence of AAb against both the BP180 N- and C-terminal of the ectodomain was found to be associated with the presence of mucosal lesions (12), or not (17). When presenting with mucosal involvement, BP may suggest the BP-like, inflammatory form of epidermolysis bullosa acquisita (EBA), another skin autoimmune disease characterized by AAb directed against type VII collagen, a protein of the basement membrane zone beneath the stratified squamous epithelia (8, 18–22). However, no study has investigated yet the presence of serum anti-type VII collagen AAb at time of diagnosis of BP patients with mucosal involvement and whether these AAb are associated with BP outcome.

Loss of immune self-tolerance eventually leads to the generation of AAb (22). Among the different AAb in BP, the pathogenicity was mainly attributed to those directed against the NC16A domain of BP180 (9, 10, 23, 24). Actually, high serum level of anti-BP180 NC16A AAb was correlated with disease activity at time of diagnosis and was shown as an independent risk factor for BP relapse after cessation of therapy (14, 15, 25–27). In addition, approximately 30% of the BP patients relapse during the first year of treatment (16, 28–31). For those latter BP patients, it was shown that the variations of serum IgG AAb directed toward the BP180 NC16A domain after 2 months of therapy may be useful to predict BP outcome (16, 32). However up to now, a comparison in BP patients of the AAb profile at time of relapse vs. baseline has been investigated neither with respect to mucosal involvement nor with respect to antibody directed against other epidermal basement membrane autoantigens such as the type VII collagen.

In this study, we investigated certain autoimmunity markers which result from the disturbance of self-tolerance in BP both at baseline with respect to mucous membrane involvement and at time of relapse. To that purpose, the presence of AAb against type VII collagen was evaluated at time of diagnosis both in the serum of BP patients with and without mucosal involvement. Furthermore, serum anti-type VII collagen AAb titer at baseline and at time of relapse was analyzed according to the initial clinical BP phenotype at baseline and compared with anti-BP180 and anti-BP230 AAb serum profiles.

MATERIALS AND METHODS

Patients

This retrospective study was conducted in the Dermatology Department of the University Hospital of Reims, belonging to the French Reference Center for Autoimmune Bullous Diseases. From January 2011 to July 2015, 71 consecutive patients with newly diagnosed BP were included in this study when they fulfilled the following criteria: (1) blistering skin dermatosis fulfilling at least three of four clinical criteria for BP (33) and (2) linear IgG and/or C3 deposits along the epidermal basement membrane zone

by skin direct immunofluorescence microscopy of perilesional skin. Patients fulfilling less than three clinical criteria, were also included if they demonstrated high anti-BP180/230 titers and at least roof labeling by indirect immunofluorescence (IIF) on salt split skin (SSS). During a 1-year follow-up, the number and dates of potential relapses were recorded. Relapse was defined as the reappearance of at least three daily new blisters along with pruriginous, erythematous, or urticarial plaques (34). The disease activity was assessed at baseline using the Bullous Pemphigoid Disease Area Index (BPDAI) (34). Patients were separated into two groups and then analyzed according to the presence or the absence of mucosal involvement as recorded in the BPDAI evaluation.

AAbs Detection

For each patient, blood samples were collected in clot activator tubes. Sample collection was realized at the time of diagnosis in all BP patients, and at the time of relapse on therapy during the follow-up (mean time of 39 weeks after the beginning of treatment), or at an equivalent follow-up visit for patients with ongoing remission. All sera were stored at -20°C until analysis. The detection of serum anti-type VII collagen AAb was performed with a commercially available ELISA (MBL, Nagoya, Japan). This assay uses the NC1 and NC2 domains of type VII collagen as substrates (35). The 6 U/mL cutoff value recommended by the manufacturer was used. Anti-BP180-NC16A and anti-BP230 AAb were detected in serum samples using commercially available ELISA tests (MBL, Nagoya, Japan) (36, 37). ELISA values are expressed as Units per milliliter of serum with the cutoff value of 9 U/mL for both ELISAs as recommended by the manufacturer. Serum anti-basement membrane zone AAb were detected by IIF on monkey SSS, according to the manufacturer's instructions (Euroimmun, Lübeck, Germany).

Statistical Analysis

Quantitative variables were described as mean \pm SD and qualitative data as number and percentage. Comparisons between groups were performed using Wilcoxon rank test, Fisher exact test or χ^2 test, as appropriate. A P -value <0.05 was considered statistically significant. All analyses were performed using Microsoft Excel and GraphPad Prism software.

RESULTS

Clinical and Immunological Characteristics of BP Patients According to Mucosal Involvement at Baseline

To investigate the potential implication of serum anti-type VII collagen AAb at the time of diagnosis and during follow-up on therapy in patients with BP, 71 patients were included in the study. Clinical and immunological characteristics of BP patients according to mucosal involvement at baseline are detailed in **Table 1**. Among the 71 BP patients, 47 (66%) had a typical clinical presentation and 24 (34%) also had mucous membrane involvement, i.e., blisters or erosions of the oral cavity in all 24 cases in addition with genital erosions in 1 case. We evidenced that the

presence of oral lesions was associated with higher total (Table 1, $P < 0.001$) and skin BPDAl scores ($P < 0.01$). We also found that anti-BP180 AAb were indifferently present in BP patients with or without mucosal lesions, with similar titers (Table 1), while the percentage of patients with positive anti-BP230 antibodies titer was lower in patients with mucosal involvement compared with patients without mucosal involvement (29 and 55%, respectively; $P < 0.05$). Further investigation of AAb expression in BP patients with and without mucosal involvement showed that, although not statistically significant, anti-type VII collagen AAb were only detected at diagnosis in the serum of patients presenting mucosal lesions (2/24 patients with mucosal involvement vs. 0/47 in patients without mucosal involvement, Table 1). In those two patients, the presence of high titers of serum anti-BP180 (66 and 109 U/mL) but low titers of anti-type VII collagen (13 and 8 U/mL) AAb confirmed the diagnosis of BP.

Clinical and Immunological Characteristics of BP Patients With Relapse

Of the 71 BP patients included in this study, 56 had both a complete clinical and biological follow-up of at least 1 year, including

BPDAl scores and serum AAb detection every 2 months and also in case of relapse. At baseline, 24 out of those 56 BP patients experienced at least one relapse during this follow-up period. Both total and skin BPDAl scores tended to be higher for BP patients who relapsed than for BP patients who remained in clinical remission upon therapy (Table 2). Mucosal involvement at time of diagnosis was not more frequent in patients with later relapse than in BP patients with ongoing remission (46 and 31%, respectively; $P = 0.26$). For those 24 BP patients, the relapse occurred in a mean delay of 39 weeks after starting treatment, without any difference between patients with and without mucosal involvement at baseline (Table 2). At the time of relapse, the level of serum anti-type VII collagen AAb was higher compared with baseline values (mean value at relapse 7.8 U/mL vs. mean value at baseline 3.6 U/mL, $P < 0.01$) (Figure 1A). Such an increase was not observed in the serum of BP patients with ongoing remission when analyzed after a similar median duration of treatment (Figure 1B).

Anti-Type VII Collagen Antibody Positivity in a Subset of BP Patients at Time of Relapse

We then attempted to further characterize the subset of BP patients who relapsed during the follow-up. Serological analysis revealed that 10 (41%) of those 24 BP patients had increased serum levels of anti-type VII collagen AAb at time of relapse (mean value at time of relapse 15.4 U/mL) (Figure 1C), whereas the other relapsing patients did not (Figure 1D). Among those 10 BP patients with detectable serum anti-type VII collagen AAb at time of relapse, 6 had mucosal involvement at time of diagnosis. Other clinical features at baseline, including mean age, sex ratio, and BPDAl scores, were not different in patients with positive anti-type VII collagen at time of relapse from those who did not express this latter AAb (Table 3). Conversely, biological investigation of serum anti-BP180/230 AAb profiles between baseline and relapse highlighted that anti-BP180 ELISA scores remained elevated the 10 BP patients with concomitant positive anti-type VII collagen AAb, while they decreased in the subgroup of BP patients without anti-type VII collagen antibodies (Figures 2A,C). Such a difference in the serum AAb profiles was not observed when analyzing the variations in anti-BP230 antibody titers (Figures 2B,D).

TABLE 1 | Clinical and immunological characteristics of BP patients at baseline.

	Total	Patients without mucosal involvement	Patients with mucosal involvement	P value
Patients no.	71	47	24	
Mean age \pm SD (range), years	80.3 \pm 10.1 (45–95)	82.6 \pm 7.9 (53–95)	75.8 \pm 12.6 (45–92)	0.05
Sex ratio (female/male)	1.7	2.1	1.2	0.25
Total BPDAl (mean \pm SD)	46.0 \pm 27.4	38.0 \pm 22.9	61.3 \pm 29.7	0.001
Skin BPDAl (mean \pm SD)	44.8 \pm 26.1	38.0 \pm 22.9	57.9 \pm 28.2	0.004
Relapse, patients no. (%)	24 (33.8)	13 (27.6)	11 (45.8)	0.12
Serum autoantibodies				
COL7 Ab				
Positive ELISA value, No (%)	2 (2.8)	0 (0.0)	2 (8.3)	0.11
Mean \pm SD (U/mL)	2.6 \pm 1.8	2.4 \pm 1.3	3.1 \pm 2.6	0.29
BP180 Ab				
Positive ELISA value, no. (%)	59 (83.1)	40 (85.1)	19 (79.2)	0.52
Mean \pm SD (U/mL)	67.1 \pm 50.0	65.8 \pm 48.6	69.5 \pm 56.2	0.80
BP230 Ab				
Positive ELISA value, no. (%)	33 (46.5)	26 (55.3)	7 (29.2)	0.046
Mean \pm SD (U/mL)	26.2 \pm 36.9	31.4 (39.7)	16.1 \pm 28.8	0.10
IIF-SSS, no. (%)				
Roof labeling	47 (66.2)	33 (70.3)	14 (58.3)	
Roof and floor labeling	4 (5.6)	3 (6.4)	1 (4.2)	
Floor labeling	0 (0.0)	0 (0.0)	0 (0.0)	
Negative	20 (28.2)	11 (23.4)	9 (37.5)	

Clinical and immunological characteristics at baseline from the whole population and comparison of these characteristics between BP patients with and without mucosal involvement.

BPDAl, Bullous Pemphigoid Disease Area Index; COL7 Ab, anti-type VII collagen autoantibodies; BP180 Ab, anti-BP180 antibodies; BP230 Ab, anti-BP230 antibodies; IIF-SSS, indirect immunofluorescence on salt split skin; BP, bullous pemphigoid.

TABLE 2 | Clinical characteristics of BP patients at baseline according to disease outcome.

	Ongoing remission	Relapse	P value
Patients no.	32	24	
Mean age \pm SD (range), years	77.9 \pm 10.3 (53–90)	80.8 \pm 11.5 (45–95)	0.19
Sex ratio (female/male)	1.66	1.67	1.00
Total BPDAl (mean \pm SD)	42.3 \pm 28.8	54.4 \pm 25.0	0.08
Skin BPDAl (mean \pm SD)	40.3 \pm 27.5	52.6 \pm 24.9	0.06
Mucosal involvement, no. (%)	10 (31.2)	11 (45.8)	0.26

BPDAl, Bullous Pemphigoid Disease Area Index.




TABLE 3 | Clinical characteristics at baseline of BP patients who further relapsed under treatment.

	Patients with secondary appearance of COL7 Ab at relapse (N = 10)	Patients without secondary appearance COL7 Ab at relapse (N = 14)	P value
Mean age \pm SD (range), years	79.8 \pm 14.3 (45–95)	81.4 \pm 9.6 (57.95)	1.00
Sex ratio (female/male)	2.5	1.3	0.52
Total BPDAl (mean \pm SD)	57.3 \pm 34.0	52.3 \pm 17.0	0.72
Skin BPDAl (mean \pm SD)	54.2 \pm 32.4	51.5 \pm 16.7	1.00
Mucosal involvement, no. (%)	6 (60.0)	5 (35.7)	0.24


The clinical characteristics at baseline of BP patients who further relapsed under treatment were analyzed according to the secondary appearance of anti-type VII collagen autoantibodies at relapse (COL7 Ab).

BPDAl, Bullous Pemphigoid Disease Area Index; BP, bullous pemphigoid.

DISCUSSION

The present study demonstrated that investigating both anti-BP180 and anti-type VII collagen antibodies serum concentrations was useful to exclude EBA diagnosis in BP patients with mucosal involvement at time of diagnosis. Furthermore, anti-type VII collagen AAb were identified in a subgroup of BP patients at time of relapse.

At diagnosis, anti-type VII collagen antibodies were detected only in 2 of 24 BP patients with oral involvement. However, high serum anti-BP180 titers compared to those of anti-type VII collagen antibodies advocated for a BP rather than an EBA for which the presence of serum anti-type VII collagen antibody remains the immunological hallmark at baseline (20, 35, 38). Although serum anti-type VII collagen antibodies were detected in the subgroup of BP patients with mucosal involvement, such a low frequency of antibody expression cannot be considered as a biological marker of mucosal subepidermal blistering in BP. Of note, this very low frequency of BP patients with serum anti-type VII collagen AAb and their low titers are in accordance with previous studies evaluating the performance of anti-type VII collagen antibody ELISA, in which 1–8% of patients with BP were positively tested (35, 39, 40). This is also in setting with the low prevalence of anti-type VII collagen antibodies in other autoimmune and autoinflammatory diseases, such as inflammatory bowel disease (16%) and pemphigus (9.5%), but also in healthy subjects (38). Furthermore, the low titer of anti-type VII collagen AAb in comparison with the high titer of anti-BP180 antibodies in our two patients with mucous membrane involvement, like in these other diseases, rather points out the production of anti-type VII collagen antibodies in BP as an epiphenomenon, as previously suggested (31). A possible diversification of the AAb response in BP could be related at least in part to disease severity, as BP patients with mucosal involvement had higher total and skin



BPDAI scores, especially the blisters/erosions activity score, as compared with the BPDAI score from patients with a typical form of BP (41).

Serum anti-type VII collagen antibodies were evidenced in about 40% of patients at the time of relapse and their titers were increasing. This is in line with a previous study which showed that the variation in the AAb profile, called epitope spreading, occurred as an early event in about 50% of BP patients (42). Our results complete our knowledge on BP-associated epitope spreading, by showing that the production of AAb against type VII collagen occurred in relapsing patients but not in patients with ongoing remission. This further illustrated that targets of immune responses in BP can be extended not only to other epitopes on the hemidesmosome protein (42, 43) but also to other proteins in their vicinity. The production of anti-type VII collagen AAb at time of relapse was not related to mucosal involvement at baseline, but to disease severity. Of note, it has been previously showed that the main predictive risk factor of relapse is the number of new daily blister at baseline (16). By contrast, the increase in the skin BPDAI score in BP patients with mucosal involvement was rather related to erosions than to blister formation. Thus, anti-type VII collagen production may result from situations in which tissue damage, induced by proteases activity linked to blister formation, causes the release and exposure of a previously encrypted antigen, thereby leading to a secondary autoimmune response against the newly exposed antigen as proposed for other autoimmune diseases (38, 39).

Noteworthy, serum anti-type VII collagen AAb was evidenced only in a subgroup of BP patients at the time of relapse. Then, the presence of anti-type VII collagen AAb may be an associated risk factor characterizing this subgroup of relapsing BP patients. Interestingly, this subgroup was also characterized by a high persistent titer of BP180 antibodies. Anti-BP180 antibody concentration has been correlated with BP disease activity (14, 15, 25, 26), and the decrease in anti-BP180 AAb levels after 2 months of treatment was lower in patients with further relapse in comparison with patients with ongoing remission (16). In the present study, if the serum level of anti-BP180 antibody remained elevated in all BP patients at time of relapse, we noticed that BP relapsing patients with serum anti-type VII collagen AAb displayed a sustained level of anti-BP180 antibody, suggesting that the immune response is still highly active in those patients. We also previously showed that the inflammatory response remained elevated in BP patients who relapsed during the first year of treatment. Especially, the production of IL-17, IL-23, CXCL10, and ECP also remained elevated after 2 months of treatment in the serum of relapsing BP patients (44–47). Then, one can hypothesize that the inflammatory response may also display a specific profile in this subgroup of BP patients with serum anti-type VII collagen AAb and high serum level of anti-BP180 at time of relapse. Of note, regulatory T (Treg) cells, a major regulatory system of autoimmunity, demonstrated plasticity and can convert to Th₁₇ cells according to the cytokine environment (48). Accordingly, a previous study evidenced that Treg cells were upregulated (49),

whereas another study proposed a Treg cell downregulation in BP (50). Knowing that Treg activity and IL-17-producing cells may have opposite effects on autoimmunity (51, 52), an impaired Treg activity in an IL-17/IL-23 context (53, 54) could result in an imbalance between the pro-inflammatory and the regulatory cytokines levels which control the tolerance breakdown limit and therefore favors the production of anti-type VII collagen AAb and the sustained concentration of anti-BP180 AAb (47, 53, 55–57). In this line, it is worth to note that all of these inflammatory molecules promote matrix metalloproteinase MMP-9 and neutrophil elastase production and therefore participate to tissue degradation (44, 45, 53). Although further investigations are still needed to explain why some BP patients had circulating anti-type VII collagen AAb at time of relapse and not the other relapsing patients, our results support the hypothesis that chronic and dysregulated inflammation in line with persistent tissue damage and exposure of autoantigens may lead to tolerance breakdown and to autoimmunity.

In conclusion, we here showed the presence of AAb against the type VII collagen in the serum of relapsing BP patients who had a severe and difficult to treat disease. Actually, this is the first study demonstrating in those relapsing patients that the immune response is still dysregulated, probably due to prolonged epidermal/dermal damages which may sustain the immune tolerance breakdown process. Based on this observation, it will be interesting in a future prospective study to evaluate whether the autoimmune response spreads to other autoantigens identified in the subepidermal autoimmune-mediated blistering diseases such as the laminin-332 and the laminin gamma-1 chain (58, 59). Furthermore, it will also be of interest to determine whether the presence of serum anti-type VII collagen antibodies could be a predictive factor for relapse by analyzing the concentration of these AAbs at an earlier time point during the patients' follow-up in future prospective longitudinal studies. Finally, the pathogenicity of those anti-type VII collagen AAb in association

with anti-BP180 NC16A AAb will also need to be investigated in animal models of BP (21, 60).

ETHICS STATEMENT

This study was carried out in accordance with the recommendations of the “Commission Nationale de l'Informatique et des libertés (CNIL)” and approved by the Ethic Committee of the University Hospital of Reims (institutional review board 14-04-2009). All subjects gave written informed consent in accordance with the Declaration of Helsinki.

AUTHOR CONTRIBUTIONS

DG and GG wrote the main manuscript text. DG, GG, and SJ conducted the experiments and statistical analyses. PB and JP participated to patients' care and clinical follow-up. FA, PB, and B-NP supervised this work and manuscript redaction. All the authors reviewed the article.

ACKNOWLEDGMENTS

We thank Dr Michael Maizières for his technical assistance during the study.

FUNDING

This study was supported by the French Department of Health's “Projet Hospitalier de Recherche Clinique (PHRC) Interregional” 2009, and by the university of Reims Champagne-Ardenne. The funding source had no role in the design and conduct of the study; collection, management, analysis, and interpretation of data; preparation, review, or approval of the manuscript; and decision to submit the manuscript for publication.

REFERENCES

- Lever WF. Pemphigus. *Medicine* (1953) 32:1–123. doi:10.1097/00005792-195302000-00001
- Schmidt E, Zillikens D. Pemphigoid diseases. *Lancet* (2013) 381:320–32. doi:10.1016/S0140-6736(12)61140-4
- Bernard P, Antonicelli F. Bullous pemphigoid: a review of its diagnosis, associations and treatment. *Am J Clin Dermatol* (2017) 18(4):513–28. doi:10.1007/s40257-017-0264-2
- Amber KT, Murrell DF, Schmidt E, Joly P, Borradori L. Autoimmune subepidermal bullous diseases of the skin and mucosae: clinical features, diagnosis, and management. *Clin Rev Allergy Immunol* (2018) 54(1):26–51. doi:10.1007/s12016-017-8633-4
- Diaz LA, Ratrie H, Saunders WS, Futamura S, Squiquera HL, Anhalt GJ, et al. Isolation of a human epidermal cDNA corresponding to the 180-kD autoantigen recognized by bullous pemphigoid and herpes gestationis sera. Immunolocalization of this protein to the hemidesmosome. *J Clin Invest* (1990) 86:1088–94. doi:10.1172/JCI114812
- Labib RS, Anhalt GJ, Patel HP, Mutasim DF, Diaz LA. Molecular heterogeneity of the bullous pemphigoid antigens as detected by immunoblotting. *J Immunol* (1986) 136:1231–5.
- Stanley JR. Pemphigus and pemphigoid as paradigms of organ-specific, autoantibody-mediated diseases. *J Clin Invest* (1989) 83:1443–8. doi:10.1172/JCI114036
- Goletz S, Zillikens D, Schmidt E. Structural proteins of the dermal-epidermal junction targeted by autoantibodies in pemphigoid diseases. *Exp Dermatol* (2017) 26:1154–62. doi:10.1111/exd.13446
- Liu Y, Li L, Xia Y. BP180 is critical in the autoimmunity of bullous pemphigoid. *Front Immunol* (2017) 8:1752. doi:10.3389/fimmu.2017.01752
- Giudice GJ, Emery DJ, Zelicson BD, Anhalt GJ, Liu Z, Diaz LA. Bullous pemphigoid and herpes gestationis autoantibodies recognize a common non-collagenous site on the BP180 ectodomain. *J Immunol* (1993) 151:5742–50.
- Perriard J, Jaunin F, Favre B, Büdinger L, Hertl M, Saurat JH, et al. IgG autoantibodies from bullous pemphigoid (BP) patients bind antigenic sites on both the extracellular and the intracellular domains of the BP antigen 180. *J Invest Dermatol* (1999) 112:141–7. doi:10.1046/j.1523-1747.1999.00497.x
- Hofmann S, Thoma-Uszynski S, Hunziker T, Bernard P, Koebnick C, Stauber A, et al. Severity and phenotype of bullous pemphigoid relate to autoantibody profile against the NH2- and COOH-terminal regions of the BP180 ectodomain. *J Invest Dermatol* (2002) 119:1065–73. doi:10.1046/j.1523-1747.2002.19529.x
- Di Zenzo G, Thoma-Uszynski S, Fontao L, Calabresi V, Hofmann SC, Hellmark T, et al. Multicenter prospective study of the humoral autoimmune response in bullous pemphigoid. *Clin Immunol* (2008) 128:415–26. doi:10.1016/j.clim.2008.04.012
- Roussel A, Benichou J, Randriamanantany ZA, Gilbert D, Drenovska K, Houivet E, et al. Enzyme-linked immunosorbent assay for the combination of

- bullous pemphigoid antigens 1 and 2 in the diagnosis of bullous pemphigoid. *Arch Dermatol* (2011) 147:293–8. doi:10.1001/archdermatol.2011.21
15. Charneau J, Lorin J, Vitry F, Antonicelli F, Reguiai Z, Barbe C, et al. Usefulness of BP230 and BP180-NC16a enzyme-linked immunosorbent assays in the initial diagnosis of bullous pemphigoid: a retrospective study of 138 patients. *Arch Dermatol* (2011) 147:286–91. doi:10.1001/archdermatol.2011.23
 16. Fichel F, Barbe C, Joly P, Bedane C, Vabres P, Truchetet F, et al. Clinical and immunologic factors associated with bullous pemphigoid relapse during the first year of treatment: a multicenter, prospective study. *JAMA Dermatol* (2014) 150:25–33. doi:10.1001/jamadermatol.2013.5757
 17. Nakatani C, Muramatsu T, Shirai T. Immunoreactivity of bullous pemphigoid (BP) autoantibodies against the NC16A and C-terminal domains of the 180 kDa BP antigen (BP180): immunoblot analysis and enzyme-linked immunosorbent assay using BP180 recombinant proteins. *Br J Dermatol* (1998) 139:365–70. doi:10.1046/j.1365-2133.1998.02396.x
 18. Chen M, Kim GH, Prakash L, Woodley DT. Epidermolysis bullosa acquisita: autoimmunity to anchoring fibril collagen. *Autoimmunity* (2012) 45:91–101. doi:10.3109/08916934.2011.606450
 19. Kim JH, Kim YH, Kim S-C. Epidermolysis bullosa acquisita: a retrospective clinical analysis of 30 cases. *Acta Derm Venereol* (2011) 91:307–12. doi:10.2340/00015555-1065
 20. Ludwig RJ. Clinical presentation, pathogenesis, diagnosis, and treatment of epidermolysis bullosa acquisita. *ISRN Dermatol* (2013) 2013:812029. doi:10.1155/2013/812029
 21. Bieber K, Koga H, Nishie W. In vitro and in vivo models to investigate the pathomechanisms and novel treatments for pemphigoid diseases. *Exp Dermatol* (2017) 26:1163–70. doi:10.1111/exd.13415
 22. Ludwig RJ, Vanhoorelbeke K, Leyboldt F, Kaya Z, Bieber K, McLachlan SM, et al. Mechanisms of autoantibody-induced pathology. *Front Immunol* (2017) 8:603. doi:10.3389/fimmu.2017.00603
 23. Liu Z, Sui W, Zhao M, Li Z, Li N, Thresher R, et al. Subepidermal blistering induced by human autoantibodies to BP180 requires innate immune players in a humanized bullous pemphigoid mouse model. *J Autoimmun* (2008) 31:331–8. doi:10.1016/j.jaut.2008.08.009
 24. Zillikens D, Rose PA, Balding SD, Liu Z, Olague-Marchan M, Diaz LA, et al. Tight clustering of extracellular BP180 epitopes recognized by bullous pemphigoid autoantibodies. *J Invest Dermatol* (1997) 109:573–9. doi:10.1111/1523-1747.ep12337492
 25. Schmidt E, Obe K, Bröcker EB, Zillikens D. Serum levels of autoantibodies to BP180 correlate with disease activity in patients with bullous pemphigoid. *Arch Dermatol* (2000) 136:174–8. doi:10.1001/archderm.136.2.174
 26. Haase C, Büdinger L, Borradori L, Yee C, Merk HF, Yancey K, et al. Detection of IgG autoantibodies in the sera of patients with bullous and gestational pemphigoid: ELISA studies utilizing a baculovirus-encoded form of bullous pemphigoid antigen 2. *J Invest Dermatol* (1998) 110:282–6. doi:10.1038/sj.jid.5602955
 27. Bernard P, Reguiai Z, Tancrede-Bohin E, Cordel N, Plantin P, Pauwels C, et al. Risk factors for relapse in patients with bullous pemphigoid in clinical remission: a multicenter, prospective, cohort study. *Arch Dermatol* (2009) 145:537–42. doi:10.1001/archdermatol.2009.53
 28. Joly P, Roujeau J-C, Benichou J, Delaporte E, D'Incan M, Dreno B, et al. A comparison of two regimens of topical corticosteroids in the treatment of patients with bullous pemphigoid: a multicenter randomized study. *J Invest Dermatol* (2009) 129:1681–7. doi:10.1038/jid.2008.412
 29. Joly P, Roujeau J-C, Benichou J, Picard C, Dreno B, Delaporte E, et al. A comparison of oral and topical corticosteroids in patients with bullous pemphigoid. *N Engl J Med* (2002) 346:321–7. doi:10.1056/NEJMoa011592
 30. Kyriakis KP, Pappas VA, Panteleos DN, Tosca AD. Re-evaluation of the natural course of bullous pemphigoid. A prospective study. *Int J Dermatol* (1999) 38:909–13. doi:10.1046/j.1365-4362.1999.00754.x
 31. Brulefert A, Le Jan S, Plée J, Durlach A, Bernard P, Antonicelli F, et al. Variation of the epidermal expression of glucocorticoid receptor-beta as potential predictive marker of bullous pemphigoid outcome. *Exp Dermatol* (2017) 26:1261–6. doi:10.1111/exd.13444
 32. Cai SCS, Lim YL, Li W, Allen JC, Chua SH, Tan SH, et al. Anti-BP180 NC16A IgG titres as an indicator of disease activity and outcome in Asian patients with bullous pemphigoid. *Ann Acad Med Singapore* (2015) 44:119–26.
 33. Vaillant L, Bernard P, Joly P, Prost C, Labeille B, Bedane C, et al. Evaluation of clinical criteria for diagnosis of bullous pemphigoid. French Bullous Study Group. *Arch Dermatol* (1998) 134:1075–80. doi:10.1001/archderm.134.9.1075
 34. Murrell DF, Daniel BS, Joly P, Borradori L, Amagai M, Hashimoto T, et al. Definitions and outcome measures for bullous pemphigoid: recommendations by an international panel of experts. *J Am Acad Dermatol* (2012) 66:479–85. doi:10.1016/j.jaad.2011.06.032
 35. Marzano AV, Cozzani E, Fanoni D, De Pittà O, Vassallo C, Berti E, et al. Diagnosis and disease severity assessment of epidermolysis bullosa acquisita by ELISA for anti-type VII collagen autoantibodies: an Italian multicentre study. *Br J Dermatol* (2013) 168:80–4. doi:10.1111/bjd.12011
 36. Kobayashi M, Amagai M, Kuroda-Kinoshita K, Hashimoto T, Shirakata Y, Hashimoto K, et al. BP180 ELISA using bacterial recombinant NC16a protein as a diagnostic and monitoring tool for bullous pemphigoid. *J Dermatol Sci* (2002) 30:224–32. doi:10.1016/S0923-1811(02)00109-3
 37. Yoshida M, Hamada T, Amagai M, Hashimoto K, Uehara R, Yamaguchi K, et al. Enzyme-linked immunosorbent assay using bacterial recombinant proteins of human BP230 as a diagnostic tool for bullous pemphigoid. *J Dermatol Sci* (2006) 41:21–30. doi:10.1016/j.jdermsci.2005.11.002
 38. Licarete E, Ganz S, Recknagel MJ, Di Zenzo G, Hashimoto T, Hertl M, et al. Prevalence of collagen VII-specific autoantibodies in patients with autoimmune and inflammatory diseases. *BMC Immunol* (2012) 13:16. doi:10.1186/1471-2172-13-16
 39. Seta V, Aucouturier F, Bonnefoy J, Le Roux-Villet C, Pendaries V, Alexandre M, et al. Comparison of 3 type VII collagen (C7) assays for serologic diagnosis of epidermolysis bullosa acquisita (EBA). *J Am Acad Dermatol* (2016) 74:1166–72. doi:10.1016/j.jaad.2016.01.005
 40. Schmidt T, Hoch M, Lotfi Jad SS, Solimani F, Di Zenzo G, Marzano AV, et al. Serological diagnostics in the detection of IgG autoantibodies against human collagen VII in epidermolysis bullosa acquisita – a multicenter analysis. *Br J Dermatol* (2017) 177(6):1683–92. doi:10.1111/bjd.15800
 41. Clapé A, Muller C, Gatouillat G, Le Jan S, Barbe C, Pham BN, et al. Mucosal involvement in bullous pemphigoid is mostly associated with disease severity and to absence of anti-BP230 autoantibody. *Front Immunol* (2018) 9:479. doi:10.3389/fimmu.2018.00479
 42. Di Zenzo G, Thoma-Uszynski S, Calabresi V, Fontao L, Hofmann SC, Lacour J-P, et al. Demonstration of epitope-spreading phenomena in bullous pemphigoid: results of a prospective multicenter study. *J Invest Dermatol* (2011) 131:2271–80. doi:10.1038/jid.2011.180
 43. Chan LS, Vanderlugt CJ, Hashimoto T, Nishikawa T, Zone JJ, Black MM, et al. Epitope spreading: lessons from autoimmune skin diseases. *J Invest Dermatol* (1998) 110:103–9. doi:10.1046/j.1523-1747.1998.00107.x
 44. Plée J, Le Jan S, Giustiniani J, Barbe C, Joly P, Bedane C, et al. Integrating longitudinal serum IL-17 and IL-23 follow-up, along with autoantibodies variation, contributes to predict bullous pemphigoid outcome. *Sci Rep* (2015) 5:18001. doi:10.1038/srep18001
 45. Riani M, Le Jan S, Plée J, Durlach A, Le Naour R, Haegeman G, et al. Bullous pemphigoid outcome is associated with CXCL10-induced matrix metalloproteinase 9 secretion from monocytes and neutrophils but not lymphocytes. *J Allergy Clin Immunol* (2016) 139(3):863–72.e3. doi:10.1016/j.jaci.2016.08.012
 46. Giusti D, Gatouillat G, Le Jan S, Plée J, Bernard P, Antonicelli F, et al. Eosinophil Cationic Protein (ECP), a predictive marker of bullous pemphigoid severity and outcome. *Sci Rep* (2017) 7:4833. doi:10.1038/s41598-017-04687-5
 47. Giusti D, Le Jan S, Gatouillat G, Bernard P, Pham BN, Antonicelli F. Biomarkers related to bullous pemphigoid activity and outcome. *Exp Dermatol* (2017) 26(12):1240–7. doi:10.1111/exd.13459
 48. Zhou X, Tang J, Cao H, Fan H, Li B. Tissue resident regulatory T cells: novel therapeutic targets for human disease. *Cell Mol Immunol* (2015) 12:543–52. doi:10.1038/cmi.2015.23
 49. Gambichler T, Tsiolakidou A, Skrygan M, Höxtermann S, Susok L, Hessam S. T regulatory cells and other lymphocyte subsets in patients with bullous pemphigoid. *Clin Exp Dermatol* (2017) 42:632–7. doi:10.1111/ced.13135
 50. Antiga E, Quaglino P, Volpi W, Pierini I, Del Bianco E, Bianchi B, et al. Regulatory T cells in skin lesions and blood of patients with bullous pemphigoid. *J Eur Acad Dermatol Venereol* (2014) 28:222–30. doi:10.1111/jdv.12091

51. Bieber K, Sun S, Witte M, Kasprick A, Beltsiou F, Behnen M, et al. Regulatory T cells suppress inflammation and blistering in pemphigoid diseases. *Front Immunol* (2017) 8:1628. doi:10.3389/fimmu.2017.01628
52. Schmidt T, Sitaru C, Amber K, Hertl M. BP180- and BP230-specific IgG autoantibodies in pruritic disorders of the elderly: a preclinical stage of bullous pemphigoid? *Br J Dermatol* (2014) 171:212–9. doi:10.1111/bjd.12936
53. Le Jan S, Plée J, Vallerand D, Dupont A, Delanez E, Durlach A, et al. Innate immune cell-produced IL-17 sustains inflammation in bullous pemphigoid. *J Invest Dermatol* (2014) 134:2908–17. doi:10.1038/jid.2014.263
54. Arakawa M, Dainichi T, Ishii N, Hamada T, Karashima T, Nakama T, et al. Lesional Th17 cells and regulatory T cells in bullous pemphigoid. *Exp Dermatol* (2011) 20:1022–4. doi:10.1111/j.1600-0625.2011.01378.x
55. D'Auria L, Cordiali Fei P, Ameglio F. Cytokines and bullous pemphigoid. *Eur Cytokine Netw* (1999) 10:123–34.
56. D'Auria L, Mussi A, Bonifati C, Mastroianni A, Giacalone B, Ameglio F. Increased serum IL-6, TNF-alpha and IL-10 levels in patients with bullous pemphigoid: relationships with disease activity. *J Eur Acad Dermatol Venereol* (1999) 12:11–5. doi:10.1111/j.1468-3083.1999.tb00801.x
57. Ameglio F, D'Auria L, Bonifati C, Ferraro C, Mastroianni A, Giacalone B. Cytokine pattern in blister fluid and serum of patients with bullous pemphigoid: relationships with disease intensity. *Br J Dermatol* (1998) 138:611–4. doi:10.1046/j.1365-2133.1998.02169.x
58. Bernard P, Antonicelli F, Bedane C, Joly P, Le Roux-Villet C, Duvert-Lehembre S, et al. Prevalence and clinical significance of anti-laminin 332 autoantibodies detected by a novel enzyme-linked immunosorbent assay in mucous membrane pemphigoid. *JAMA Dermatol* (2013) 149:533–40. doi:10.1001/jamadermatol.2013.1434
59. Commin M-H, Schmidt E, Duvert-Lehembre S, Lasek A, Morice C, Estival J-L, et al. Clinical and immunological features and outcome of anti-p200 pemphigoid. *Br J Dermatol* (2016) 175:776–81. doi:10.1111/bjd.14629
60. Antonicelli F, Ludwig RJ. New insights into pemphigoid diseases. *Exp Dermatol* (2017) 26:1151–3. doi:10.1111/exd.13469

Conflict of Interest Statement: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2018 Giusti, Gatouillat, Le Jan, Plée, Bernard, Antonicelli and Pham. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.