



**HAL**  
open science

## Clinical reporting following the quantification of cerebrospinal fluid biomarkers in Alzheimer's disease: An international overview

Constance Delaby, Charlotte Teunissen, Kaj Blennow, Daniel Alcolea, Ivan Arisi, Elodie Bouaziz Amar, Anne Beaume, Aurélie Bedel, Giovanni Bellomo, Edith Bigot-corbel, et al.

### ► To cite this version:

Constance Delaby, Charlotte Teunissen, Kaj Blennow, Daniel Alcolea, Ivan Arisi, et al.. Clinical reporting following the quantification of cerebrospinal fluid biomarkers in Alzheimer's disease: An international overview. *Alzheimer's & Dementia: the Journal of the Alzheimer's Association*, In press, 10.1002/alz.12545 . hal-03599269

**HAL Id: hal-03599269**

**<https://hal.univ-reims.fr/hal-03599269>**

Submitted on 7 Apr 2022

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial - NoDerivatives | 4.0  
International License

## FEATURED ARTICLE

# Clinical reporting following the quantification of cerebrospinal fluid biomarkers in Alzheimer's disease: An international overview

Constance Delaby<sup>1,4</sup>  | Charlotte E. Teunissen<sup>2</sup> | Kaj Blennow<sup>3</sup> | Daniel Alcolea<sup>4</sup> | Ivan Arisi<sup>5</sup> | Elodie Bouaziz Amar<sup>6</sup> | Anne Beaume<sup>7</sup> | Aurélie Bedel<sup>8</sup> | Giovanni Bellomo<sup>9</sup> | Edith Bigot-Corbel<sup>10</sup> | Maria Bjerke<sup>11,18</sup> | Marie-Céline Blanc-Quintin<sup>12</sup> | Mercè Boada<sup>13,59</sup> | Olivier Bousiges<sup>14</sup> | Miles D Chapman<sup>15</sup> | Mari L. DeMarco<sup>16</sup> | Mara D'Onofrio<sup>5</sup> | Julien Dumurgier<sup>6</sup> | Diane Dufour-Rainfray<sup>17</sup> | Sebastiaan Engelborghs<sup>18,19</sup> | Hermann Esselmann<sup>20</sup> | Anne Fogli<sup>21</sup> | Audrey Gabelle<sup>1</sup> | Elisabetta Galloni<sup>22</sup> | Clémentine Gondolf<sup>23</sup> | Frédérique Grandhomme<sup>23</sup> | Oriol Grau-Rivera<sup>24,25,26,27</sup> | Melanie Hart<sup>15</sup> | Takeshi Ikeuchi<sup>28</sup> | Andreas Jeromin<sup>29</sup> | Kensaku Kasuga<sup>28</sup> | Ashvini Keshavan<sup>30</sup> | Michael Khalil<sup>31</sup> | Peter Körtvelyessy<sup>32</sup> | Agnieszka Kulczynska-Przybik<sup>33</sup> | Jean-Louis Laplanche<sup>6</sup> | Piotr Lewczuk<sup>33,34</sup> | Qiao-Xin Li<sup>35</sup> | Alberto Lleó<sup>4</sup> | Catherine Malaplate<sup>36</sup> | Marta Marquié<sup>13,59</sup> | Colin L. Masters<sup>35</sup> | Barbara Mroczko<sup>33</sup> | Léonor Nogueira<sup>37</sup> | Adelina Orellana<sup>13,59</sup> | Markus Otto<sup>38</sup> | Jean-Baptiste Oudart<sup>39</sup> | Claire Paquet<sup>6</sup> | Federico Paolini Paoletti<sup>9</sup> | Lucilla Parnetti<sup>9</sup> | Armand Perret-Liaudet<sup>40</sup> | Katell Peoc'h<sup>41</sup> | Koen Poesen<sup>42</sup> | Albert Puig-Pi Joan<sup>25,26</sup> | Isabelle Quadrio<sup>40</sup> | Muriel Quillard-Muraine<sup>43</sup> | Benoit Rucheton<sup>44</sup> | Susanna Schraen<sup>45</sup> | Jonathan M. Schott<sup>30</sup> | Leslie M. Shaw<sup>46</sup> | Marc Suárez-Calvet<sup>24,25,26,27</sup> | Magda Tsolaki<sup>47</sup> | Hayrettin Tumani<sup>38</sup> | Chinedu T Udeh-Momoh<sup>48</sup> | Lucie Vaudran<sup>49</sup> | Marcel M Verbeek<sup>50</sup> | Federico Verde<sup>51,52</sup> | Lisa Vermunt<sup>2</sup> | Jonathan Vogelgsang<sup>20,53</sup> | Jens Wiltfang<sup>20,54,55</sup> | Henrik Zetterberg<sup>3,56,57,58</sup> | Sylvain Lehmann<sup>1</sup>

<sup>1</sup> LBPC-PPC, Univ Montpellier, CHU Montpellier, INSERM, Montpellier, France<sup>2</sup> Neurochemistry Lab, Department of Clinical Chemistry, Amsterdam Neuroscience, Amsterdam UMC, Vrije Universiteit Amsterdam, Amsterdam, Netherlands<sup>3</sup> Institute of Neuroscience and Physiology, The Sahlgrenska Academy at the University of Gothenburg, Gothenburg, Sweden<sup>4</sup> Hospital de la Santa Creu i Sant Pau - Biomedical Research Institute Sant Pau - Universitat Autònoma de Barcelona, Barcelona, Spain<sup>5</sup> European Brain Research Institute (EBRI) "Rita Levi-Montalcini", Roma, Italy<sup>6</sup> Université de Paris, Cognitive Neurology Center, GHU APHP Nord Lariboisière Fernand-Widal Hospital, Paris, France

This is an open access article under the terms of the [Creative Commons Attribution-NonCommercial-NoDerivs](https://creativecommons.org/licenses/by-nc-nd/4.0/) License, which permits use and distribution in any medium, provided the original work is properly cited, the use is non-commercial and no modifications or adaptations are made.

© 2021 The Authors. *Alzheimer's & Dementia* published by Wiley Periodicals LLC on behalf of Alzheimer's Association

- <sup>7</sup> CHU Poitiers, Poitiers, France
- <sup>8</sup> CHU Bordeaux, Bordeaux, France
- <sup>9</sup> Lab of Clinical Neurochemistry, Section of Neurology, Dept. of Medicine and Surgery, University of Perugia, Perugia, Italy
- <sup>10</sup> Laboratoire de Biochimie, Hôpital G & R Laënnec, CHU de Nantes, Nantes, France
- <sup>11</sup> Vrije Universiteit Brussel, Center for Neurosciences and Department of Clinical Biology, Clinical Neurochemistry Laboratory, Universitair Ziekenhuis Brussel, Brussels, Belgium
- <sup>12</sup> APHP Centre—Université de Paris, Cochin Hospital, Paris, France
- <sup>13</sup> Research Center and Memory Clinic, Fundació ACE, Institut Català de Neurociències Aplicades and Universitat Internacional de Catalunya (UIC), Barcelona, Spain
- <sup>14</sup> Laboratoire de Biochimie et Biologie Moléculaire, et CNRS, ICube Laboratory UMR 7357 and FMTS (Fédération de Médecine Translationnelle de Strasbourg), Team IMIS, Hôpitaux Universitaires de Strasbourg, Strasbourg, France
- <sup>15</sup> Department of Neuroimmunology, National Hospital for Neurology and Neurosurgery, UCL Queen Square, London, UK
- <sup>16</sup> Department of Pathology and Laboratory Medicine, St. Paul's Hospital, Providence Health Care, Vancouver, Canada & Department of Pathology & Laboratory Medicine, University of British Columbia, Vancouver, Canada
- <sup>17</sup> CHU Tours, UMR 1253- iBrain, Tours, France
- <sup>18</sup> Department of Biomedical Sciences, Institute Born-Bunge, University of Antwerp, Antwerp, Belgium
- <sup>19</sup> Vrije Universiteit Brussel, Universitair Ziekenhuis Brussel, Center for Neurosciences and Department of Neurology, Brussels, Belgium
- <sup>20</sup> Department of Psychiatry and Psychotherapy, University Medical Center Goettingen (UMG, Goettingen, Germany)
- <sup>21</sup> CHU Clermont-Ferrand, Clermont-Ferrand, France
- <sup>22</sup> Dipartimento di Neuroscienze Divisione di Neurologia, Vicenza, Italy
- <sup>23</sup> CHU Cote de Nacre Caen, Caen, France
- <sup>24</sup> Barcelonaβeta Brain Research Center (BBRC), Pasqual Maragall Foundation, Barcelona, Spain
- <sup>25</sup> Servei de Neurologia, Hospital del Mar, Unitat de deteriorament cognitiu i trastorns del moviment, Barcelona, Spain
- <sup>26</sup> IMIM (Hospital del Mar Medical Research Institute), Barcelona, Spain
- <sup>27</sup> Centro de Investigación Biomédica en Red de Fragilidad y Envejecimiento Saludable (CIBERFES), Madrid, Spain
- <sup>28</sup> Dept. of Molecular Genetics, Center for Bioresources, Brain Research Institute, Niigata University, Niigata, Japan
- <sup>29</sup> Cohen Veterans Bioscience, Cambridge, Massachusetts, USA
- <sup>30</sup> Dementia Research Centre, UCL Queen Square Institute of Neurology, London, UK
- <sup>31</sup> Medical University of Graz, Graz, Austria
- <sup>32</sup> Freie Universität Berlin and Humboldt-Universität zu Berlin, Department of Neurology, German Center for Neurodegenerative Diseases, Magdeburg, Germany and Charité-Universitätsmedizin Berlin, Berlin, Germany
- <sup>33</sup> Department of Neurodegeneration Diagnostics, Medical University of Białystok, Białystok, Poland
- <sup>34</sup> Lab for Clinical Neurochemistry and Neurochemical Dementia Diagnostics, Universitätsklinikum Erlangen and Friedrich-Alexander Universität Erlangen-Nürnberg, Erlangen, Germany
- <sup>35</sup> Florey Institute and The University of Melbourne, Melbourne, Victoria, Australia
- <sup>36</sup> CHRU de Nancy, Laboratoire de Biochimie, Biologie Moléculaire et Nutrition/ Université de Lorraine, Nancy, France
- <sup>37</sup> Laboratoire de Biologie Cellulaire et Cytologie, CHU PURPAN, Toulouse, France
- <sup>38</sup> Department of Neurology and CSF Laboratory, University of Ulm, Ulm, Germany
- <sup>39</sup> CNRS UMR 7369, CHU Reims, Université de Reims Champagne-Ardenne, Reims, France
- <sup>40</sup> Lyon Neuroscience Research Center BIORAN Team - CNRS UMR 5292, INSERM U1028, Lyon University Hospital, Lyon, France
- <sup>41</sup> Université de Paris GHU APHP Nord Beaujon Hospital, Paris, France
- <sup>42</sup> Laboratory for Molecular Neurobiomarker Research (LaMoN), Department of Neurosciences, KU Leuven, Leuven Brain Institute, Leuven, Belgium
- <sup>43</sup> UNIROUEN, Rouen University Hospital, Department of Clinical biology, Biochemistry laboratory, Normandie Univ, Rouen, France
- <sup>44</sup> DMU BioGeM, AP-HP Sorbonne University, Paris, France
- <sup>45</sup> Inserm, CHU Lille, U1172-LiNCog, LICEND, LabEx DISTALZ, Université de Lille, Lille, France
- <sup>46</sup> Department of Pathology and Laboratory Medicine Hospital, University of Pennsylvania, Pennsylvania, USA
- <sup>47</sup> 1st Department of Neurology, School of Medicine, Faculty of Health of Sciences, Aristotle University of Thessaloniki, Thessaloniki, Greece
- <sup>48</sup> Imperial College London, London, UK
- <sup>49</sup> CHU Lille, Univ. Lille, Lille, France
- <sup>50</sup> Donders Institute for Brain, Cognition and Behaviour, Radboud Alzheimer Centre, Departments of Neurology and Laboratory Medicine, Radboud University Medical Center, Nijmegen, The Netherlands
- <sup>51</sup> Department of Neurology - Stroke Unit and Laboratory of Neuroscience, IRCCS Istituto Auxologico Italiano, Milan, Italy

<sup>52</sup> Department of Pathophysiology and Transplantation, "Dino Ferrari" Center, Università degli Studi di Milano, Milan, Italy

<sup>53</sup> McLean Hospital, Translational Neuroscience Laboratory, Harvard Medical School, Belmont, Massachusetts, USA

<sup>54</sup> German Center for Neurodegenerative Diseases (DZNE), Goettingen, Germany

<sup>55</sup> Neurosciences and Signaling Group, Institute of Biomedicine (iBiMED), Department of Medical Sciences, University of Aveiro, Aveiro, Portugal

<sup>56</sup> Clinical Neurochemistry Laboratory, Sahlgrenska University Hospital, Mölndal, Sweden

<sup>57</sup> UK Dementia Research Institute at UCL, London, UK

<sup>58</sup> Department of Neurodegenerative Disease, UCL Institute of Neurology, London, UK

<sup>59</sup> Centro de Investigación Biomédica en Red de Enfermedades Neurodegenerativas (CIBERNED), Instituto de Salud Carlos III, Madrid, Spain

#### Correspondence

Sylvain Lehmann, Laboratoire de Biochimie Protéomique Clinique, CHU Montpellier, Hôpital St Eloi, IRMB 80 av A Fiche, 34295 Montpellier, France.  
Email: [s-lehmann@chu-montpellier.fr](mailto:s-lehmann@chu-montpellier.fr)

#### Abstract

**Introduction:** The current practice of quantifying cerebrospinal fluid (CSF) biomarkers as an aid in the diagnosis of Alzheimer's disease (AD) varies from center to center. For a same biochemical profile, interpretation and reporting of results may differ, which can lead to misunderstandings and raises questions about the commutability of tests.

**Methods:** We obtained a description of (pre-)analytical protocols and sample reports from 40 centers worldwide. A consensus approach allowed us to propose harmonized comments corresponding to the different CSF biomarker profiles observed in patients.

**Results:** The (pre-)analytical procedures were similar between centers. There was considerable heterogeneity in cutoff definitions and report comments. We therefore identified and selected by consensus the most accurate and informative comments regarding the interpretation of CSF biomarkers in the context of AD diagnosis.

**Discussion:** This is the first time that harmonized reports are proposed across worldwide specialized laboratories involved in the biochemical diagnosis of AD.

#### KEYWORDS

Alzheimer's disease, cerebrospinal fluid biomarkers, clinical report, consensus approach, harmonization

## 1 | INTRODUCTION

Alzheimer's disease (AD) has gradually become one of the major global public health issues due to its prevalence, which increases with age and life expectancy, and the economic cost of caring for patients whose cognitive decline progressively leads to loss of functional autonomy.<sup>1</sup>

The diagnosis of AD is based on a multidisciplinary approach, involving, among other things, evaluation of the medical history together with clinical symptoms and signs, neuropsychological tests, and neuroimaging. The quantification of cerebrospinal fluid (CSF) core biomarkers (amyloid beta peptides [ $A\beta_{1-40}$  and  $A\beta_{1-42}$ ], total tau [t-tau] and its phosphorylated form on threonine 181 [p-tau(181)]) has progressively proven useful for the diagnosis of AD and its prodromal forms.<sup>1</sup> CSF biomarkers are now included in international guidelines for the diagnosis of AD in research settings and clinical practice<sup>2,3</sup> and the Alzheimer's Association appropriate use criteria for the use of lumbar puncture and CSF testing in the diagnosis of AD have been published.<sup>4</sup> Such biochemical diagnostics are currently implemented in many specialized centers around the world. Different methods of analysis have

been developed over the last decade and each laboratory has implemented the one best suited to its own practice. Related to this diversity there are also variations in pre-analytical and analytical conditions (such as sample tubes, storage, dilution of the biological sample, definition of cut-off values) between centers. The subsequent interpretation of the analytical results may depend on the calculation of ratios (such as t-tau/ $A\beta_{1-42}$  or  $A\beta_{1-42}/A\beta_{1-40}$ <sup>5-7</sup>), the use of scales (PLM,<sup>8</sup> Erlangen<sup>9</sup> scores), or on additional experiments (eg, dilution if t-tau is above the limit for detection<sup>10</sup>). Some laboratories mentioned the use of the A/T/N<sup>11</sup> classification, which is, however, based on data additional to CSF biomarkers, and is used more in the research setting than in the clinic. Depending on the laboratory, the type of report sent back to physicians (prescribing or referring physicians, and general practitioners) varies greatly, which may raise questions about the commutability of the tests and cause misunderstanding. It is therefore very important to harmonize comments on the reporting of results, so that the conclusions are similar regardless of where the analysis is performed.

Our work provides an overview of the procedures used in 40 centers worldwide performing CSF analysis to support AD diagnosis. For

each clinical laboratory participating in this work, we report the pre-analytical (eg, type of sample tubes, storage conditions, potential non-compliance with pre-specified local protocols) and analytical (quantified biomarkers, methods) conditions. We also detail each partner's post-analytical procedures, such as cutoff values, and use of ratios or scales for the interpretation of the results. Then, we list the clinical reports (for each biochemical profile) sent to the physicians responsible for the prescriptions. On the basis of the most frequently used reports and in-depth exchange and discussion between the participants, we propose harmonized reports adapted to each biochemical CSF profile.

This work is an essential step towards a consensual harmonization of clinical reporting after CSF analysis in the context of AD diagnosis, as advocated by the Biofluid Based Biomarkers Professional Interest Area (BBB-PIA) working group of the Alzheimer's Association. This harmonization is of great importance given the prevalence of AD and the increasing number of laboratories performing these diagnostic assays worldwide.

## 2 | METHODS

### 2.1 | Partners involved

Centers and laboratories specialized in AD diagnosis were contacted through the French Society of Clinical Biology (SFBC, <https://www.sfbc-asso.fr/>), the International Society to Advance Alzheimer's Research and Treatment (ISTAART) BBB-PIA, or the Society for Neurochemistry and Clinical CSF analysis (<http://www.neurochem.info/>). A total of 40 centers (17 French and 23 from 15 different countries, see authors' affiliations and Supplementary Figure S1 in the Supporting Information) provided different levels of information regarding their practice. For the interpretation of the surveys, each laboratory was anonymized. No personal or clinical patient data were used for this project, which therefore did not require ethical clearance. Data were collected between June and December 2020.

### 2.2 | Inquiries

Clinical laboratories performing CSF testing were asked to provide information on the pre-analytical and analytical protocols in their clinical practice (eg, type of tubes used, centrifugation or storage protocol, type of kit) and their criteria for non-conformity with local protocols. Post-analytical information was also requested, such as the cutoff values of the analytes, and the use of ratios or scales. All potential combinations of amyloid  $\beta$ , t-tau and p-tau(181) were then regrouped in eight different profiles labeled as follows: (1) "all normal," in which amyloid ( $A\beta_{1-42}$  or ratio  $A\beta_{1-42}/A\beta_{1-40}$ ), t-tau, and p-tau(181) are within reference range values; (2) "all pathological," in which all biomarkers show pathological values; (3) "amyloid," in which only amyloid values are pathological; (4) "t-tau," in which only t-tau values are pathological; (5) "amyloid & t-tau," in which amyloid and t-tau values are patholog-

### HIGHLIGHTS

- Diversity of global practices in cerebrospinal fluid biomarker quantification
- Definition of a common terminology for biochemical biomarkers' reporting
- Harmonization of clinical reporting for Alzheimer's disease diagnosis

### RESEARCH IN CONTEXT

1. **Systematic review:** The quantification of cerebrospinal fluid (CSF) biomarkers (beta-amyloid peptides, total tau, and phosphor-tau proteins) is gradually being implemented in specialized clinical laboratories as an aid in the diagnosis of Alzheimer's disease (AD). However, for the same biochemical profile, the interpretation and reporting of results may differ from one center to another, which can lead to misunderstandings and raises questions about the commutability of tests.
2. **Interpretation:** Consensus approach to provide the most accurate and informative comments regarding the interpretation of CSF biomarkers will benefit AD care, prevention, and treatment strategies.
3. **Future direction:** This work is an essential first step towards harmonization of the clinical reporting of CSF biomarkers panel for the diagnosis of AD. We also consider this work useful to new CSF biomarkers and from the perspective of defining reporting comments for emerging blood biomarkers of AD.

ical; (6) "amyloid & p-tau(181)," in which amyloid and p-tau(181) values are pathological; (7) "t-tau & p-tau(181)," in which values for both tau biomarkers are pathological; and (8) "p-tau(181)," in which only p-tau(181) values are pathological. These profiles were also associated with their corresponding values of the scales and ratios (see Supplementary Table S3). We asked the participants to provide the different reporting texts they used according to the most common biochemical profiles they encountered.

### 2.3 | Data processing and decision making

Clinical comments were compiled into a single table and returned to participants for selection. The percentage of similar reports and an initial vote to identify the two most relevant comments resulted in a short list of comments for each biochemical profile. This list was then commented on electronically and a series of video conferences was held to reach a consensus on the proposed comments for different profiles.

**TABLE 1** Pre-analytical and analytical features used by the different centers (in %) for the measurement of AD biomarkers in CSF

	Participant centers (in %)
Use of similar collection tube	94.0
Measurement on frozen sample	97.0
Automated immunoassay analyzer	88.2
Cutoff based solely on manufacturer's information	16.6
Dilution of t-tau if above limits	59.4
Systematic measurement of Ab <sub>1-40</sub>	58.1
Use of the Ab <sub>1-42</sub> /Ab <sub>1-40</sub> ratio	82.3
Use of other ratios than Ab <sub>1-42</sub> /Ab <sub>1-40</sub>	57.6
Use of scales	38.2
Turnover <= 1 week	34.5

The various comments were then translated into the different national languages of the participants. The laboratory of Prof. Sylvain Lehmann (France) in Montpellier was in charge of piloting the study and preparing, collecting, and analyzing all the survey responses.

## 2.4 | Role of the funding/sponsoring source

This research did not receive any specific grant from funding agencies in the public, commercial, or not-for-profit sectors. The study supporters had no role in the design and conduct of the study; collection, management, analysis, and interpretation of the data; preparation, review, or approval of the manuscript; or decision to submit the manuscript for publication.

## 3 | RESULTS

### 3.1 | Pre-analytical and analytical conditions overview

Table 1 summarizes the current practice across participating centers and Supplementary Table S1 provides detailed data on the analytical and pre-analytical procedures of each participating laboratory. The majority of centers (94%) used very similar polypropylene (PP) tubes for sample collection, and storage conditions at  $-20^{\circ}\text{C}$  for short term and  $-80^{\circ}\text{C}$  for long term. This is in line with standard operating procedures (SOPs) and the various experimental works in the field.<sup>12-15</sup> The use of collection tubes different from those recommended by each laboratory was identified as a non-conformity by 97% of centers. For secondary tubes that have a lower impact on pre-analytical variability,<sup>13</sup> we observed a greater diversity of origin, but these were PP microtubes from 0.5 to 2 mL in most cases.

For the analytical part, Fujirebio immunoassays (Lumipulse) were used in 76.5% of the centers for the quantification of the four analytes (A $\beta_{1-42}$ , A $\beta_{1-40}$ , t-tau, and p-tau[181]). Other laboratories measured the four analytes using Roche (Elecys), Euroimmun (ELISA), Fujirebio (ELISA), IBL (ELISA), or MSD (V-Plex), either alone or in combination. One center used a liquid chromatography mass spectrometry (LC-MS) approach for A $\beta_{1-42}$ . Overall, 88.2% used automated immunoassay analyzers. We observed an important heterogeneity of the cutoffs selected for the different analytes, not only for A $\beta_{1-42}$ , but also for t-tau and p-tau(181).

The selection of cutoffs by the centers was based on information given solely by manufacturers in 16.6% of cases, on literature (4.2%), on other laboratories/colleagues (12.5%), on internal data (45.8%), or on a combination of these approaches (25%). Cutoffs for A $\beta_{1-40}$  have generally not been defined, as this analyte is mainly used to calculate the A $\beta_{1-42}$ /A $\beta_{1-40}$  ratio, which has its own cutoff. A $\beta_{1-40}$  was systematically quantified in 58.1% of laboratories or added in case of discordance (ie, when A $\beta_{1-42}$  was normal but tau biomarkers were abnormal, or vice versa) in 22.6% of laboratories (Table 1 and Supplementary Table S1). Seventeen percent of the laboratories mentioned the use of a "grey zone," which corresponds to profiles where the values, often close to the cutoffs, correspond to a situation that remains undetermined.

Dilution of samples, when the upper limit of detection was exceeded, was only performed for t-tau and only in 59.4% of the laboratories. The A $\beta_{1-42}$ /p-tau(181) and t-tau/p-tau(181) ratios were computed in 39.4% and 30.3% of the laboratories, respectively. The validated PLM<sup>8</sup> and Erlangen<sup>9</sup> scales combining biomarkers are used by only 17.6% of the participating laboratories, a low percentage that is certainly an underestimate depending on the country, the Erlangen score being widely used in Germany for example. CSF biomarkers were also used by seven centers to establish the ATN<sup>11</sup> research classification that also includes imaging information.

### 3.2 | Clinical reports according to CSF biomarker profiles

Supplementary Table S3 (row 6) shows the mean and minimum/maximum frequency of the eight biochemical profiles, observed in consecutive series of samples (55 to 3000 samples) in 15 of the participating laboratories. Supplementary Table S2 lists the comments initially provided by participants for the biochemical profiles: the most frequent ones ("all normal," "all pathological," "amyloid," "t-tau") and the less frequent ("amyloid & t-tau or amyloid & p-tau[181]," "t-tau & p-tau[181]" and "p-tau[181]"). We observed very similar comments provided for "all normal," "all pathological," and "amyloid" profiles in 30%, 26%, and 36.6% of the centers, respectively. The comments for the other profiles showed more heterogeneity. In a second step, each laboratory was asked to select, from among all the reports previously listed, the two that best reflected their own current clinical practice. They were also given the opportunity to add additional comments based on available clinical information.

**TABLE 2** Summary of consensus comments for interpretation of biochemical profiles of AD biomarkers in CSF

amyloid	t-tau	p-tau(181)	Consensus comments
N	N	N	Biochemical profile not consistent with Alzheimer's disease.
P	P	P	Biochemical profile consistent with Alzheimer's disease.
P	N	N	Biochemical profile consistent with an amyloidopathy.
N	P	N	Biochemical profile not consistent with Alzheimer's disease; may be consistent with other neurodegenerative disease and/or neuronal damage. (If t-tau is close to/above upper limit of detection with a high t-tau/p-tau[181] ratio, the profile may indicate Creutzfeldt-Jakob disease)
P	P	N	Atypical biochemical profile; may be consistent with Alzheimer's disease.
P	N	P	Atypical biochemical profile; consistent with Alzheimer's disease.
N	P	P	Atypical biochemical profile; not consistent with Alzheimer's disease.
N	N	P	Atypical biochemical profile; not consistent with Alzheimer's disease.

Note to be added to all comments: This biochemical profile must be interpreted in its clinical context and in conjunction with a physician.

Abbreviations: N, normal; P, pathological; p-tau(181), tau phosphorylated at threonine 181; t-tau, total tau.

On the basis of these proposals, and after several rounds of exchanges (electronically and by videoconference) with all the partners, we generated harmonized comments for each profile, associated in some cases with additional information (Table 2 and Supplementary Table S3). We also translated these comments into 11 languages corresponding to the different countries of the participating laboratories (Supplementary Table S4). It should be noted that our survey found that less than 5% of centers use plots/graphs in addition to numerical values. In our group discussion, the consensus was not to use additional plots/graphs that might interfere or make the commentary be returned with the numeric values less clear.

## 4 | DISCUSSION

In this work, we collected information from 40 centers located in 15 different countries (Supplementary Figure S1) that measure CSF amyloid and tau biomarkers in the clinical setting. Moving from the measurement of biomarkers for clinical research, which is mainly performed on retrospective cohorts, to routine clinical measurement is a real challenge.<sup>4</sup> Even with established SOPs<sup>12</sup> and international guidelines for the handling of CSF,<sup>15</sup> pre-analytical and analytical deviations may be present in real world settings, affecting the result, and some-

times going unreported. To provide high-quality results, it is important to ensure that tests achieve sufficient levels of performance to make a meaningful contribution to diagnosis and ultimately to patient care. Finally, a critical step in the medical use of CSF testing is how the results are communicated to clinicians and, when appropriate, to patients themselves. The harmonization of reporting is in this regard essential to avoid misunderstandings while comparing results between centers and to provide accurate, informative, and harmonized information that will have an impact on prevention, care and treatment strategies.<sup>1,16</sup>

We have focused our work only on CSF biomarkers currently used in clinical practice, which are part of official guidelines<sup>2,3</sup> and measured using IVD (In Vitro Diagnostics) assays. The main context of use (COU) of these tests is the diagnosis of AD. It is particularly important to keep this COU in mind because it influences the choice of pathological cutoffs (which may vary depending on the clinical question). This also explains why comments in our proposed consensus clinical reports focus on the diagnosis of AD rather than dementia or neurodegenerative diseases in general.

The methodology to be used in the development of clinical practice guidelines is well established.<sup>17</sup> The first phase generally involves conducting a systematic review and synthesis of the literature. Different works focus on the interpretation of biochemical profiles,<sup>8,9,15</sup> but we could not find previous publications dealing with the clinical reporting of CSF results for AD diagnosis. This observation is not surprising since these tests have only recently been widely used in clinical routine. Therefore, we employed a "consensus" methodological approach directly based on the agreement among experts through iterative ratings with feedback.

Our review of the pre-analytical protocols of the different centers firstly shows that study over the last ten years of the confounding factors related to this phase<sup>12</sup> and the definition and harmonization of SOPs<sup>14,15</sup> for CSF AD biomarkers have been successfully implemented. Indeed, most of the centers use very similar PP tubes (only differing in their size and shape), which have previously shown low adsorption with amyloid  $\beta$  peptides.<sup>18</sup> However, the volume of stored samples differs among centers, and this may still affect amyloid  $\beta$  quantification. Pre-analytical procedures, including centrifugation, secondary aliquots in microtubes and freezing at  $-20^{\circ}\text{C}$  for a few days or at  $-80^{\circ}\text{C}$  were also very similar, with the exception of the secondary tubes. Rather than using fresh samples, use of frozen secondary samples may represent a more easy-to-use protocol,<sup>19</sup> which probably adapts to current numbers of tests requested.

Regarding the analytical part, in addition to the three core biomarkers  $A\beta_{1-42}$ , t-tau and p-tau(181), we observed that more than 83% of the laboratories also measure  $A\beta_{1-40}$  and thereafter compute the  $A\beta_{1-42}/A\beta_{1-40}$  ratio. The rationale is likely related to the fact that the ratio improves diagnostic performance<sup>6,20</sup> and reduces biases linked to collection tube, volume of sample, and storage. In terms of detection method, it is also unsurprising that more than 88% of the laboratories are using automated chemiluminescence immunoassays that have a reduced analytical variability<sup>21,22</sup> and offer a more flexible test throughput. Despite this, one striking finding was the wide dispersion

of the cutoff values used by the different laboratories, even for  $A\beta_{1-42}$ , which will benefit from a metrological harmonization as a result of the development of both mass spectrometry reference methods and certified reference materials.<sup>23</sup> This diversity is linked on the one hand to the different assays/protocols used, but also to the fact that most centers adapt the values proposed by the test providers using results from their own cohort (16.6% vs 45.8%, Supplementary Table S1). In addition, there is a highly variable distribution of the percentages of CSF profiles between centers (Supplementary Table S3). This is consistent with the prevalence of AD, which varies widely between centers, from a low of 25% to a high of 53%. This indirectly leads to different optimal threshold definitions; therefore, the positive and negative predictive values of the tests also vary considerably.

Regarding the interpretation and reporting of the CSF biomarker results, the inquiries from the different centers (Supplementary Table S2) showed first of all that the different comments involve "Alzheimer's disease," which is in line with the COU of the CSF biomarkers. We also note the term "biochemical profile." It should be kept in mind that if CSF biomarkers do mainly reflect the amyloid and tau pathologies of AD, they may show pathologic change without clinical symptoms and this may occur sometimes more than a decade before clinical manifestation of AD.<sup>1</sup> Nevertheless, in accordance with the National Institute on Aging and Alzheimer's Association Research Framework, AD can be defined as a biological construct,<sup>24</sup> which led us in the comments to refer to AD per se, rather than with its pathological changes.

There are several points to consider in the derivation of the consensus comments from the initial reporting of the biomarker profiles. First, when all biomarkers are pathological, most initial comments from the centers indicated that the profile is "suggestive" or "consistent" with AD, showing some caution in asserting the diagnosis. The performance of CSF biomarkers for AD is very good,<sup>1</sup> but there may still be room for improvement using other p-tau biomarkers such as p-tau(217)<sup>25</sup> or p-tau(231).<sup>26</sup> In addition, the physician's diagnosis of AD remains multidisciplinary, combining clinical history, symptomatology, neuropsychological testing, imaging, and biology,<sup>1</sup> and accordingly CSF biomarkers alone are not diagnostic of the disease. The consensus comment in this pathological situation was therefore "Biochemical profile consistent with Alzheimer's disease," and we have chosen to mention in all cases that "This biochemical profile must be interpreted in its clinical context and in conjunction with a physician" (Table 2).

Second, when all biomarkers are normal, one may clearly indicate that the profile is "not consistent" with AD. This makes sense because AD is intrinsically associated with amyloid and tau pathology and retrospective studies show that normal CSF profiles virtually rule out AD.<sup>27</sup>

Third, when only amyloid biomarkers are pathological there is an obvious consensus to indicate that the biochemical profile is consistent with an "amyloidopathy" (Table 2 and Supplementary Table S3). However, this pathological situation may also be considered indicative of the presence of AD in the disease continuum.<sup>24</sup>

Fourth, when  $A\beta_{1-40}$  is assessed, there is usually no defined cutoff value for this analyte alone. Variations of  $A\beta_{1-40}$  in frontotemporal dementia<sup>28</sup> or cerebral amyloid angiopathy<sup>29</sup> have been described but

they are minimal and therefore only useful when associated with clinical information in favor of these diagnoses. However, assessment of  $A\beta_{1-40}$  in combination with  $A\beta_{1-42}$  ( $A\beta_{1-42}/A\beta_{1-40}$  ratio) has proven to be highly informative for AD diagnosis in clinical routine<sup>6,7,20</sup> and this ratio is currently used in 88.2% of the centers.

Fifth, the profile showing an isolated increase in t-tau is present in 2.7% to 15.7% of cases, depending on the center. Initial comments indicated that this profile was not consistent with AD but rather with other neurodegeneration and/or neuronal damage (such as cerebrovascular disease or—if strongly increased—Creutzfeldt-Jakob disease [CJD]). For the latter possibility, it should be noted that a very high level of t-tau (close to or above the high detection limit of the assays) associated with a high t-tau/p-tau(181) ratio is strongly in favor of this diagnosis, if other causes of major neuronal injury, for example, stroke and encephalitis, are excluded.<sup>10</sup> The consensus comment for this profile could therefore refer to this particular situation if clinical information or the diagnostic hypothesis suggests the presence of CJD (see Table 2).

Sixth, to cover all situations, we also defined consensus comments for the less frequent profiles (< 5% of cases). Thus, when amyloid biomarkers are pathological and associated with an increase in p-tau(181) but not t-tau, one may consider that this atypical profile "is consistent" with AD. When both amyloid and t-tau biomarkers are pathological (with p-tau[181] normal), the consensus is to consider that this atypical profile "may be consistent" with AD. On the other hand, profiles with normal amyloid but abnormal t-tau and/or p-tau(181) are considered as "not consistent" with AD (Table 2). This is an important consensus decision emphasizing that AD can exist only in the presence of amyloid pathology. This situation is also reminiscent of the "suspected non-Alzheimer's disease pathophysiology (SNAP)."<sup>30</sup> It should be noted that in these cases, special attention should be paid to the search for amyloidopathy using in particular the  $A\beta_{1-42}/A\beta_{1-40}$  ratio.

A limitation of this work is that we are not exhaustive in consulting laboratories using CSF biomarkers. Therefore, some results such as the percentage of use of different ratios or scales may be biased. In addition, CSF biomarkers are not always used for the diagnosis of AD and, therefore, many centers and countries are missing from this international study.

In conclusion, this work is an essential first step towards harmonization of the clinical reporting of the CSF biomarkers panel for the diagnosis of AD. The proposed framework is adaptable and applicable to new CSF biomarkers passing regulatory criteria and prospective validations for clinical application. We also consider this work useful from the perspective of defining reporting comments for emerging blood biomarkers of AD.

## ACKNOWLEDGEMENTS

This manuscript was facilitated by the Alzheimer's Association International Society to Advance Alzheimer's Research and Treatment (ISTAART), through the Biofluid Based Biomarkers Professional Interest Area (BBB-PIA). The views and opinions expressed in this publication represent those of the authors and do not necessarily reflect those of the BBB-PIA membership, ISTAART, or the Alzheimer's Association.



The authors thank the French Society of Clinical Biology, the ISTAART BBB-PIA, and the Society for Neurochemistry and Clinical CSF analysis for their help in initiating this work.

Constance Delaby received no support for the present manuscript.

Charlotte Teunissen received the following grants during the last 36 months: Research of CET is supported by the European Commission (Marie Curie International Training Network, grant agreement No 860197 [MIRIADE], and JPND), Health Holland, the Dutch Research Council (ZonMw), Alzheimer Drug Discovery Foundation, The Selfridges Group Foundation, Alzheimer Netherlands, Alzheimer Association. CT and WF are recipients of ABOARD, which is a public-private partnership receiving funding from ZonMw (#73305095007) and Health~Holland, Topsector Life Sciences & Health (PPP-allowance; #LSHM20106). More than 30 partners participate in ABOARD. ABOARD also receives funding from Edwin Bouw Fonds and Gieskes-Strijbisfonds. IV is appointed on a research grant by Alzheimer Nederland (NL-17004). She received no support for the present manuscript.

Kaj Blennow is supported by the Swedish Alzheimer Foundation (#AF-742881) and Hjärnfonden, Sweden (#FO2017-0243). He received no support for the present manuscript.

Daniel Alcolea received funding from Institute of Health Carlos III (ISCIII), Spain PI18/00435 and INT19/00016, and by the Department of Health Generalitat de Catalunya PERIS program SLT006/17/125. He received no support for the present manuscript.

Ivan Arisi was partly supported by: Fondo Ordinario Enti (FOE D.M 865/2019) funds in the framework of a collaboration agreement between the Italian National Research Council and EBRI (2019-2021); POR (Operative Program Lazio Region, Italy) FESR (European Program Regional Development) 2014-2020, Public Notice "LIFE 2020" MODIAG Project.

Elodie Bouaziz-Amar received no support for the present manuscript.

Anne Beaume received no support for the present manuscript.

Aurélié Bedel received no support for the present manuscript.

Giovanni Bellomo is supported by University of Perugia, CIRMMF, IRST Istituto Romagnolo per lo Studio e la Cura dei Tumori, Innovatech srl. He received no support for the present manuscript.

Edith Bigot-Corbel received no support for the present manuscript.

Maria Bjerke received no support for the present manuscript.

Marie-Céline Blanc-Quintin received no support for the present manuscript.

Mercè Boada received support from LA CAIXA, IMI, ISCIII H2020 the European Union/EFPIA Innovative Medicines Initiative Joint Undertaking MOPEAD project (Grants No. 115985) and is also supported by national grant, PI17/01474 from Acción Estratégica en Salud, integrated into the Spanish National Plan of R+D+I and founded by ISCIII (Instituto de Salud Carlos III)-Subdirección General de Evaluación and the Fondo Europeo de Desarrollo Regional (FEDER- "Una manera de Hacer Europa"). She received no support for the present manuscript.

Olivier Bousigues is supported by A2MCL Alsace Alzheimer. He received no support for the present manuscript.

Miles D. Chapman received no support for the present manuscript.

Mari L. DeMarco is supported by the Michael Smith Foundation for Health Research, Brain Canada (Canada Brain Research Fund), Health Canada, Women's Brain Health Initiative, Alzheimer Society of Canada, St. Paul's Foundation, Djavad Mowafaghian Centre for Brain Health at the University of British Columbia and the Canadian Consortium for Neurodegeneration and Aging; with all funds provided to the University of British Columbia.

Mara D'Onofrio was partly supported by Fondo Ordinario Enti (FOE D.M 865/2019) funds in the framework of a collaboration agreement between the Italian National Research Council POR (Operative Program Lazio Region, Italy) FESR (European Program Regional Development) 2014-2020, Public Notice "LIFE 2020", MODIAG Project. She received no support for the present manuscript.

Julien Dumurgier received no support for the present manuscript.

Diane Dufour received no support for the present manuscript.

Sebastiaan Engelborghs is supported for various projects: Research Project GSKE/FMRE, Research Project FWO Vlaanderen, PhD fellowship FWO Vlaanderen (n = 3), VLAIO PhD fellowship. He received no support for the present manuscript.

Hermann Esselmann received no support for the present manuscript.

Anne Fogli received no support for the present manuscript.

Audrey Gabelle received no support for the present manuscript.

Elisabetta Galloni received no support for the present manuscript.

Clémentine Gondolf received no support for the present manuscript.

Frédérique Grandhomme received no support for the present manuscript.

Oriol Grau-Rivera receives grants from the Spanish Ministry of Science, Innovation and Universities (FJCI-2017-33437), and from the Alzheimer's Association Research Fellowship Program (2019-AARF-644568). He received no support for the present manuscript.

Melanie Hart receives UCLH Biomedical Research Centre support from NIHR (Neurosciences). She received no support for the present manuscript.

Takeshi Ikeuchi received the following grants to his institution AMED: 21dk0207049, 21dk0207045, 21ek0109545, 20ek0109350, 20ek0109392. He received no support for the present manuscript.

Andreas Jeromin received no support for the present manuscript.

Kensaku Kasuga received no support for the present manuscript.

Ashvini Keshavan is supported by the Weston Brain Institute/Selfridges Foundation grant UB170045. She received no support for the present manuscript.

Michael Khalil received unrestricted research grants from Biogen and Novartis. He received no support for the present manuscript.

Peter Körtvelyessy received no support for the present manuscript.

Agnieszka Kulczynska-Przybik received no support for the present manuscript.

Jean-Louis Laplanche received no support for the present manuscript.

Qiao-Xin Li received no support for the present manuscript.

Alberto Lleó is supported by grants from Generalitat de Catalunya (PERIS SLT002/16/00408), Instituto de Salud Carlos III (PI17/01896), Instituto de Salud Carlos III CIBERNED. He received no support for the present manuscript.

Catherine Malaplate received no support for the present manuscript.

Marta Marquí is supported by the Instituto de Salud Carlos III (ISCIII) Acción Estratégica en Salud, integrated in the Spanish National RCDCI Plan and financed by ISCIII-Subdirección General de Evaluación and the Fondo Europeo de Desarrollo Regional (FEDER-Una manera de hacer Europa) grant PI19/00335. She received no support for the present manuscript.

Colin M. Masters received no support for the present manuscript.

Barbara Mroczo received a grant from The Binding Site Group and Biokom Diagnostyka. She received no support for the present manuscript.

Léonor Nogueira received no support for the present manuscript.

Adelina Orellana received no support for the present manuscript.

Markus Otto is supported by BMBF, Thierry Latran foundation, EU, DFG. He received no support for the present manuscript.

Jean-Baptiste Oudart is supported by Astrazeneca grant for a research program on EGFR targeted therapy resistance in NSCLC. He received no support for the present manuscript.

Claire Paquet received no support for the present manuscript.

Federico Paolini received no support for the present manuscript.

Lucilla Parnetti received no support for the present manuscript.

Armand Perret-Liaudet received no support for the present manuscript.

Katell Peoc'h is supported by contract with Siemens Healthineers. She received no support for the present manuscript.

Koen Poesen received no support for the present manuscript.

Albert Puig-Pijoan received no support for the present manuscript.

Isabelle Quadrio received no support for the present manuscript.

Murielle Quillard-Muraine received no support for the present manuscript.

Benoît Rucheton received no support for the present manuscript.

Susanne Schraen received no support for the present manuscript.

Jonathan M. Schott is supported by the Medical Research Council, Alzheimer's Research UK, Weston Brain Institute, British Heart Foundation, Alzheimer's Association. He received no support for the present manuscript.

Leslie M. Shaw is supported by NIA for ADNI Biomarker Core, UPENN ADRC Biomarker Corr, MJ Fox PPMI for CSF Biomarker measurements, Roche for IIS study of AD biomarkers. He received no support for the present manuscript.

Marc Suárez-Calvet receives funding from the European Research Council (ERC) under the European Union's Horizon 2020 research and innovation program (Grant agreement No. 948677). He also receives funding from the Instituto de Salud Carlos III (PI19/00155) and from the Spanish Ministry of Science, Innovation and Universities (Juan de la Cierva Programme grant IJC2018-037478-I). He received no support for the present manuscript.

Magda Tsolaki received no support for the present manuscript.

Hayrettin Tumani is supported by Alexion, Bayer, Biogen, Celgene, Genzyme-Sanofi, Merck, Novartis, Roche, Teva. He received no support for the present manuscript.

Chinedu T Udeh-Momoh is supported by Alzheimer research UK (Project Grant). He received no support for the present manuscript.

Lucie Vaudran received no support for the present manuscript.

Marcel M. Verbeek is supported by the BIONIC project (no. 733050822, which has been made possible by ZonMw within the framework of "Memorabel," the research and innovation program for dementia, as part of the Dutch national "Deltaplan for Dementia": zonmw.nl/dementiaresearch), and the CAFÉ project (the National Institutes of Health, USA, grant number 5R01NS104147-02). The BIONIC project is a consortium of Radboudumc, LUMC, ADX Neurosciences, and University of Rhode Island University. MM Verbeek is also supported by a grant from the Selfridges Group Foundation. He received no support for the present manuscript.

Federico Verde received no support for the present manuscript.

Lisa Vermunt is supported by Alzheimer Nederland. She received no support for the present manuscript.

Jonathan Volgesgang is supported by Clinician Scientist Research Fellowship, German Research Foundation (Deutsche Forschungsgesellschaft, project number: 413501650; 09/01/2019-02/28/2021); Eric Dorris Memorial Fellowship (McLean Hospital; 07/01/2020-06/30/2021). He received no support for the present manuscript.

Jens Wiltfang is supported by Federal Ministry of Education and Research (BMBF) State of Lower Saxony; He received no support for the present manuscript.

Henrik Zetterberg is a Wallenberg Scholar supported by grants from the Swedish Research Council (#2018-02532), the European Research Council (#681712), Swedish State Support for Clinical Research (#ALFGBG-720931), the Alzheimer Drug Discovery Foundation (ADDF), USA (#201809-2016862), the AD Strategic Fund and the Alzheimer's Association (#ADSF-21-831376-C, #ADSF-21-831381-C and #ADSF-21-831377-C), the Olav Thon Foundation, the Erling-Persson Family Foundation, Stiftelsen för Gamla Tjänarinnor, Hjärnfonden, Sweden (#FO2019-0228), the European Union's Horizon 2020 research and innovation program under the Marie Skłodowska-Curie grant agreement No 860197 (MIRIADE), and the UK Dementia Research Institute at UCL. He received no support for the present manuscript.

Sylvain Lehmann received no support for the present manuscript.

## CONFLICTS OF INTERESTS

Charlotte Teunissen's institution received consulting fees from Roche and received Kits for blood-based biomarkers from Quanterix and ADX Neurosciences. She leads the Global Biomarker Standardisation Consortium, Blood based Biomarker PIA, CSF Society.

Kaj Blennow has served as a consultant (and received fees), at advisory boards, or at data monitoring committees for Abcam, Axon, Biogen, JOMDD/Shimadzu, Julius Clinical, Lilly, MagQu, Novartis, Roche Diagnostics, and Siemens Healthineers, and is a co-founder of Brain Biomarker Solutions in Gothenburg AB (BBS), which is a part of the GU Ventures Incubator Program, all outside the present work.

Daniel Alcoléa participated in advisory boards from Fujirebio-Europe and Roche Diagnostics and received speaker honoraria from Fujirebio-Europe, Roche Diagnostics, Nutricia, Krka Farmacéutica S.L., and Esteve Pharmaceuticals S.A. Dr Alcoléa declares a filed patent application (WO2019175379 A1 Markers of synaptopathy in neurodegenerative disease).

Elodie Bouaziz-Amar received fees from Roche Pharma/Roche Diagnostics France (Board Alzheimer 23 octobre 2019).

Giovanni Bellomo received fees from Innuvatech srl.

Mercè Boada received consulting fees from Roche, Grifols, Araclon, Biogen, Merck, Lilly, Nutricia, Cotexyme, Renew Research.

Olivier Bousigues received consulting fees from KeyQuest Health Ltd, Azure Knowledge Corporation, and benefited from manuscript writing fees from Spectra diagnostic and Revue francophone des laboratoires.

Miles D. Chapman is a member of UK NEQAS SAG Committee.

Sebastian Engelborghs received consulting fees from Biogen, Danone, Eisai, icometrix, Pfizer, Novartis, Nutricia, Roche, and received research funding from ADx Neurosciences and Janssen Pharmaceutica. He received fees as an advisory Boards from Biogen, Danone, Eisai, icometrix, Pfizer, Novartis, Nutricia, Roche. He serves as Secretary-General of Belgian Neurological Society, Vice-President of Belgian Dementia Council, ExCo member of European Alzheimer Disease Consortium.

Melanie Hart is member of the Royal College of Pathologists Special Advisory Board for Immunology—clinical scientist representative.

Takeshi Ikeuchi received a honoraria for lectures from the following pharmaceutical companies: Eisai, Takeda, Novartis, Roche, Daichi-Sankyo, Ono, Chugai.

Andreas Jeromin, has served as an advisor to Quanterix Corp and holds stock options.

Michael Khalil has received speaker honoraria from Bayer, Novartis, Merck, Biogen Idec, and Teva Pharmaceutical Industries Ltd. He serves on scientific advisory boards for Biogen Idec, Merck Serono, Roche, Novartis, Bristol-Myers Squibb, and Gilead.

Agnieszka Kulczynska-Przybik received a consulting and/or lecture honoraria from Roche company.

Piotr Lewczuk received consulting and/or lecture honoraria from IBL International, Fujirebio Europe, AJ Roboscreen, and Roche.

Alberto Lleó has served as a consultant or at advisory boards and received fees from Roche, Biogen, Nutricia, Zambon, MassConsultoria, SEMNIM. In addition, Dr. Lleó has a patent WO2019175379 A1 Markers of synaptopathy in neurodegenerative disease issued. He received royalties from a European Patent licensed.

Barbara Mroczko has received consulting and/or lecture honoraria from Abbott, Wiener, Roche, Cormay and Biameditek.

Léonor Nogueira is a co-author of a patent concerning the detection of antibodies toward citrullinated protein for the diagnosis of Rheumatoid Arthritis.

Adelina Orellana has submitted MMP-10 CSF biomarker in the European Patent Office (Patent number 21382305 7, priority date April 9th 2021)

Markus Otto received fees as consultant or advisory board from Axon, Roche, Axon, Biogen. He is a member of the CSF Society.

Jean-Baptiste Oudart received fees as an expert advisory board for Osimertinib (Astrazeneca).

Claire Paquet received fees as a consultant from Pharmaceutical laboratories Biogen and Roche concerning anti-AD treatments, as a lecturer for Roche, as advisory board member from Biogen and Roche.

Armand Perret-Liaudet received consulting fees from QUADRANT consulting (Paris), support from Fujirebio company for attending meetings.

Katell Peoc'h received consulting fees from Vifor France.

Albert Puig-Pijoan received support from Nutricia (payment of inscription fees to attend meetings).

Jonathan M. Schott received royalties from Oxford University Press, Henry Stewart Talks, Alzheimer's Research and Therapy, consulting fees from Eli Lilly, Biogen, GE, Roche, Alzheimer's Research UK, UK Dementia Research Institute Ltd and as advisory board from Axon Neuroscience SE. He serves as Chair ISTAART Advisory Council.

Leslie M. Shaw received fees from Biogen teaching program.

Marc Suárez-Calvet received fees as a consultant and at advisory boards for Roche Diagnostics International Ltd, as a lecturer in symposia sponsored by Roche Diagnostics, S.L.U and Roche Farma, S.A. and as member of advisory boards for Roche Diagnostics International Ltd.

Magda Tsolaki is member of Alzheimer Hellas Greek Alzheimer's Disease Federation.

Hayrettin Tumani received fees or honoraria for lectures and advisory boards from Alexion, Bayer, Biogen, Celgene, Genzyme-Sanofi, Merck, Novartis, Roche, Teva. He is member of DGLN (German Society of CSF Diagnostics and Clinical Neurochemistry).

Chinedu T Udeh-Momoh received support (travel grant) from Alzheimer's Association and British society for neuroendocrinology. He is trustee for British society for neuroendocrinology.

Lisa Vermunt received support (travel grant) from the Alzheimer Association.

Jens Wiltfang received fees as a consultant from Roche Pharma GmbH, Boehringer Ingelheim, Immungenetics AG, MSD Sharp & Dohm, Abbott AG, Biogen, as a lecturer from Pfizer, Janssen-Cilag GmbH, med Update GmbH, AGNP e. V., CSF Society, Roche Pharma, Vitos Klinikum Kurhessen, AWO Psychiatrie Akademie, Actelion Pharmaceuticals, Amgen, Fachverband Rheumatologische Fachass., Neuroakademie e. V., Guangzhou Glorylen Medical Technology Co. (China), Beijing Yibai Science and Technology Ltd. He has Patents (no.: EP1270592B1; patent no.: US 6,849,416 B2; patent no.: EP2095128B1; EP3105589 A1). He is member of AGNP.

Henrik Zetterberg is a co-founder of Brain Biomarker Solutions in Gothenburg AB (BBS), which is a part of the GU Ventures Incubator Program, which holds licenses (unrelated to the current submission). He served at scientific advisory boards for Eisai, Denali, Roche Diagnostics, Wave, Samumed, Siemens Healthineers, Pinteon Therapeutics, Nervgen, AZTherapies, and CogRx. He has given lectures in symposia sponsored by Cellectricon, Fujirebio, Alzecure, and Biogen. He is chair of the Alzheimer's Association Global Biomarker Standardization

Consortium and the Alzheimer's Association Biofluid-Based Biomarker Professional Interest Area.

Sylvain Lehmann received consulting fees from Fujirebio, Euroimmun, and Roche diagnostics.

All other authors declare that they have no competing interests and nothing to disclose.

## AUTHOR CONTRIBUTIONS

S. Lehmann and C. Delaby had full access to all of the data in the study and take responsibility for the integrity of the data and the accuracy of the data analysis.

Concept and design: S. Lehmann, C. Delaby, C. E. Teunissen, H. Zetterberg.

Acquisition, analysis, or interpretation of data: All authors.

Drafting of the manuscript: S. Lehmann and C. Delaby.

Critical revision of the manuscript for important intellectual content: All authors.

## ORCID

Constance Delaby  <https://orcid.org/0000-0002-8606-6814>

## REFERENCES

- Scheltens P, De Strooper B, Kivipelto M, et al. Alzheimer's disease. *Lancet*. 2021;397:1577-1590. S0140673620322054. [https://doi.org/10.1016/S0140-6736\(20\)32205-4](https://doi.org/10.1016/S0140-6736(20)32205-4).
- Dubois B, Feldman HH, Jacova C, et al. Advancing research diagnostic criteria for Alzheimer's disease: the IWG-2 criteria. *Lancet Neurol*. 2014;13:614-629. [https://doi.org/10.1016/S1474-4422\(14\)70090-0](https://doi.org/10.1016/S1474-4422(14)70090-0).
- McKhann GM, Knopman DS, Chertkow H, et al. The diagnosis of dementia due to Alzheimer's disease: recommendations from the National Institute on Aging-Alzheimer's Association workgroups on diagnostic guidelines for Alzheimer's disease. *Alzheimers Dement*. 2011;7:263-269. <https://doi.org/10.1016/j.jalz.2011.03.005>.
- Shaw LM, Arias J, Blennow K, et al. Appropriate use criteria for lumbar puncture and cerebrospinal fluid testing in the diagnosis of Alzheimer's disease. *Alzheimers Dement*. 2018;14:1505-1521. <https://doi.org/10.1016/j.jalz.2018.07.220>.
- Lehmann S, Dumurgier J, Ayrignac X, et al. Cerebrospinal fluid A beta 1-40 peptides increase in Alzheimer's disease and are highly correlated with phospho-tau in control individuals. *Alzheimers Res Ther*. 2020;12:123. <https://doi.org/10.1186/s13195-020-00696-1>.
- Hansson O, Lehmann S, Otto M, Zetterberg H, Lewczuk P. Advantages and disadvantages of the use of the CSF Amyloid  $\beta$  (A $\beta$ ) 42/40 ratio in the diagnosis of Alzheimer's Disease. *Alzheimers Res Ther*. 2019;11:34. <https://doi.org/10.1186/s13195-019-0485-0>.
- Lehmann S, Gabelle A, Paquet C. Can we rely only on ratios of cerebrospinal fluid biomarkers for AD biological diagnosis? *Alzheimers Dement*. 2015;11:1125-1126. <https://doi.org/10.1016/j.jalz.2014.09.003>.
- Lehmann S, Dumurgier J, Schraen S, et al. A diagnostic scale for Alzheimer's disease based on cerebrospinal fluid biomarker profiles. *Alzheimers Res Ther*. 2014;6:38. <https://doi.org/10.1186/alzrt267>.
- Lewczuk P, Zimmermann R, Wiltfang J, Kornhuber J. Neurochemical dementia diagnostics: a simple algorithm for interpretation of the CSF biomarkers. *J Neural Transm*. 2009;116:1163-1167. <https://doi.org/10.1007/s00702-009-0277-y>.
- Lehmann S, Paquet C, Malaplate-Armand C, et al. Diagnosis associated with Tau higher than 1200 pg/mL: insights from the clinical and laboratory practice. *Clin Chim Acta*. 2019;495:451-456. <https://doi.org/10.1016/j.cca.2019.04.081>.
- Jack CR Jr, Bennett DA, Blennow K, et al. A/T/N: an unbiased descriptive classification scheme for Alzheimer disease biomarkers. *Neurology*. 2016;87:539-547. <https://doi.org/10.1212/WNL.0000000000002923>.
- del Campo M, Mollenhauer B, Bertolotto A, et al. Recommendations to standardize preanalytical confounding factors in Alzheimer's and Parkinson's disease cerebrospinal fluid biomarkers: an update. *Biomark Med*. 2012;6:419-430. <https://doi.org/10.2217/bmm.12.46>.
- Hansson O, Mikulskis A, Fagan AM, et al. The impact of preanalytical variables on measuring cerebrospinal fluid biomarkers for Alzheimer's disease diagnosis: a review. *Alzheimers Dement*. 2018;14:1313-1333. <https://doi.org/10.1016/j.jalz.2018.05.008>.
- Fourier A, Portelius E, Zetterberg H, Blennow K, Quadrio I, Perret-Liaudet A. Pre-analytical and analytical factors influencing Alzheimer's disease cerebrospinal fluid biomarker variability. *Clin Chim Acta*. 2015;449:9-15. <https://doi.org/10.1016/j.cca.2015.05.024>.
- Hansson O, Batrla R, Brix B, et al. The Alzheimer's Association international guidelines for handling of cerebrospinal fluid for routine clinical measurements of amyloid  $\beta$  and tau. *Alzheimers Dement*. 2021;17:1575-1582.
- Molinuevo JL, Blennow K, Dubois B, et al. The clinical use of cerebrospinal fluid biomarker testing for Alzheimer's disease diagnosis: a consensus paper from the Alzheimer's Biomarkers Standardization Initiative. *Alzheimers Dement*. 2014;10:808-817. <https://doi.org/10.1016/j.jalz.2014.03.003>.
- Wang X, Chen Y, Yang N, et al. Methodology and reporting quality of reporting guidelines: systematic review. *BMC Med Res Methodol*. 2015;15:74. <https://doi.org/10.1186/s12874-015-0069-z>.
- Perret-Liaudet A, Pelpel M, Tholance Y, et al. Risk of Alzheimer's disease biological misdiagnosis linked to cerebrospinal collection tubes. *J Alzheimers Dis*. 2012;31:13-20. <https://doi.org/10.3233/JAD-2012-120361>.
- Hansson O, Rutz S, Zetterberg H, et al. Pre-analytical protocol for measuring Alzheimer's disease biomarkers in fresh CSF. *Alzheimers Dement (Amst)*. 2020;12:e12137. <https://doi.org/10.1002/dad2.12137>.
- Lehmann S, Delaby C, Boursier G, et al. Relevance of A $\beta$ 42/40 ratio for detection of Alzheimer disease pathology in clinical routine: the PLM<sub>R</sub> scale. *Front Aging Neurosci*. 2018;10:138. <https://doi.org/10.3389/fnagi.2018.00138>.
- Bittner T, Zetterberg H, Teunissen CE, et al. Technical performance of a novel, fully automated electrochemiluminescence immunoassay for the quantitation of  $\beta$ -amyloid (1-42) in human cerebrospinal fluid. *Alzheimers Dement*. 2016;12:517-526. <https://doi.org/10.1016/j.jalz.2015.09.009>.
- Agnello L, Piccoli T, Vidali M, et al. Diagnostic accuracy of cerebrospinal fluid biomarkers measured by chemiluminescent enzyme immunoassay for Alzheimer disease diagnosis. *Scand J Clin Lab Invest*. 2020;80:313-317. <https://doi.org/10.1080/00365513.2020.1740939>.
- Andreasson U, Kuhlmann J, Pannee J, et al. Commutability of the certified reference materials for the standardization of  $\beta$ -amyloid 1-42 assay in human cerebrospinal fluid: lessons for tau and  $\beta$ -amyloid 1-40 measurements. *Clin Chem Lab Med*. 2018;56:2058-2066. <https://doi.org/10.1515/cclm-2018-0147>.
- Jack CR Jr, Bennett DA, Blennow K, et al. NIA-AA Research Framework: Toward a biological definition of Alzheimer's disease. *Alzheimers Dement*. 2018;14:535-562. <https://doi.org/10.1016/j.jalz.2018.02.018>.
- Barthélemy NR, Bateman RJ, Hirtz C, et al. Cerebrospinal fluid phospho-tau T217 outperforms T181 as a biomarker for the differential diagnosis of Alzheimer's disease and PET amyloid-positive patient identification. *Alzheimers Res Ther*. 2020;12:26. <https://doi.org/10.1186/s13195-020-00596-4>.

26. Suárez-Calvet M, Karikari TK, Ashton NJ, et al. Novel tau biomarkers phosphorylated at T181, T217 or T231 rise in the initial stages of the preclinical Alzheimer's continuum when only subtle changes in A $\beta$  pathology are detected. *EMBO Mol Med.* 2020;12:e12921. <https://doi.org/10.15252/emmm.202012921>.
27. Vercausse O, Paquet C, Gabelle A, et al. Relevance of follow-up in patients with core clinical criteria for Alzheimer disease and normal CSF biomarkers. *Curr Alzheimer Res.* 2018;15:691-700. <https://doi.org/10.2174/1567205015666180110113238>.
28. Gabelle A, Roche S, Gény C, et al. Decreased sA $\beta$ PP $\beta$ , A $\beta$ 38, and A $\beta$ 40 cerebrospinal fluid levels in frontotemporal dementia. *J Alzheimers Dis.* 2011;26:553-563. <https://doi.org/10.3233/JAD-2011-110515>.
29. Charidimou A, Friedrich JO, Greenberg SM, Viswanathan A. Core cerebrospinal fluid biomarker profile in cerebral amyloid angiopathy: a meta-analysis. *Neurology.* 2018;90:e754-e762. <https://doi.org/10.1212/WNL.0000000000005030>.
30. Jack CR Jr, Knopman DS, Chételat G, et al. Suspected non-Alzheimer disease pathophysiology—concept and controversy. *Nat Rev Neurol.* 2016;12:117-124. <https://doi.org/10.1038/nrneuro.2015.251>.

#### SUPPORTING INFORMATION

Additional supporting information may be found in the online version of the article at the publisher's website.

**How to cite this article:** Delaby C, Teunissen CE, Blennow K, et al. Clinical reporting following the quantification of cerebrospinal fluid biomarkers in Alzheimer's disease: An international overview. *Alzheimer's Dement.* 2021;1-12. <https://doi.org/10.1002/alz.12545>