



**HAL**  
open science

# The structure of hemicalide from the marine sponge *Hemimycale* sp.

Laurence Marcourt, Georges Massiot

► **To cite this version:**

Laurence Marcourt, Georges Massiot. The structure of hemicalide from the marine sponge *Hemimycale* sp.. *European Journal of Organic Chemistry*, 2022, pp.e202200208. 10.1002/ejoc.202200208 . hal-03635190

**HAL Id: hal-03635190**

**<https://hal.univ-reims.fr/hal-03635190>**

Submitted on 27 Feb 2024

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

**Content:**

Procedure for isolation of hemicalide

**Figure S1:** HPLC profile and UV of hemicalide

**Figure S2:**  $^1\text{H}$  NMR spectrum of hemicalide in  $\text{CD}_3\text{OD}$  at 500 MHz

**Figure S3:** COSY NMR spectrum of hemicalide in  $\text{CD}_3\text{OD}$

**Figure S4:** HOHAHA NMR spectrum of hemicalide in  $\text{CD}_3\text{OD}$

**Figure S5:**  $^{13}\text{C}$ -DEPTQ NMR spectrum of hemicalide in  $\text{CD}_3\text{OD}$  at 126 MHz

**Figure S6:** Edited-HSQC NMR spectrum of hemicalide in  $\text{CD}_3\text{OD}$

**Figure S7:** HMBC NMR spectrum of hemicalide in  $\text{CD}_3\text{OD}$

**Isolation of hemicalide**

The fresh sponge (5 kg) was lyophilized to give 650 g of a wax, which was macerated for 6 hours with chloroform and then twice with a 9/1 mixture of methanol and water. The filtrates were pooled and concentrated to an aqueous syrup, which was partitioned with dichloromethane and ethyl acetate. After evaporation of the organic solutions (42 g), the extract was solubilized in a 9/1 methanol/water mixture and partitioned with hexane. The "defatted" methanolic extract was chromatographed on a column of Sigel with mixtures of ethyl acetate and methanol of increasing polarities. All these steps were monitored by a bio-assay and by LC/MS. The activity was located in the more polar fractions, which were pooled and subjected first to a LH20 exclusion chromatography, then to RP18 HPLC with a gradient of water in acetonitrile. It was thus obtained 0.5 mg of hemicalide. HRMS:  $m/z$  1061.6780, calc. for  $[\text{M-H}]^-$  1061.6782 in the negative ESI mode, corresponding to  $\text{C}_{59}\text{H}_{98}\text{O}_{16}$  (calc. 1061.6782). UV (DAD detection) 238, 306 nm.





Figure S1: HPLC profile and UV of hemicalide



**Figure S2:** <sup>1</sup>H NMR spectrum of hemicalide in CD<sub>3</sub>OD at 500 MHz











Figure S5:  $^{13}\text{C}$ -DEPTQ NMR spectrum of hemicalide in  $\text{CD}_3\text{OD}$  at 126 MHz



**Figure S6:** Edited-HSQC NMR spectrum of hemicalide in CD<sub>3</sub>OD







Figure S7: HMBC NMR spectrum of hemicalide in CD<sub>3</sub>OD



