



HAL
open science

Bis-iridoids and other constituents from *Scabiosa semipapposa*

Samia Bendamene, Naima Boutaghane, Charlotte Sayagh, Abdulmagid Alabdul Magid, Zahia Kabouche, Chawki Bensouici, Laurence Voutquenne-Nazabadioko

► **To cite this version:**

Samia Bendamene, Naima Boutaghane, Charlotte Sayagh, Abdulmagid Alabdul Magid, Zahia Kabouche, et al.. Bis-iridoids and other constituents from *Scabiosa semipapposa*. *Phytochemistry Letters*, 2022, 49, pp.202-210. 10.1016/j.phytol.2022.04.005 . hal-03653763

HAL Id: hal-03653763

<https://hal.univ-reims.fr/hal-03653763>

Submitted on 28 Apr 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Bis-iridoids and other constituents from *Scabiosa semipapposa*

Samia Bendamene^{a,b,c}, Naima Boutaghane^a, Charlotte Sayagh^b, Abdulmagid Alabdul Magid^b,
Zahia Kabouche^a, Chawki Bensouici^c, Laurence Voutquenne-Nazabadioko^{b,*}

^a*Université des frères Mentouri-Constantine 1, Département de chimie, Laboratoire d'Obtention des Substances Thérapeutiques (LOST), Campus Chaabet-Ersas, 25000 Constantine, Algérie.*

^b*Université de Reims Champagne Ardenne, CNRS, ICMR UMR 7312, 51097 Reims, France*

^c*Centre de Recherche en Biotechnologie, Ali Mendjli Nouvelle Ville UV 03, Constantine, Algérie.*

Abstract

Six undescribed bis-iridoids named Semipapposiridoids A–F (**1-6**) together with seventeen known compounds were obtained from the roots and aerial parts of *Scabiosa semipapposa* Salzm. ex DC.. Among the known compounds, three lignans were isolated for the first time in *Scabiosa* species, whereas the other known constituents including iridoids, phenol glycosides and flavonoids were previously isolated from *Scabiosa* sp. Their structures were assigned on the basis of extensive 1D- and 2D-NMR experiments (^1H , ^{13}C , COSY, TOCSY, NOESY, HSQC, HMBC), mass spectrometry HR-ESI-MS and by comparison of their spectral data with those of literature values. To the best of our knowledge, this should be the first report on the bis-iridoids content of this plant; furthermore, the fact that the plant is rich in mono- and bis-iridoids will support chemotaxonomy of the genus. The total phenolic and flavonoid contents of the 80% MeOH extracts and fractions were evaluated, with their antioxidant activities using three different methods, namely DPPH, ABTS and CUPRAC. Most of the fractions showed a moderate to good antioxidant activity compared to the standards BHA and BHT.

Keywords:

Scabiosa semipapposa Salzm. ex DC., Caprifoliaceae, Bis-iridoid glucosides, Semipapposiridoids A-F, Antioxidant activity.

1. Introduction

The Caprifoliaceae, also known as the honeysuckle family, is a dicotyledonous flowering plant consisting of about 860 species in approximately 42 genera, with a nearly cosmopolitan distribution, with the centers of diversity in eastern North America and eastern Asia, while absent in tropical and southern Africa (Xuand Chang, 2017). Plants of this family are usually shrubs and vines and rarely herbaceous. Many genera of this family like *Cephalaria* (Tabatadze et al., 2007; Pasi et al., 2009; Mustafayeva et al., 2010; Sarikahya and Kirmizigul, 2010; Sarikahya et al., 2021), *Valeriana* (Ming et al., 1997; Tang et al., 2002; Dong et al., 2015; Quan et al., 2020; Liu et al., 2021), *Lonicera* (Lin et al., 2008; Liu et al., 2015; Li et al., 2019; Qiu et al., 2021), and *Scabiosa* (Lehbili et al., 2018a; Lehbili et al., 2018b; Bendamene et al., 2020; Kılınç et al., 2020) have attracted the attention of various authors for their phytochemical and biological properties. The *Scabiosa* genus, belonging to this family, is widespread in the Mediterranean area, including 12 species in Algeria (Quezel and Santa, 1963). Many *Scabiosa* species are used in folk medicine for the treatment of various illnesses such as asthma, influenza, bronchitis, bronchial pneumonia, liver diseases, neurodegenerative diseases, and to treat certain dermatoses, in particular scabies (Girre, 1980; Bammi and Douira, 2002; Rigat et al., 2007; Kose et al., 2015; Moteetee et al., 2016; Pinto et al., 2018). Numerous *Scabiosa* species yielded a great variety of secondary metabolites such as triterpene saponins (Alimbaeva et al., 1977; Baykal et al., 1998; Zheng et al., 2004; Lehbili et al., 2018a; Kılınç et al., 2020; Bendamene et al., 2020), flavonoids, coumarins (Garaev et al., 2008; Al-Qudah et al., 2017; Lehbili et al., 2018b), mono- and bis-iridoid glucosides (Papalexandrou et al., 2003; Polat et al., 2010; Lehbili et al., 2018b) and phenolic compounds (Akar, 2022). In our continuous researches on the chemical constituents of *Scabiosa* species growing in Algeria (Lehbili et al., 2018a; Lehbili et al., 2018b; Bendamene et al., 2020), we have investigated *Scabiosa semipapposa* Salzm. Ex DC. roots and aerial parts. After previous isolation of triterpenoid

saponins from the roots of this plant (Bendamene et al., 2020), new phytochemical investigations were performed for its iridoids content. In this paper, we thus describe the isolation and structure elucidation of six undescribed bis-iridoids (**1–6**), namely semipapposiridoids A-F (Fig. 1), along with seventeen (**7–23**) known compounds. Chemotaxonomy of this species was discussing. In addition, the antioxidant capacity of the 80% methanol extracts and fractions were evaluated using DPPH, ABTS and CUPRAC methods.

2. Results and discussion

The 80% methanol extract of *S. semipapposa* roots was fractionated by vacuum-liquid chromatography (VLC) on RP-C₁₈ to give five fractions (R_I to R_V) which were purified by successive chromatographic techniques, including flash chromatography, as well as semi-preparative and preparative high performance liquid chromatography (HPLC) to yield eleven known compounds (**9–19**). In the same manner, the 80% methanol extract of the aerial parts afford six undescribed compounds (**1–6**) (Fig.1) and six known ones (**7, 8, 20–23**). Their structures were established by 1D- and 2D-NMR analysis (¹H, ¹³C, COSY, TOCSY, NOESY, HSQC, HMBC) in combination with mass spectrometry ESI- and HR-ESI-MS, and by comparison of their physical and spectral data with those of literature values. The known compounds were identified as sylvestroside II (**7**) (Jensen et al., 1979), sylvestroside I (**8**) (Jensen et al., 1979; Abdallah,1991), triplotoside A (**9**) (Graikou et al.,2002; Gülcemal et al.,2010), sweroside (**10**) (Jensen et al., 1979), loganin (**11**) (Calis et al., 1984a), secoxyloganin (**12**) (Calis and sticher. 1984b), vanilloloside (**13**) (El-Ghazooly et al., 2003), vanillyl- β -D-glucopyranoside (**14**) (Kanho et al., 2005), caffeic acid methylester (**15**)(Rani et Devanan.,2013), 3,4,5-trimethoxyphenyl-1-*O*- β -apiofuranosyl-(1" \rightarrow 6')- β -glucopyranoside (= kelampayosideA) (**16**) (Kanchanapoom et al., 2002; Qingwei et al., 2012), (+)-8-hydroxypinoresinol (**17**) (Tsukamoto et al.,1984a;Tsukamoto et al.,1984b; Cowan et al.,2001),

(+)-8-hydroxypinoresinol-4'-*O*- β -D-glucopyranoside) (**18**) (Deyama et al., 1986), (+)-8-hydroxypinoresinol-4-*O*- β -D-glucopyranoside) (**19**) (Tsukamoto et al., 1984a; Tsukamoto et al., 1985), apigenin (**20**) (Wawer et al., 2001), luteolin (**21**) (Okamura et al., 1994), lutéoline-7-*O*- β -D-glucopyranoside (**22**) (Chiruvella et al., 2007), and kaempferol 3-*O*-(3'',6''-di-*O*-*E*-*p*-coumaroyl)- β -D-glucopyranoside (**23**) (Yang et al., 2010).

Compound **1** was obtained as a yellow amorphous powder, with a molecular formula of C₄₄H₅₆O₂₂ based on HR-ESI-MS (*m/z* 959.3170 [M+Na]⁺, calcd for 959.3161). The infrared (IR) spectrum indicated the presence of the hydroxyl functions (3356 cm⁻¹), and α,β unsaturated ester carbonyl (1695 and 1629 cm⁻¹). The ¹H and ¹³C NMR spectra (Table 1) in combination with the HSQC and ¹H-¹H COSY correlations showed two distinct parts, indicated as units A and B (Fig. 1). The signals corresponding to unit A included one olefinic methine [δ_{H} 7.52, s (H-3a); δ_{C} 153.9 (C-3a)], one vinyl [δ_{H} 5.80, ddd (*J*=17.4, 10.4, 8.6 Hz, H-8a); δ_{C} 135.7 (C-8a), δ_{H} 5.28, dd (*J*=10.4, 1.5 Hz, H-10a), 5.33, d(*J*=17.4 Hz, H-10a); δ_{C} 119.6 (C-10a)], one methylene oxy [δ_{H} 4.11, 4.14, (H-7a); δ_{C} 64.1 (C-7a)], one methylene [δ_{H} 1.86, td (*J*=13.8, 7.4 Hz, H-6a), 2.12, dd (*J*=13.8, 7.2 Hz, H-6a); δ_{C} 30.1(C-6a)], two methines [δ_{H} 2.90, q (*J*=6.3 Hz, H-5a); δ_{C} 31.3 (C-5a), δ_{H} 2.70, dt (*J*=8.3, 6.1 Hz, H-9a); δ_{C} 45.4 (C-9a)], one acetal [δ_{H} 5.58, d (*J*=6.6 Hz, H-1a); δ_{C} 97.7 (C-1a)], one carboxy group [δ_{C} 168.2 (C-11a)] and one carbomethoxy group [δ_{C} 173.0 (C-12a)].

This ¹H and ¹³C NMR spectroscopic data of **1** showed that unit A contained a secoiridoid moiety, almost identical to secologanic acid (Kocsis et al., 1993), except that compound **1** had a carbomethoxy group at C-7a [δ_{C} 64.1 (C-7a)] instead of the aldehyde group of secologanic acid, which was also detected from the HMBC correlations from H-5a and H-6a to C-7a (Fig. 2).

The remaining resonances in the ¹H and ¹³C NMR spectra of **1** attributed to unit B, particularly those arising from an olefinic methine [δ_{H} 7.47, s (H-3b); δ_{C} 152.5 (C-3b)], one methylene

[$\delta_{\text{H}}1.78$, ddd ($J=14.1, 7.8, 5.1$ Hz), 2.33, ddd ($J=14.1, 7.7, 1.5$ Hz)(H-6b); $\delta_{\text{C}}40.3$ (C-6b)], three methines [$\delta_{\text{H}}3.15$, q($J=7.9$ Hz,H-5b); $\delta_{\text{C}}32.6$ (C-5b), $\delta_{\text{H}}2.18$, m (H-8b); $\delta_{\text{C}}41.0$ (C-8b); $\delta_{\text{H}}2.13$, ddd ($J=17.4, 8.8, 4.8$ Hz,H-9b); $\delta_{\text{C}}47.1$ (C-9b)], one oxygenated methine [$\delta_{\text{H}}5.24$, t ($J=4.5$ Hz, H-7b); $\delta_{\text{C}}78.4$ (C-7b)], one acetal [$\delta_{\text{H}}5.33$, d ($J=4.6$ Hz, H-1b); $\delta_{\text{C}}97.6$ (C-1b)], one secondary methyl [$\delta_{\text{H}}1.10$, d ($J=6.8$ Hz, H-10b); $\delta_{\text{C}}13.8$ (C-10b)], one methoxy [$\delta_{\text{H}}3.71$, s (MeO-12b); $\delta_{\text{C}}51.7$ (MeO-12b)] and one carboxy group [$\delta_{\text{C}}169.3$ (C-11b)], indicated the second unit to be loganic-type iridoid (Kocsis et al., 1993). The major differences between them is the downfield shifts of H-7b ($\delta_{\text{H}}5.24$) and C-7b ($\delta_{\text{C}}78.4$). The attachment site between unit A and B was found to be ester linkage between the C-7 of unit B and the carboxyl group (C-11a) of unit A as deduced by the HMBC correlation observed between the oxygenated methine proton (H-7b) and the carboxy carbon (C-11a).

Furthermore, two anomeric protons resonances corresponding to *O*-linked sugars were observed in the ^1H NMR spectrum of **1** as two doublets at $\delta_{\text{H}}4.74$ ($J = 7.9$ Hz) and 4.76 ($J = 7.9$ Hz). The sugar units were identified after analysis of COSY and HSQC spectra as two glucoses [glc-A ($\delta_{\text{H}-1'}$ 4.74 and $\delta_{\text{C}-1'}$ 100.1) and glc-B ($\delta_{\text{H}-1''}$ 4.76 and $\delta_{\text{C}-1''}$ 100.2)] (Table 1). The β -anomeric configurations were defined by the $^3J_{\text{H}1,\text{H}2}$ coupling constants and the comparison of ^{13}C -NMR chemical shifts with those in the literature (Agrawal, 1992) (Table 1). The glc-A unit was linked to C-1a as deduced from the long-range correlation observed between H-1' and C-1a in the HMBC spectrum, and the glc-B unit was linked to C-1b as deduced from the long-range correlation observed between H-1'' and C-1b in the HMBC spectrum (Fig. 2). These ^1H and ^{13}C NMR spectroscopic data of **1** are very similar to sylvestroside II (Soeren et al., 1979), except for the signals for an additional *trans-p*-coumaroyl unit in **1** with two coupled *trans* double-bond protons [$\delta_{\text{H}}6.39$, H-8''' and 7.69 , H-7''', each d, $J = 15.9$ Hz] and four aromatic protons at $\delta_{\text{H}}7.49$ and 6.83 (each 2H, d, $J = 8.6$ Hz) assignable to H-2'''/6''' and H-3'''/5''' respectively, as well as to an ester carbonyl ($\delta_{\text{C}}168.5$) in the ^{13}C NMR spectrum. The chemical

shifts of C-4'' in the glucopyranosyl moiety (glc-B) were downfield shifted from δ_C 71.6 to δ_C 72.4 relative to those of sylvestroside II (Soeren et al., 1979). Additionally, the location of the *E-p*-coumaroyl unit was established by the key cross-peaks between the H-4'' (δ_H 4.87) of glucosyl and carbonyl carbon C-9''' (δ_C 168.5) of the coumaroyl in the HMBC spectrum. The relative configuration of **1** was further determined by the NOESY experiment. The NOE correlations observed between δ_H 2.90 (H-5a) with 2.70 (H-9a), 2.13 (H-9b) with 1.10 (H-10b), 3.15 (H-5b) with 2.13 (H-9b) and 5.24 (H-7b) with 2.18 (H-8b) reveal that they were β -oriented, as is usually observed in the iridoid skeleton (Jensen et al., 1979; Kocsis et al., 1993; Tomassini et al., 2000; Dinda et al., 2006; Ji et al., 2012). Based on the above findings, compound **1** was identified as 4''-*O-E-p*-coumaroyl-sylvestroside II, named semipapposiridoid A shown in Fig. 1 and differs from sylvestroside II, reported in *Dipsacus sylvestris* (Soeren et al., 1979), by an additional *E-p*-coumaroyl group at C-4''.

Semipapposiridoid B (**2**) was isolated as a yellow powder. The HR-ESI-MS, m/z 959.3169 [M+Na]⁺ was identical to that obtained for compound **1**, suggesting that both compounds were isomers with molecular formula C₄₄H₅₆O₂₂. The IR spectrum indicated the presence of the hydroxyl functions (3352cm⁻¹) and α,β unsaturated ester carbonyl (1688 and 1630 cm⁻¹), which are responsible for the maximum UV absorption of the compound at 231 nm (Tian et al., 2006). Comparison of ¹H and ¹³C NMR values and the analysis of the ¹H-¹H-COSY, NOESY, HSQC and HMBC revealed that compound **2** is structurally closely related to **1** and contains the same skeleton (sylvestroside II) (Table 1). The only significant difference is the location of the *E-p*-coumaroyl group. The *E-p*-coumaroyl group was linked to the C-6'' of glc-B unit in **2** instead of the C-4'' of glc-B in **1**. The HMBC spectrum gave further evidence of this by showing long-range correlations between the C-6'' methylene protons at δ_H 4.45 and 4.54 of glucosyl and the carbonyl carbon C-9''' at δ_C 168.9. The relative configuration of **2** was established as identical

to **1** by the NOESY data of **2**. Hence, the structure of **2** was elucidated as 6''-*O-E-p*-coumaroyl-sylvestroside II (Fig. 1).

Semipapposiridoid C (**3**) was isolated as a mixture with semipapposiridoid A (**1**) in the ratio 1:1(*Z:E*), based on the integral intensity of the corresponding signals in their ¹H NMR. Both components of the mixture had the same molecular formula C₄₄H₅₆O₂₂ [HR-ESI-MS, *m/z* 959.3169 [M+ Na]⁺]. The ¹H and ¹³C NMR spectra (Table 1) of **3** were extremely close to those of compound **1**. The only significant difference was the value of the vinyl proton coupling constants [δ_{H} 5.83 and 6.94 (each d, *J* = 12.8 Hz)]. HMBC correlations of the ester carbonyl at δ_{C} 167.3 to the *Z*-vinyl protons [δ_{H} 5.83 and 6.94 (each d, *J* = 12.8 Hz)] and Glc-H-4'' [4.86 (t, *J* = 9.5 Hz)] indicated that the structure of semipapposiridoid C (**3**) was 4''-*O-Z-p*-coumaroyl-sylvestroside II (Fig. 1).

The ¹H-NMR and ¹³C-NMR spectra (Table 1) of semipapposiridoid D (**4**) were very similar to those of semipapposiridoid B (**2**) and again showed that they were a mixture of sylvestroside II esterified at the C-6'' position and revealed the presence of *Z*- and *E-p*-coumaroyl groups. They had the same molecular formula, C₄₄H₅₆O₂₂, according to their HR-ESI-MS. The *Z-p*-coumaroyl was located at the C-6'' position in **4**, based on HMBC correlations from the ester carbonyl at δ_{C} 170.8 to the *Z*-vinyl protons [δ_{H} 5.80 and 6.90 (each d, *J* = 12.9 Hz)] and Glc-H-6'' [4.37 (dd, *J* = 12.0, 6.5 Hz) and 4.50 (dd, *J* = 12.0, 2.1 Hz)]. The relative configuration of **4** was established as identical to **2** by the NOESY data of **4**. Therefore, the chemical structure of semipapposiridoid D was determined to be 6''-*O-Z-p*-coumaroyl-sylvestroside II (Fig. 1).

Semipapposiridoid E (**5**) was obtained as a yellowish amorphous powder. Its molecular formula C₂₆H₄₀O₁₂ was obtained from its HR-ESI-MS (*m/z* 567.2413; [M + Na]⁺, calcd 587.3298). The IR spectrum of **5** showed absorption bands of the hydroxy group at 3377cm⁻¹ and the carbonyl groups at 1692 and 1629cm⁻¹, respectively. The UV spectrum showed absorption maxima at λ 235 nm. The ¹H- and ¹³C-NMR spectroscopic signals of **5** were almost identical to those of

abelioside B (Murai et al., 1985) except for the presence of signals at $\delta_{\text{H}}3.42$ (3H, s, 3-OCH₃), $\delta_{\text{C}}56.0$ attributed to the methoxy group, instead of the signal for an ester carbonyl group in abelioside B. In the HMBC spectrum, a correlation between the protons at $\delta_{\text{H}}3.42$ and carbons at $\delta_{\text{C}}102.9$ (C-3) were observed, indicating that this methoxy group was located at C-3. The relative configuration of **5** was established by analysis of the NOESY spectrum showing correlations between H-5 and H-9 β axial and between H-1 β axial and CH₃-10 which indicated that H-5, H-9 and CH₃-10 were all β -oriented, and between H-7 and H-8 α -oriented. The NOESY correlations observed between H-3 and H-1 β axial indicated a β -axial orientation of H-3 and an α -axial orientation of methoxyl group at the C-3 position. Based on the above evidence, the structure of semipapposiridoid E (**5**) was established as depicted in Fig. 1.

Semipapposiridoid F (**6**) was isolated as a yellowish amorphous powder. It has the same molecular formula as compound **5** (C₂₆H₄₀O₁₂) and differs from the latter only by the orientation of the methoxy group at C-3. The IR spectrum displayed absorption bands at 3376, 1693 and 1629 cm⁻¹ attributed to the hydroxyl and carbonyl functions. The ¹³C NMR spectra (see Table 2) of **6** showed great similarities to those of **5**, except for the signals of C-1 $\delta_{\text{C}}59.3$ (Δ -4.9 ppm), C-3 $\delta_{\text{C}}99.3$ (Δ -3.6 ppm) and C-4 $\delta_{\text{C}}33.1$ (Δ -2.9 ppm). The lack of correlation between H-3 and H-1 in the NOESY spectrum indicates the β -axial orientation of the methoxyl group at C-3 (Breitmaier and Voelter, 1987). Therefore, the structure of **6** (semipapposiridoid F) was elucidated as shown in Fig. 1.

The total phenolic and flavonoid contents of the 80% MeOH extracts of the roots (R) and aerial parts (A) of *S. semipapposa* and their fractions R_I to R_V (for the roots), A_I to A_V (for the aerial parts), obtained after VLC fractionation on RP-C₁₈, were quantitatively estimated using a calibration curve established in terms of gallic acid (GAE) and quercetin equivalents (QE), respectively (Table 3). Moderate phenolic and flavonoid contents were found in the 80% MeOH extracts with values of 109.0 μg GAE/mg and 12.8 μg QE/mg for the roots and 169.6 μg

GAE/mg and 19.5 μg QE/mg for the aerial parts. The highest contents of total phenolics and flavonoids were found in the fraction R_I with values of 361.7 μg GAE/mg and 74.6 μg QE/mg, while the fraction R_V containing saponins (Bendamene et al., 2020) was the poorest (32.3 μg GAE/mg and 5.7 μg QE/mg). Previous research carried out on organic extracts of the species *S. tschiliensis* (Wang et al. 2013), *S. arenaria* (Hlila et al. 2015) and *S. stelatta* (Mouffouk et al. 2018), indicated lower levels of phenol and flavonoid contents compared to the results of the present study. Here, we identified in the roots (fractions R_I-R_{IV}), three phenyl glucosides (13-14, 16), caffeic acid methyl ester (15), three lignans (17-19) and four iridoids (9-12), and in the aerial parts (A_{III} to A_V), eight bis-iridoids (1-8) and four flavonoids (20-23), thus justifying these results.

Several *in vitro* test procedures were carried out to estimate the antioxidant activity of crude plant extracts, but there is no simple universal method to assess antioxidant capacity accurately and quantitatively. Therefore, in the present study, three methods (DPPH, ABTS and CUPRAC assays), based on different principles, were applied to determine the antioxidant capacity of samples (Table 3). The antioxidant activity of the 80% MeOH extracts and fractions varied in a dose-dependent manner in all tests. The antioxidant activity of *S. semipapposa* samples ranged from 4.9 to 315.8 $\mu\text{g}/\text{mL}$ when measured by the DPPH assay, from 8.9 to 124.0 $\mu\text{g}/\text{mL}$ when measured by the ABTS^{•+} assay, and from 10.2 to 378.5 $\mu\text{g}/\text{mL}$ when measured by the CUPRAC assay. These results are in agreement with those reported by Mouffouk et al. (2018). The total antioxidant capacity values from ABTS^{•+} were higher than those of the DPPH[•] and CUPRAC assays. This may result from the presence of antioxidants that interact more selectively with fast reacting ABTS^{•+} radicals. Many studies reported that the solubility of the tested extract to different test systems and the stereoselectivity of radicals affect the ability of extracts to react against different radicals (Yu et al. 2002). The moderate antioxidant activity of extract and fractions of the aerial parts is correlated with the presence of amounts of iridoids, which in their

molecule have no (or few) phenolic OH groups, which neutralize free radicals. This is consistent with the study of [Pacifico et al. \(2009\)](#) and [Wu et al. \(2009\)](#) who reported that sweroside (**10**) and iridoids did not exhibit good radical scavenging ability.

Among the fractions tested, the fraction R_I showed significant antioxidant activity in all tests. Moreover, the results also indicated that this fraction showed higher activity (FRS₅₀ 4.9 µg/mL ([Table 3](#)) than those of the antioxidant standards BHA and BHT (FRS₅₀ 8.7, 15.9 µg/mL, respectively) in the DPPH assays. The good antioxidant activity of the fraction R_I could be associated with its high content of phenolic compounds (361.7±1.2 µg GAE/mg). Indeed, phenolic compounds have been the most studied for their antioxidant activity and have been reported to protect against many diseases through their ability to neutralize free radicals. ([Jayaprakasha et al., 2001](#); [Gali and Bdjou, 2019](#)). In the fraction R_I, the iridoids sweroside (**10**) and secologanine (**12**) were purified as well as the phenolic glucosides (**13**, **14**, **16**). Vanilloloside (**13**) was found to be a moderately active scavenger of DPPH FRS₅₀ 43.3 µg/mL and moderate CUPRAC activity (0.397 mM TRg⁻¹) ([Sarikahya et al., 2011](#)), while vanillyl β-D-glucopyranoside (**14**), showed DPPH scavenger activity (FRS₅₀ 30.88 µg/mL) ([Shi et al., 2012](#)).

3. Conclusions

In summary, the systematic phytochemical study on the roots and the aerial parts of *S. semipapposa* afforded six undescribed components, semipapposiridoids A-F (**1–6**) ([Fig. 1](#)), and three known bis-iridoid glucosides (**7–9**), three iridoids (**10–12**), three phenolic glucosides (**13–14**, **16**), one caffeic acid methyl ester (**15**), three lignans (**17–19**) and four flavonoids (**20–23**) ([Fig. S49](#) see supplementary data). These results expand our knowledge of chemical diversity and provide evidence to further analysis of their potential chemotaxonomic significance.

Compounds **7–12**, **20–23**, have been previously reported from different species of the genus *Scabiosa* ([Horn et al., 2001](#); [Papalexandrou et al., 2003](#); [Christopoulou et al., 2008](#); [Wang](#)

etal.,2015; Lehbili et al.,2018a) and might reveal the close chemotaxonomic relationships between *S. semipapposa* and other *Scabiosa* species. It is also interesting to note that lignans (17–19) are described for the first time in the genus *Scabiosa*. They could serve as potential chemotaxonomic markers to differentiate *Scabiosa* from other genera in the Caprifoliaceae family.

Among the bis-iridoids, compounds (1–4, 7–9) isolated from *S. semipapposa* were found to possess secoiridoid/iridoid-subtype skeletons consisting of secologanic acid condensed to the 7-OH of loganin or loganin-like iridoids. These results also showed the close relationship between *Scabiosa* species (Papalexandrou et al.,2003; Lehbili et al.,2018a).

In addition, the antioxidant activities of the 80% MeOH extracts and the fractions (R_I to R_V and A_I to A_{IV}) were evaluated by DPPH, ABTS and CUPRAC assays. Our results demonstrate that the fraction R_I has particularly high antioxidant and free radical scavenging activities, probably due to the presence of vanillyl alcohol glucosides.

4. Experimental

4.1. General experimental procedures

UV spectra were measured on a Shimadzu UV/Vis U-2450 spectrophotometer. Optical rotations of pure compounds were measured in CH₃OH using a MCP 5100 AntonPaar Polarimeter. IR spectra were recorded on a Nicolet Avatar 320 FT-IR spectrometer with KBr disks. The 1D and 2D NMR spectra (¹H and ¹³C NMR, ¹H-¹H COSY, NOESY, HSQC and HMBC) were performed using a Bruker Avance III 600 spectrometer (¹H at 600 MHz and ¹³C at 150 MHz) equipped with a 5 mm TCI cryoprobe. 2D-NMR experiments were performed using standard Bruker microprograms (TopSpin 3.5 software). ESI-MS and HR-ESI-MS experiments were performed using a Micromass Q-TOF micro instrument (Manchester, UK). Flash chromatography (FC) was carried out on a Grace Reveleris system equipped with dual UV and ELSD detection using Grace® cartridges (Silica gel or RP-C₁₈). The medium-pressure

liquid chromatography (MPLC) was employed using a Buchi pump system AP250/500 (Buchi, France), with a RP-C₁₈ silica gel MERCK column (15 × 230 and 26 × 460 mm). Semi-preparative HPLC was realized on a Dionex apparatus equipped with an ASI-100 automated sample injector, a STH 585 column oven, a P580 pump, a diode array detector UVD 340S and the Chromeleon[®] software version 6.8. A prepacked RP-C₁₈ column (Phenomenex 250 x 10 mm, Luna 5 μ) was used for semi-preparative HPLC. The eluting mobile phase consisted of H₂O with TFA (0.0025%) and CH₃CN with a flow rate of 5 mL/min and the chromatogram was monitored at 205 and 215 nm. Analytical HPLC experiments were performed using a Thermofisher Ultimate 3000 (Thermo Fischer Scientific, Villebon sur Yvette, France), equipped with a 4 ways pump LPG 3400 SD, an automatic injector WPS 3000 SL, a UV/visible diode array detector 3000 and the Chromeleon[®] software version 6.8. A prepacked C₁₈ column Uptisphere Strategy C₁₈ (Interchim, 4.6×250 mm, 5μ) was used for analytical HPLC and the mobile phase consisted of H₂O with TFA (0.0025% v/v) and CH₃CN. A gradient elution method was applied from 5% to 80% of CH₃CN in 30 min with a flow rate of 1 mL/min and the chromatograms were monitored at 205, 254, 300 and 360 nm. TLC was performed on pre-coated silicagel 60 F₂₅₄ Merck and compounds were visualized by spraying the dried plates with 50% H₂SO₄, followed by heating.

The antioxidant activity assays were carried out at the Center of Biotechnology Research (Contantine, Algeria) on a 96-well microplate reader, Perkin Elmer Multimode Plate Reader EnSpire (Perkin Elmer, France). Butylated hydroxyanisole (BHA), butylated hydroxytoluene (BHT), 1,1-diphenyl-2-picrylhydrazyl (DPPH[•]), 2,2'-azinobis (3-ethyl-benzothiazoline-6-suphonic acid) diammonium salt (ABTS^{•+}) and Ciocalteureagent (FCR) were obtained from Sigma Chemical Co. (Sigma-Aldrich GmbH, Sternheim, Germany). All other chemicals and solvents were of analytical grade.

4.2. Plant material

The whole plant *Scabiosa semipapposa* Salzm. ex DC. was collected in May 2017 from Alguemas, Constantine, North-Eastern Algerian (latitude 36.3479 and longitude 6.650773). The plant material was identified by Mr. Kamel Kabouche. A voucher specimen (LOST Ss.05/17) has been deposited at the herbarium of LOST Laboratory, University Frères Mentouri-Constantine, Algeria.

4.3. Extraction and isolation

The aerial parts and the roots of *S. semipapposa* were studied apart from each other. The dried and powdered roots of *S. semipapposa* (1 kg) were macerated in 80% MeOH (3 × 10 L, 24 h) at room temperature, followed by rotary evaporation at 45 °C under reduced pressure to give the crude extract (100 g). A part of this (55 g) was subjected to RP-C₁₈ vacuum liquid chromatography (VLC) eluted successively with MeOH-H₂O (3:7, 4:6, 6:4, 8:2 and 10:0), to provide five major fractions (R_I to R_V, respectively). Fraction R_I (1.3 g) was subjected to High Performance Flash Chromatography (HPFC), over silica gel, eluted by a gradient system of CH₂Cl₂-MeOH-H₂O (10:0:0 to 7:3:0.5), in 39 min to afford 69 sub-fractions (R_{I-1} to R_{I-69}). Sub-fractions R_{I-29} to R_{I-34} were combined (27.5 mg) and purified by semi-preparative HPLC using a gradient (10-30% CH₃CN, in 20 min) to furnish compounds **14** (1.4 mg, *t_R* 5.9 min), **10** (3.5 mg, *t_R* 11.1 min). Then the combined sub-fractions R_{I-36-37} (23 mg) were purified by semi-preparative HPLC (10-30% CH₃CN, in 20 min) to afford compounds **13** (1 mg, *t_R* 4.5 min), **16** (1.9 mg, *t_R* 13.7 min) and **12** (3.4 mg, *t_R* 18.4 min). Fraction R_{II} (1.3 g) was fractionated by silica gel CC, using a gradient of EtOAc-MeOH (1:0-6:4) to give 230 sub-fractions. Sub-fractions R_{II-50-55} contained the pure compound **11** (31.3 mg). Sub-fractions R_{II-59-63} (38.8 mg) was purified by semi prep HPLC using an isocratic elution of 15% CH₃CN during 25 min to give compounds **10** (14.3 mg, *t_R* 9.9 min) and **18** (6.4 mg, *t_R* 19.4 min) whereas the purification of the R_{II-70-80} (58.2 mg) in the same condition gave **19** (2.9 mg, *t_R* 22.1 min). Fraction R_{III} (1.2 g)

was applied to HPFC on normal phase silica gel, eluted with the system CH₂Cl₂-MeOH-H₂O [10:0:0 to 60:40:7], in 48 min to afford 60 sub-fractions. The purification of the sub-fractions R_{III-10-14} (24.8 mg) by semi prep HPLC in isocratic elution with 25% CH₃CN during 20 min, gave compounds **17** (4.7 mg, *t_R* 11.1 min), and **15** (6.9 mg, *t_R* 8.1 min). Fraction R_{IV} (198 mg) was purified by silica gel CC, eluted with the gradient of the mixture of CH₂Cl₂-MeOH-H₂O [95:5:0-70:30:5], to afford compound **9** (11.3 mg) as a white precipitate in pure form. Fraction R_V was purified previously ([Bendamene et al., 2020](#)).

The same protocol was used for the extraction of the aerial parts of *S. semipapposa* (700 mg). A part of the resulting 80% MeOH extract (50 g) was submitted to VLC over RP-C₁₈ eluted successively with MeOH-H₂O (3:7, 4:6, 6:4, 8:2 and 10:0), to give 5 main fractions (A_I to A_V, respectively) and a pure compound **22** (3mg) obtained as a precipitate in fraction A_{III}. This latter (4 g) was fractionated by flash chromatography over silica gel, using a binary gradient of CH₂Cl₂-MeOH [10:0 to 6:4] in 55 min to afford 146 sub-fractions. Sub-fractions A_{III-51-52} (51 mg), A_{III-57-60} (50 mg) and A_{III-72} (33.4 mg), were purified each one by MPLC on RP-C₁₈ eluted by gradient system 10%–50% CH₃CN, in 45 min to yield compounds **5** (4.4 mg) and **6** (2.6 mg), from A_{III-51-52}, **10** (8.5 mg), and a mixture of **3** and **1** (1.7 mg) from A_{III-57-60}, and **8** (15.2 mg) from A_{III-72}. Fraction A_{IV} (1.9g) was purified by flash chromatography over silica gel, eluted by a gradient system of CH₂Cl₂-MeOH [10:0 to 6:4] in 45 min to afford 100 sub-fractions. Compounds **21**(1mg), **1** (23.2 mg) and **7** (24.4 mg) were obtained as pure compounds in fractions A_{IV-25}, A_{IV-45} and A_{IV-66-67}, respectively, while the sub-fraction A_{IV-22} (31.5 mg) was purified by MPLC (20-55% CH₃CN, in 18 min) to yield 3.4 mg of compound **20**. Sub-fractions A_{IV-32} to A_{IV-35} (30 mg) were pooled and purified by MPLC (30-60% CH₃CN, in 20 min) to afford **23** (6 mg). The remaining sub-fraction A_{IV-53-54} (28.5 mg) was chromatographed by MPLC (20-50% CH₃CN, in 20 min) to furnish the pure compound **2** (8.5 mg), and the mixture of compounds **3** and **1** (1.9 mg), and **4** and **2** (1 mg).

4.3.1. *Semipapposiridoid A (1)*

Yellowish solid; $[\alpha]_D^{25}$ -97.7(c1, MeOH); IR (KBr) ν_{\max} (cm⁻¹):3356, 2924, 1695, 1629, 1603, 1514, 1440, 1371, 1260; UV λ_{\max} (MeOH) (log ϵ): 231 (0.177), 313 (0.193) nm; ¹H (600 MHz, CD₃OD) and ¹³C NMR (150 MHz, CD₃OD) data: see [Table1](#). HR-ESI-MS m/z : 959.3170[M+Na]⁺ (calcd for C₄₄H₅₆O₂₂, 959.3161).

4.3.2. *Semipapposiridoid B (2)*

Yellowish solid; $[\alpha]_D^{25}$ -40(c 0.16, MeOH); IR (KBr) ν_{\max} (cm⁻¹):3352, 2925, 1688, 1629, 1605, 1515, 1440, 1380, 1282; UV λ_{\max} (MeOH) (log ϵ): 228 (0.845), 308 (0.250) nm; ¹H (600 MHz, CD₃OD) and ¹³C NMR (150 MHz, CD₃OD) data: see [Table1](#). HR-ESI-MS m/z : 959.3169 [M+Na]⁺ (calcd for C₄₄H₅₆O₂₂, 959.3161).

4.3.3. *Semipapposiridoid C (3)*

Yellowish solid; $[\alpha]_D^{25}$ -66(c 0.17, MeOH); IR (KBr) ν_{\max} (cm⁻¹): 3357, 2924, 1695, 1630, 1067; UV λ_{\max} (MeOH) (log ϵ): 231 (1.266), 312 (0.630) nm; ¹H (600 MHz, CD₃OD) and ¹³C NMR (150 MHz, CD₃OD) data: see [Table 1](#). HR-ESI-MS m/z : 959.3156 [M+Na]⁺ (calcd for C₄₄H₅₆O₂₂,959.3161).

4.3.4. *Semipapposiridoid D(4)*

Yellowish solid; $[\alpha]_D^{25}$ 13.3(c0.18, MeOH); IR(KBr) ν_{\max} (cm⁻¹):3358, 2920, 1695, 1630,1068; UV λ_{\max} (MeOH) (log ϵ): 231 (0.724), 312 (0.304) nm; ¹H (600 MHz, CD₃OD) and ¹³C NMR (150 MHz, CD₃OD) data: see [Table1](#). HR-ESI-MS m/z : 959.3169 [M+Na]⁺ (calcd for C₄₄H₅₆O₂₂, 959.3161).

4.3.5. *Semipapposiridoid E(5)*

Yellowish solid: $[\alpha]_D^{25}$ -58 (c0.44, MeOH); IR (KBr) ν_{\max} (cm⁻¹): 3377, 2926, 1692, 1629, 1449, 1386, 1277; UV λ_{\max} (MeOH) (log ϵ): 235 (0.580) nm; ¹H (600 MHz, CD₃OD) and ¹³C NMR

(150 MHz, CD₃OD) data: see [Table 2](#). HR-ESI-MS m/z : 567.2413 [M+Na]⁺ (calcd for C₂₆H₄₀O₁₂, 567.2417).

4.3.6. *Semipapposiridoid F (6)*

Yellowish solid; $[\alpha]_D^{25}$ -14 (c0.26, MeOH); IR (KBr) ν_{\max} (cm⁻¹): 3376, 2926, 1693, 1629, 1451, 1387, 1262; UV λ_{\max} (MeOH) (log ϵ): 235 (1.761) nm; ¹H (600 MHz, CD₃OD) and ¹³C NMR (150 MHz, CD₃OD) data: see [Table 2](#). HR-ESI-MS m/z : 567.2408 [M+Na]⁺ (calcd for C₂₆H₄₀O₁₂, 567.2417).

4.4. Determination of total phenolic (TPC) and flavonoid contents (TFC)

Total phenolic content (TPC) of 80% MeOH extracts of aerial parts, roots, and fractions (R_{I-V} and A_{I-IV}) of *S. semipapposa* ([Table 3](#)) was determined spectrophotometrically using Folin-Ciocalteu method as previously reported ([Singleton and Rossi, 1965](#)) with slight modifications ([Muller et al., 2010](#)). The absorbance was measured at 765 nm and the results were expressed as micrograms of gallic acid equivalent per milligram of extract ($\mu\text{g GAE/mg}$).

The total flavonoid content (TFC) was analyzed using an adapted aluminum chloride colorimetric reported by [Topçu et al. \(2007\)](#). The absorbance was measured at 415 nm and the results were expressed as mg of quercetin equivalents per milligram of extract ($\mu\text{g QE/mg}$).

4.5. Antioxidant activities

Evaluation of the antioxidant activities of 80% MeOH extracts of aerial parts, roots, and fractions (R_{I-RV} and A_{I-AIV}) was carried out by three different methods including scavenging of the free radicals of 2,2-diphenyl-1-picrylhydrazyl (DPPH) and 2,2'-azinobis-(3-ethylbenzothiazoline-6-sulfonate) (ABTS) and cupric reducing antioxidant capacity (CUPRAC) assays.

4.5.1. DPPH free radical scavenging activity

The free radical scavenging activity of the 80% MeOH extracts and the fractions of *S. semipapposa* was determined by the DPPH assay described by [Blois \(1958\)](#) with few

modifications as previously described (Boutaghane et al., 2018). Briefly, 40 μL of either standard or sample solutions was mixed with 160 μL of DPPH solution in microplate. The mixture was then incubated at room temperature in the dark for 30 min at 37°C, and the absorbance was measured at λ 517 nm and expressed as percentage of DPPH radical scavenging calculated as follows: $\text{DPPH scavenging\%} = [(\text{Ab}_{\text{control}} - \text{Ab}_{\text{sample}})/\text{Ab}_{\text{control}}] \times 100$.

The sample concentration providing 50% free radical scavenging activity (FRS₅₀) was calculated from the graph of DPPH scavenging effect percentage against sample concentration. BHA and BHT were used as positive controls. All the tests were conducted in triplicate. The experimental data were expressed as mean \pm standard deviation.

4.5.2. ABTS radical scavenging activity

The ABTS scavenging activity of the 80% MeOH extracts and the obtained fractions was performed spectrophotometrically according to a literature method (Re et al. 1999) with a little modification. The ABTS^{•+} was produced by the reaction between 7 mM ABTS in water and 2.45 mM potassium persulfate, stored in the dark at room temperature for 12 h. Before usage, the ABTS^{•+} solution was diluted to get an absorbance of 0.70 ± 0.02 at 734 nm with methanol. Then 40 μL of either standard or sample solutions was added to 160 μL of ABTS^{•+} solution in microplate). After 10 min, the percentage inhibition at λ 734 nm was calculated for each concentration relative to a blank absorbance (methanol). The scavenging capability of ABTS^{•+} was calculated using the following equation: $\text{ABTS}^{\bullet+}\text{scavenging\%} = [(\text{Ab}_{\text{control}} - \text{Ab}_{\text{sample}})/\text{Ab}_{\text{control}}] \times 100$.

The sample concentration providing 50% cation radical scavenging activity (CRS₅₀) was calculated from the graph of ABTS^{•+} scavenging effect percentage against sample concentration. BHA and BHT were used as positive controls. All the tests were conducted in triplicate. The experimental data were expressed as mean \pm standard deviation.

4.5.3. Power cupric ion reducing (CUPRAC)

80% MeOH extracts and fractions were tested for their power cupric ion reducing (CUPRAC) capacity assay using the method of [Apak et al. \(2004\)](#) with some modifications. Sixty microliters of ammonium acetate buffer solution (1 M, pH = 7.0) were added to 40 μ L of a solution of copper (II) chloride (10 mM) and 50 μ L of the neocuprine solution (7.5 mM). Different concentrations of extracts, fractions and standards were added to the initial mixture to make a final volume of 200 μ L. After 1 h, the absorbance at 450 nm was recorded against a reagent blank by using a 96-well microplate reader. The power cupric ion reducing was calculated as follows: $[1 - A_0 / (A_1 - A_2)] \times 100$, where A_0 is absorbance of the control (without sample), A_1 is absorbance in the presence of the sample and A_2 is absorbance of the blank. All the tests were conducted in triplicate and the reduction capacity of the extracts was compared with BHA and BHT standards.

Declaration of competing interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Acknowledgements

The authors are grateful to the DGRSDT of the Ministry of Higher Education and Scientific Research of Algeria for PRFU Project (B00L01UN250120180001), Biotechnology Research Center (CRBt) (Constantine-Algeria), Conseil Regional Champagne Ardenne, Conseil General de la Marne, Ministry of Higher Education and Research (MESR) in France, and to the PLANET CPER project for financial support.

Appendix A. Supplementary data

Supplementary material related to this article can be found, in the online version.




Fig. 1. Structures of undescribed bis-iridoids **1**–**6** isolated from *S. semipapposa*




Fig. 2. Selected HMBC and NOESY correlations of compound **1**

Table 1 ¹³C NMR and ¹H NMR spectroscopic data of compounds **1-4** in CD₃OD.^a

Position	1		2		3		4	
	δ_{C}	δ_{H}	δ_{C}	δ_{H}	δ_{C}	δ_{H}	δ_{C}	δ_{H}
Unit A								
1a	97.7	5.58, d (6.6)	97.7	5.56, d (6.5)	97.7	5.59, d (6.5)	97.7	5.56, d (6.6)
3a	153.9	7.52, s	153.7	7.49, d (1.3)	153.7	7.52, (s 1)	153.7	7.49, d (2.8)
4a	111.6	-	111.6	-	111.6	-	111.6	-
5a	31.3	2.90, q (6.3)	31.2	2.87, q (6.4)	31.3	2.90, q (6.3)	31.2	2.87, q (6.3)
6a	30.1	1.86, td (13.8, 7.4) 2.12, dd (13.8, 7.2)	30.1	1.81, dq (13.8, 7.2) 2.02, m	30.1	1.85, dq (13.7, 7.1) 2.04, m	30.2	1.82, dq (13.8, 6.7) 2.01, m
7a	64.1	4.11, m 4.14, m	64.1	4.08, m 4.12, m	64.1	4.11, m 4.14, m	64.1	4.08, m 4.12, m
8a	135.7	5.80, ddd (17.4, 10.4, 8.6)	135.7	5.78, ddd (14.4-10.4-8.9)	135.7	5.80, q (8.5)	135.7	5.78, dd (9.8, 8.8)
9a	45.4	2.70, dt (8.3, 6.1)	45.3	2.68, dd (8.2-6.1)	45.3	2.69, dt (8.2, 5.8)	45.3	2.67, dd (17.2, 10.4, 6.3)
10a	119.6	5.28, dd (10.4-1.5) 5.33, d (17.4)	119.6	5.26, d (10.4) 5.31, d (14.4)	119.6	5.28, d (10.7) 5.33, dd (14.7, 1.4)	119.6	5.25, d (10.4) 5.32, d (17.2)
11aC=O	168.2	-	168.1	-	168.3	-	168.2	-
12aC=O	173.0	-	173.0	-	173.0	-	173.0	-
13aCH ₃	20.9	2.04, s	21.0	2.02, s	20.9	2.04, s	21.0	2.18, s
1a-O-glc								
1'	100.1	4.74, d (7.9)	100.6	4.68, d (8.0)	100.1	4.74, d (7.9)	100.1	4.71, d (8.0)
2'	74.7	3.23, dd (9.0, 7.9)	74.6	3.22, dd (9.0, 8.0)	74.8	3.22, t (8.7)	74.7	3.22, dd (8.8, 8.0)
3'	78.0	3.40, t (9.0)	78.0	3.39, t (9.0)	78.0	3.39, t (9.1)	78.0	3.39, t (8.8)
4'	71.6	3.30, t (9.2)	71.5	3.29, t (9.5)	71.6	3.29, t (9.2)	71.6	3.29, t (9.6)
5'	78.4	3.33, m	78.4	3.34, m	78.5	3.33, m	78.4	3.35, m
6'	62.8	3.69, dd (11.9, 6.0) 3.93, dd (11.9, 2.0)	62.8	3.68, dd (11.9, 6.2) 3.92, dd (11.9, 1.9)	62.8	3.68, dd (11.9, 4.6) 3.93, dd (11.9, 1.8)	62.1	3.68, dd (12.0, 6.1) 3.92, d (12.0, 1.3)
Unit B								
1b	97.6	5.33, d (4.6)	97.3	5.08, d (5.7)	97.6	5.34, d (4.8)	98.0	5.09, d (5.5)
3b	152.5	7.47, d (1.3)	152.8	7.44, d (1.1)	152.5	7.45, d (1.2)	152.6	7.44, d (1.3)
4b	113.3	-	112.9	-	113.3	-	112.9	-
5b	32.6	3.15 q (7.9)	33.1	3.09, q (8.4)	32.6	3.15, q (7.9)	32.7	3.10, q (8.0)
6b	40.3	1.78, ddd (14.1, 7.8, 5.1) 2.33, ddd (14.1, 7.7, 1.5)	40.6	1.61, ddd (14.4, 8.9, 4.8) 2.26, dd (14.4, 7.7)	40.3	1.78, ddd (14.5, 7.7, 5.0) 2.33, td (14.5, 7.7)	40.5	1.61, ddd, (11.3, 9.0, 5.0) 2.27, m
7b	78.4	5.24, t (4.5)	78.2	5.22, t (4.7)	78.4	5.24, t (4.5)	78.4	5.23, t (4.6)
8b	41.0	2.18, m	41.2	2.12, m	41.1	2.18, m	41.1	2.09, m
9b	47.1	2.13, ddd (17.4, 8.8, 4.8)	46.9	2.02, m	47.1	2.12, ddd (13.8, 8.9, 5.0)	47.1	2.02, m
10b	13.8	1.10, d (6.8)	14.1	1.02, d (7.1)	13.8	1.10, d (6.7)	13.9	1.02, d (7.1)
11b-C=O	169.3	-	169.2	-	169.3	-	169.3	-
12b-OCH ₃	51.7	3.71, s	-	-	51.8	3.72, s	-	-
1b-O-glc								
1''	100.2	4.76, d (7.9)	100.1	4.69, d (7.9)	100.2	4.77, d (7.9)	100.6	4.66, d (7.8)
2''	74.8	3.34, dd (9.3, 7.9)	74.6	3.25, dd (9.0, 7.9)	74.8	3.33, m	74.7	3.23, dd (9.5, 7.8)
3''	75.7	3.68, t (9.2)	77.8	3.42, t (9.0)	75.7	3.68, t (9.1)	77.8	3.40, t (8.9)
4''	72.4	4.87, t (9.5)	71.7	3.39, t (9.0)	72.0	4.86, t (9.5)	71.7	3.30, t (9.0)
5''	76.5	3.60, m	75.7	3.59, m	76.5	3.59, m	75.6	3.55, m
6''	62.5	3.60, m 3.66, m	64.0	4.45, dd (11.9, 6.4) 4.54, dd (11.9, 2.1)	62.5	3.57, m 3.66, m	64.1	4.37, dd (12.0, 6.5) 4.50, dd (12.0, 2.1)
4''-O-E-P-coumaroyl								
1'''	127.1	-	127.0	-	127.5	-	127.5	-
2'''	131.3	7.49, d (8.6)	131.3	7.48, d (8.5)	134.0	7.72, d (8.6)	133.9	7.68, d (8.7)
3'''	116.9	6.83, d (8.6)	116.9	6.82, d (8.5)	115.8	6.77, d (8.6)	115.9	6.78, d (8.7)
4'''	161.4	-	161.4	-	160.3	-	160.5	-
5'''	116.9	6.83, d (8.6)	116.9	6.82, d (8.5)	115.8	6.77, d (8.6)	115.9	6.78, d (8.7)
6'''	131.3	7.49, d (8.6)	131.3	7.48, d (8.5)	134.0	7.72, d (8.6)	133.9	7.68, d (8.7)
7'''	147.3	7.69, d (15.9)	146.9	7.67, d (15.9)	146.2	6.94, d (12.8)	145.5	6.90, d (12.9)
8'''	114.7	6.39, d (15.9)	114.9	6.37, d (15.9)	115.9	5.83, d (12.8)	116.2	5.80, d (12.9)
9'''	168.5	-	168.9	-	167.3	-	170.8	-

^a in ppm, *J* in parentheses in Hz.

Table 2 ^{13}C NMR and ^1H NMR spectroscopic data of compounds **5-6** in CD_3OD .^a

Position	5		6	
	$\delta_{\text{C}}/\delta_{\text{H}}$	$\delta_{\text{C}}/\delta_{\text{H}}$	$\delta_{\text{C}}/\delta_{\text{H}}$	$\delta_{\text{C}}/\delta_{\text{H}}$
Unit A				
1a	97.7	5.57, d (6.5)	97.7	5.56, d (6.5)
3a	153.4	7.47 s	153.4	7.48, s
4a	112.0	-	112.0	-
5a	31.0	2.84, q (6.6)	31.0	2.84, q (6.4)
6a	33.1	1.73, ddd (13.7-7.7-6.2)	33.1	1.72, m
		1.88 m		1.86, m
7a	61.2	3.55, m	61.2	3.55, m
		3.60, m		3.60, m
8a	135.9	5.80, ddd (17.4-10.4-8.7)	135.9	5.80, ddd (17.3, 10.7, 8.7)
9a	45.4	2.65, dt (8.2, 6.1)	45.4	2.65, dt (8.1-6.0)
10a	119.3	5.26, dd (10.4-1.2)	119.3	5.25, d (10.7-1.3)
		5.31, dd (17.4-1.2)		5.30, d (17.3)
11-C=O	168.7	-	168.7	-
1A-O-glc				
1'	100.1	4.71, d (7.9)	100.1	4.71, d (7.9)
2'	74.6	3.21, dd (9.1-7.9)	74.6	3.20, dd (9.0-7.9)
3'	78.0	3.38, t (8.8)	78.0	3.38, t (8.8)
4'	71.5	3.28, dd (9.7-8.7)	71.6	3.28, dd (9.7-8.7)
5'	78.4	3.32 m	78.4	3.31 m
6'	62.8	3.68, dd (11.9-6.1)	62.8	3.68, dd (11.9-6.0)
		3.92, dd (11.9-2.0)		3.92, dd (11.9-2.0)
Unit B				
1b	64.2	3.79, dd (12.1, 4.8)	59.3	3.52, dd (11.9-3.7)
		3.83, dd (12.1, 5.0)		3.99, dd (11.9-1.4)
3b	102.9	4.42, dd (8.9, 3.3)	99.3	4.67, t (3.5)
4b	35.4	1.28, ddd (13.3-9.7-8.9)	33.1	1.55, ddd (14.1, 10.0, 3.9)
		1.84, m		1.68, ddd (14.2-6.2-3.2)
5b	34.4	2.34, m	31.1	2.51, m
6b	39.8	1.84, m	39.8	1.74, ddd (13.6-7.6-6.0)
		1.97, ddd (14.6-8.1-2.5)		1.86, m
7b	79.1	5.29, m	78.7	5.32, m
8b	39.3	2.13, dq (11.5, 6.3)	39.0	2.28, dq (11.1, 6.6)
9b	44.8	1.83, m	44.6	1.98, m
10b	12.8	0.88, d (6.8)	12.9	1.00, d (6.9)
11b-OCH₃	56.0	3.42, s	55.0	3.37, s

^a in ppm, *J* in parentheses in Hz.

Table 3. Phenolic, flavonoid contents and antioxidant activity by the DPPH[•], ABTS^{•+}, and CUPRAC assays of extracts and fractions from *Scabiosa semipapposa*. FRS₅₀, CRS₅₀ and A_{0.50} values represent the means ± S.D of three parallel measurements

	TPC	TFC	DPPH	ABTS	CUPRAC
Extracts	(µg GAE/mg)	(µg QE/mg)	FRS ₅₀ (µg/mL)	CRS ₅₀ (µg/mL)	A _{0.50} (µg/mL)
Roots 80% MeOH extract (R)	109.0±0.5	12.8±1.7	80.0±0.6	49.5 ±1.2	122.2±1.0
Fraction R_I	361.7±1.2	74.6±1.4	4.9 ±0.1	8.9±0.3	10.2±0.1
Fraction R_{II}	266.8±0.6	37.4±0.1	34.3±0.6	12.7±0.4	29.1±0.4
Fraction R_{III}	150.8±1.4	10.5 ±0.5	64.6 ±0.1	23.3 ±0.9	54.8±0.8
Fraction R_{IV}	158.9±1.6	43.0±0.6	62.1±0.8	36.2±0.6	44.1±1.9
Fraction R_V	32.3 ±0.3	5.7±0.4	315.8±0.7	124.0±0.9	378.5 ±1.4
Aerial parts 80% MeOH extract (A)	169.6±0.2	19.5±0.0	80.0±0.5	31.1 ±0.7	92.2 ± 1.7
Fraction A_I	190.8±0.9	20.9±1.8	45.1 ± 0.8	27.3±0.7	32.9 ± 1.1
Fraction A_{II}	243.9±0.8	44.2 ±1.6	46.0 ± 1.8	18.8 ±0.6	35.5 ± 0.5
Fraction A_{III}	146.7 ±0.8	57.2 ±1.4	42.1 ± 0.1	23.5 ±1.3	37.1 ±0.8
Fraction A_{IV}	183.9 ± 0.9	62.2 ±2.7	46.9 ± 0.6	41.7 ± 0.9	40.9 ± 0.2
BHT^a			15.9 ± 0.1	2.7±0.2	5.4±0.3
BHA^a			8.7 ±0.1	1.8 ± 0.1	3.1 ±0.1

^aUsed as a positive control

References

- Abdallah, O.M., Phenolic glucoside and other constituents of *Dipsacus laciniatus*, 1991.[https://doi.org/10.1016/0031-9422\(91\)85153-Q](https://doi.org/10.1016/0031-9422(91)85153-Q)
- Agrawal, P.K.,1992. NMR Spectroscopy in the structural elucidation of oligosaccharides and glycosides. *Phytochemistry* 31, 3307–3330.[https://doi.org/10.1016/0031-9422\(92\)83678-R](https://doi.org/10.1016/0031-9422(92)83678-R)
- Akar, Z. 2022. Chemical compositions by using LC–MS/MS and GC–MS and antioxidant activities of methanolic extracts from leaf and flower parts of *Scabiosa columbaria* subsp. *columbaria* var. *columbaria* L.Saudi. *J. Biol. Sci.*28, 6639–6644. <http://doi.org/10.1016/j.sjbs.2021.07.039>
- Alimbaeva, P.K., Akimaliev, A., Mukhamedziev, M.M., 1977. Triterpene glycosides of some representatives of the Dipsacaceae family. *Khim. Prir. Soedin.* 5, 708–709. <https://doi.org/10.1007/BF00569610>.
- Al-Qudah, M.A., Ootom, N.K., Al-Jaber, H.I., Saleh, A.M., Abu Zarga, M.H., Afifi, F.U., AbuOrabi, S.T., 2017. New flavonol glycoside from *Scabiosa prolifera* L. aerial parts with *in vitro* antioxidant and cytotoxic activities. *Nat. Prod. Res.* 31, 2865–2874. <https://doi.org/10.1080/14786419.2017.1305377>
- Apak, R., Güçlü, K., Özyürek, M., Çelik, S.E., 2008. Mechanism of antioxidant capacity assays and the CUPRAC (cupric ion reducing antioxidant capacity) assay. *Microchimica Acta* 160, 413–419. <http://doi.org/10.1007/s00604-007-0777-0>
- Bammi, J., Douira, A.,2002. Les plantes médicinales dans la forêt de L’Achach (Plateau Central, Maroc). *Acta. Bot. Malacit.*27, 131–145. <https://doi.org/10.24310/abm.v27i0.7323>
- Baykal, T., Panayir, T., Tasdemir, D., Sticher, O., Çalis, I., 1998. Triterpene saponins from *Scabiosa rotata*. *Phytochemistry* 48, 867–873. [https://doi.org/10.1016/s0031-9422\(97\)00982-5](https://doi.org/10.1016/s0031-9422(97)00982-5)

- Bendamene,S., Boutaghane,N., Bellik,Y., Sayagh, C., Alabdul Magid, A., Harakat, D., Kabouche,Z., Voutquenne-Nazabadioko,L., 2020. Semipapposides A-M, triterpenoid bidesmosides saponins from the roots of *Scabiosa semipapposa*. *Phytochemistry* 180,112526. <https://doi.org/10.1016/j.phytochem.2020.112526>
- Blois, M.S., 1958. Antioxidant determinations by the use of a stable free radical. *Nature* 181, 1199–1200. <https://doi.org/10.1038/1811199a0>
- Bonet, M.À., Parada, M., Selga, A., Vallès, J.,1999. Studies on pharmaceutical ethnobotany in the regions of L'Alt Empordà and Les Guilleries (Catalonia, Iberian Peninsula). *J.Ethnopharmacol.* 68, 145–168.[https://doi.org/10.1016/S0378-8741\(99\)00083-5](https://doi.org/10.1016/S0378-8741(99)00083-5)
- Boutaghane, N., Alabdul Magid, A., Abedini, A., Cafolla, A., Djeghim, H., Gangloff, SC.,2018. Chemical constituents of *Genista numidica* Spach aerial parts and their antimicrobial, antioxidant and antityrosinase activities. *Nat. Prod. Res.* 6419, 1–7.<https://doi.org/10.1080/14786419.2018.1437425>
- Breitmaier, E., Voelter, W., 1987. Carbon-13 NMR spectroscopy.3rd Edition. <https://doi.org/10.1002/mrc.1260260415>
- Calis, I., Lahloub, M.f., Sticher,O.,1984. Loganin, loganicacid periclymenoside, a new biosidicesteriridoid glucoside from *Lonicera periclymenum* L. (Caprifoliaceae). *Helv. Chim. Acta* 67, 160–165. <https://doi.org/10.1002/HLCA.19840670119>
- Calis, I. ,Sticher. O., 1984. Secoiridoid glucosides from *Lonicera*. *Phytochemistry* 23, 2539–2540.[https://doi.org/10.1016/s0031-9422\(00\)84094-7](https://doi.org/10.1016/s0031-9422(00)84094-7)
- Chiruvella, K.K., Mohammed, A., Dampuri, G. Ghanta, R.G., Raghavan, S.C., 2007. Phytochemical and antimicrobial studies of methyl angolensate and luteolin-7-O-glucoside isolated from callus cultures of *Soymida febrifuga*. *Intern.J. Biomed. Sci.* 3, 269-78.PMID: [23675053](https://pubmed.ncbi.nlm.nih.gov/23675053/)

- Christopoulou, C., Graikou, K., Chinou, I., 2008. Chemosystematic value of chemical constituents from *Scabiosahymettia*(Dipsacaceae). Chem. and Biodiv. 5, 318–323. <https://doi.org/10.1002/cbdv.200890029>
- Cowan, S., Stewart, M., Abbiw, D. K., Latif, Z., Sarker, S. D., Nash, R. J., 2001. Lignans from *Strophanthus gratus*. Fitoterapia72, 80–82. [https://doi.org/10.1016/S0367-326X\(00\)00240-9](https://doi.org/10.1016/S0367-326X(00)00240-9)
- Deyama, T., Ikawa, T., Kitagawa, S., Neshibe, S., 1986. The constituent of *Eucommia ulmoides* OLIV. III. Isolation and structure of a new lignan glycoside. Chem. Pharm. Bull. 32, 325–527. <https://doi.org/10.1248/cpb.34.523>
- Dinda, B., Debnath, S., Majumder, S., Arima, S., Sato, N., Harigaya Y., 2006. A new bisiridoid glucoside from *Mussaenda incana*. Chin. Chem. Lett. 10, 1331–4. <https://doi.org/10.1007/s11418-008-0273-9>
- Dong, F.W., Wu, Z.K., Yang, L., Zi, C.T., Yang, D., Ma, R.J., Liu, Z.H., Luo, H.R., Zhou, J., Hu, J.M., 2015. Iridoids and sesquiterpenoids of *Valeriana stenoptera* and their effects on NGF-induced neurite outgrowth in PC12 cells. Phytochemistry 118, 51–60. <https://doi.org/10.1016/j.phytochem.2015.08.015>
- El-Ghazooly, M.G., El-lakany, M.A., Abou-Shoer, M.I., Aly, A.H., 2003. Chemical constituents of *Helichrysum conglobatum* growing in Egypt. Nat. Prod. Sci. 9, 213–219.
- Gali, L., Bdjou, F., 2019. Antioxidant and anticholinesterase effects of the ethanol extract, ethanol extract fractions and total alkaloids from the cultivated *Ruta chalepensis*. South African J. Botany 120 (Special Issue: Enzyme Inhibitors of Natural Origin), 163–169. <https://doi.org/10.1016/j.sajb.2018.04.011>
- Garaev, E.A., Movsumov, I.S., Isaev, M.I., 2008. Flavonoids and oleanolic acid from *Scabiosa caucasica*. Chem. Nat. Compd. 44, 520–521. . <https://doi.org/10.1007/s10600-008-9108-x>

- Girre, L., 1980. Connaitre et reconnaitre les plantes médicinales. Ouest, France, Rennes.
- Graikou, K., Aliginnis, N., Chinou, I.B., Hava, C., 2002. Cantleyoside-dimethyl-acetal and other iridoid glucoside from *Pterocephalus perennis* - antimicrobial activities. Z. Naturforsch. 57C, (1-2), 95–99. <https://doi.org/10.1515/znc-2002-1-217>
- Gülcemal, D., Masullo, M., AlankuşÇalışkan, O., Karayildirim, T., Şenol, S.G., Piacente, S., Bedir, E., 2010. Monoterpenoid glucoindole alkaloids and iridoids from *Pterocephalus pinardii*. Magn. Reson. Chem. 48, 239–243. <https://doi.org/10.1002/mrc.2559>
- Hlila, M.B., Omri, A., Ben Jannet, H., Lamari, A., Aouni, M., Selmi, B., 2015. Acetylcholinesterase inhibitory and antioxidant properties of roots extracts from the Tunisian *Scabiosa arenaria* Forssk. Industrial Crops and Products 67, 62–69. <https://doi.org/10.1016/j.indcrop.2015.01.009>
- Horn, M.M., Drewes, S.E., Brown, N.J., Munro, O.Q., Marion Meyer, J.J., Mathekga, A-D.M., 2001. Transformation of naturally-occurring 1, 9-trans-9,5-cis sweroside to all trans sweroside during acetylation of sweroside aglycone. Phytochemistry 57, 51–56. [https://doi.org/10.1016/S0031-9422\(00\)00460-X](https://doi.org/10.1016/S0031-9422(00)00460-X)
- Jayaprakasha, G.K., Singh, R.P., Sakariah, K.K., 2001. Antioxidant activity of grape seed *Vitis vinifera* extracts on peroxidation models *in vitro*. Food Chem. 73, 285–290. [https://doi.org/10.1016/S0308-8146\(00\)00298-3](https://doi.org/10.1016/S0308-8146(00)00298-3)
- Jensen, S.R., Lyse-Petersen, S.E., Nielsen, B.J., 1979. Novel bis-iridoid glucosides from *Dipsacus sylvestris*. Phytochemistry 18, 273–277. [https://doi.org/10.1016/0031-9422\(79\)80069-2](https://doi.org/10.1016/0031-9422(79)80069-2)
- Ji, D., Zhang, C.F., Jing, Z., Yang, H.W., Shen, J.Y., Yang, Z.G. 2012 A new iridoid glycoside from the roots of *Dipsacus asper*. Molecules 17, 1419–24. <https://doi.org/10.3390/molecules17021419>

- Kanchanapoom, T., Kasaia, R., Yamasakia, K. 2002. Iridoid and phenolic glycosides from *Morinda coreia*. *Phytochemistry* 59, 551–556. [https://doi.org/10.1016/s0031-9422\(01\)00426-5](https://doi.org/10.1016/s0031-9422(01)00426-5)
- Kanho, H., Yaoya, S., Kawahara, N., Nakane, T., Takase, Y., Masuda, K., Kuroyanagi, M., 2005. Biotransformation of benzaldehyde-type and Acetophenone-type derivatives by *Pharbitis nil* hairy Roots. *Chem. Pharm. Bull.* 53, 361–365. <https://doi.org/10.1248/cpb.53.361>
- Kılınç, H., Masullo, M., D’Urso, G., KaraYildirim, T., Alankus, O., Piacente, S., 2020. Phytochemical investigation of *Scabiosa sicula* guided by a preliminary HPLC-ESIMSⁿ profiling. *Phytochemistry* 174, 112350. <https://doi.org/10.1016/j>
- Kocsis, A., Szabó, L. F., Podányi B., 1993. New Bis-Iridoids from *Dipsacus laciniatus*. *J. Nat. Prod.* 56, 1486–1499. <https://doi.org/10.1021/np50099a007>
- Kose, L.S., Moteetee, A., Vuuren, S.V., 2015. Ethnobotanical survey of medicinal plants used in the Maseru district of Lesotho. *J. Ethnopharmacol.* 170, 184–200. <https://doi.org/10.1016/j.jep.2015.04.047>
- Lehbil, M., AlabdulMagid, A., Kabouche, A., Voutquenne-Nazabadioko, L., Morjani, H., Harakat, D., Kabouche, Z., 2018a. Triterpenoidsaponins from *Scabiosa stelatta* collected in the North-eastern Algeria. *Phytochemistry* 150, 40–49. <https://doi.org/10.1016/j.phytochem.2018.03.005>
- Lehbil, M., AlabdulMagid, A., Hubert, J., Kabouche, A., Voutquenne-Nazabadioko, L., Renault, J.-H., Nuzillard, J.-M., Morjani, H., Abedini, A., Gangloff, S.-C., Kabouche, A., 2018b. Two new bis-iridoids isolated from *Scabiosa stellata* and their antibacterial, antioxidant, anti-tyrosinase and cytotoxic activities. *Fitoterapia* 125, 41–48. <https://doi.org/10.1016/j.fitote.2017.12.018>

- Li, Y.K., Li, W., Fu, C.M., Song, Y., Fu, Q., 2019. *Lonicera japonica* flos and *Lonicera* flos: a systematic review of ethnopharmacology, phytochemistry and pharmacology. *PhytochemRev.* 19, 1–61. <https://doi.org/10.1007/s11101-019-09655-7>
- Lin, L.W., Zhang, X.G., Zhu, J.J., Gao, H.M., Wang, Z.M., Wang, H.W., 2008. Two new triterpenoid saponins from the flowers and buds of *Lonicera japonica*. *J. Asian Nat. Prod. Res.* 10, 925–929. <https://doi.org/10.1080/10286020802217366>
- Liu, H., Liu, D., Jiang, M.Y., Zhao, X.D., Li, R.T., Li, H.M., 2021. Iridoids from *Valeriana jatamansi* with anti-inflammatory and antiproliferative properties. *Phytochemistry* 184, 112681. <https://doi.org/10.1016/j.phytochem.2021.112681>
- Liu, Z.X., Liu, C.T., Liu, Q.B., Ren, J., Li, L.Z., Huang, X.X., Wang, Z.Z., Song, S.J., 2015. Iridoid glycosides from the flower buds of *Lonicera japonica* and their nitric oxide production and α -glucosidase inhibitory activities. *J. Funct. Foods* 18, 512–519. <https://doi.org/10.1016/j.jff.2015.08.017>
- Moteetee, A., Kose, L.S., 2016. Medicinal plants used in Lesotho for treatment of reproductive and post reproductive problems. *J. Ethnopharmacol.* 194, 827–849. <https://doi.org/10.1016/j.jep.2016.10.062>
- Ming, D.S., Yu, D.Q., Yang, Y.Y., and He, C.H., 1997. The structures of three novel sesquiterpenoids from *Valeriana jatamansi* Jones. *Tetrahedron Lett.* 38, 5205–5208. [https://doi.org/10.1016/S0040-4039\(97\)01112-X](https://doi.org/10.1016/S0040-4039(97)01112-X)
- Mouffouk, C., Hambaba, L., Haba, H., Mouffouk, S., Bensouici, C., Mouffouk, S., Hachemi, M., Khadraoui, H., 2018. Acute toxicity and *in vivo* anti-inflammatory effects and *in vitro* antioxidant and anti-arthritic potential of *Scabiosa stellate*. *Oriental Pharmacy and Experimental Medicine* 18, 335–348. <https://doi.org/10.1007/s13596-018-0320-3>

- Müller, L., Gnoyke, S., Popken, A.M., Böhm, V., 2010. Antioxidant capacity and related parameters of different fruit formulations. *Food Sci. Technol.*43, 992-999. <https://doi.org/10.1016/j.lwt.2010.02.004>
- Murai, F., Tagawa, M., Matsuda, S., Kikuchi, T., Uesato, S, Inouye, H., Inouye, H.,1985. Abeliosides A and B, secoiridoid glucosides from *Abelia grandiflora*. *Phytochemistry* 24, 2329–35. [https://doi.org/10.1016/s0031-9422\(00\)83036-8](https://doi.org/10.1016/s0031-9422(00)83036-8)
- Mustafayeva, K., Giorgio,C.D., Elias,R., Kerimov,Y., Ollivier,E., De Meo, D., 2010. DNA-Damaging, mutagenic, and clastogenic activities of gentiopicroside isolated from *Cephalaria kotschyi* roots, *J. Nat. Prod.*73, 99–103. <https://doi.org/10.1021/np900322c>
- Okamura, N., Haraguchi, H., Hashimoto, K., Yagi, A., 1994. Flavonoids in *Rosmarinus officinalis* leaves. *Phytochemistry* 37, 1463–1466. [https://doi.org/10.1016/s0031-9422\(00\)90434-5](https://doi.org/10.1016/s0031-9422(00)90434-5)
- Pacifico, S.,D’Abrosca, B., Pascarella, M.T., Letizia, M.,Uzzo, P.,Piscopo, V.,Fiorentino, A. 2009.Antioxidant efficacy of iridoid and phenylethanoid glycosides from the medicinal plant *Teucrium chamaedris*in cell-free systems. *Bioorg. Med. Chem.*17, 6173–6179.<https://doi.org/10.1016/j.bmc.2009.07.065>
- Papalexandrou, A., Magiatis, P., Perdetzoglou, D., Skaltsounis, A.L., Chinou, I.B.,Harvala, C., 2003. Iridoids from *Scabiosa variifolia* (Dipsacaceae) growing in Greece. *Biochem. Syst. Ecol.* 31, 91–93. [https://doi.org/10.1016/S0305-1978\(02\)00070-4](https://doi.org/10.1016/S0305-1978(02)00070-4)
- Pasi,S., Aligiannis,N., Pratsinis, H., Skaltsounis, A.L., Chinou, I.B.,2009. Biologically active triterpenoids from *Cephalaria ambrosioides*. *Planta Med.* 75, 163–167. <https://doi.org/10.1055/s-0028-1088391>
- Polat, E., Alankus-Caliskan, O., Karayildirim, T., Bedir, E., 2010. Iridoids from *Scabiosa atropurpurea* L. subs p. *maritime* Arc. (L.). *Biochem. Syst. Ecol.* 38, 253–255. <https://doi.org/10.1016/j.bse.2010.01.004>

- Pinto, D.C.G.A., Rahmouni, N., Beghidja, N., Silva, A.M.S., 2018. *Scabiosa* genus: a rich source of bioactive metabolites. *Medicines* 5, 110. <https://doi.org/10.3390/medicines5040110>
- Qingwei, W., Xing, L., Yeqi, L., Xiaohui, X., Tao, L., Ni, Z., Kintoko, K., Renbin, H., 2012. Phenolic and lignan glycosides from the butanol extract of *Averrhoa carambola* L. Root. *Molecules* 17, 12330–12340. <https://doi.org/10.3390/molecules171012330>
- Qiu, S., Bai, M., Zhao, P., Liu, Z.X, Huang, X.X, Song, S.J., 2021. Phytochemical and network-based chemotaxonomic study of *Lonicera japonica* Thunb. *Biochem. Syst. Ecol.* 94, 104210. <https://doi.org/10.1016/j.bse.2020.104210>
- Quan, L.Q, Hegazy, A.M., Zhang, Z.J., Zhao, X.D., Li, H.M., Li, R.T., 2020. Iridoids and bis-iridoids from *Valeriana jatamansi* and their cytotoxicity against human glioma stem cells. *Phytochemistry* 175, 112372. <https://doi.org/10.1016/j.phytochem.2020.112372>
- Quezel et Santa., 1963. Nouvelle flore d'Algérie et des régions désertiques méridionales, Tome 2 C.N.R.S. Paris.
- Rahmouni, N., Pinto, D.C.G.A., Beghidja, N., Benayache, S., Silva, A. M.S. 2018. *Scabiosa stellata* L. phenolic content clarifies its antioxidant activity. *Molecules* 23, 1285. <https://doi.org/10.3390/molecules23061285>
- Rani, P.U., P., Devanan. P., 2013. Bioactivities of caffeic acid methyl ester (methyl-(*E*)-3-(3,4-dihydroxyphenyl) prop-2-enoate): a hydroxycinnamic acid derivative from *Solanum melongena* L. fruits. *J. Pest Sci.* 86, 579–589 <https://doi.org/10.1007/s10340-013-0516-8>
- Re, R., Pellegrini, N., Proteggente, A., Pannala, A., Yang, M., Rice-Evans, C., 1999. Antioxidant activity applying an improved ABTS radical cation decolorization assay. *Free Radical Biology and Medicine* 26, 1231–1237. [https://doi.org/10.1016/s0891-5849\(98\)00315-3](https://doi.org/10.1016/s0891-5849(98)00315-3)

- Rigat, M., Bonet, M.À.; Garcia, S.; Garnatje, T.; Vallès, J.,2007. Studies on pharmaceutical ethnobotany in the high river Ter valley (Pyrenees, Catalonia, Iberian Peninsula). J. Ethnopharmacol. 113, 267–277. <https://doi.org/10.1016/j.jep.2007.06.004>
- Sarikahya,N.B., Kirmizigul,S., 2010. Antimicrobial triterpenoid glycosides from *Cephalaria scoparia*.J. Nat. Prod.73, 825–830. <https://doi.org/10.1021/np900724u>
- Sarikahya, N.B., Murat, P., Arda, N., Kayce, P., Yavasoglu, N.U.K,Kirmizigul,S., 2011. Isolation and characterization of biologically active glycosides from endemic *Cephalaria* Species in Anatolia. *Phytochemistry* 4, 415-420. <https://doi.org/10.1016/j.phytol.2011.05.006>
- Sarikahya, N.B., Goren, A.C., Okkali,G.S., Kirmizigul,S., 2021. Saponins from twenty-two *Cephalaria* Species. *Rec. Nat. Prod.* 15, 537-546. <http://doi.org/10.25135/rnp.241.21.02.1985>
- Shi, A-H., Huang, J-W., Liu, Y-H., Yuan, K., 2013. Separation, antioxidant and antimicrobial activities of chemical constituents from exocarp of *Juglans mandshurica* Maxim. *Asian J. Chem.* 25, 3361-3365. <http://dx.doi.org/10.14233/ajchem.2013.13725>
- Singleton, V.L., Rossi. J.A., 1965. Colorimetry of total phenolics with phosphomolybdic-phosphotungsticacid reagents. *Am J Enol Vitic*16, 144-158.
- Soeren, R.J., Erik, L.P., Bent, J.N., 1979. Novelbis-iridoidglucosides from *Dipsacus sylvestris*. *Phytochemistry* 18, 273. [https://doi.org/10.1016/0031-9422\(79\)80069-2](https://doi.org/10.1016/0031-9422(79)80069-2)
- Tabatadze,N., Elias,R., Faure, R., Gerkens,P., De Pauw-Gillet, M.C., Kemertelidze, E., Chea, A.E., Ollivier, E.,2007. Cytotoxic triterpenoidsaponins from the roots of *Cephalaria gigantea*, *Chem. Pharm. Bull.* 55, 102–105. <https://doi.org/10.1248/cpb.55.102>
- Tang, Y.P., Liu, X., Yu, B., 2002. Iridoids from the rhizomes and roots of *Valeriana jatamansi*. *J. Nat. Prod.* 65 1949–1952. <https://doi.org/10.1021/np0203335>

- Tian, X.Y., Wang, Y.H., Yu, S.S., Fang, W.S., 2006. Two novel tetrairidoid glucosides from *Dipsacus asper*. *OrgLett.* 8, 2179–2182. <https://doi.org/10.1021/OL060676K>
- Tomassini, L., Foddai, S., Serafini, M., Cometa, M.F., 2000. Bis-iridoid glucosides from *Abelia chinensis*. *J. Nat. Prod.* 63, 998–999. <https://doi.org/10.1021/np9904713>
- Topçu, G, Ay, A, Bilici, A, Sarıkürkcü, C, Öztürk, M, Ulubelen, A. 2007. A new flavone from antioxidant extracts of *Pistacia terebinthus*. *Food Chem.* 103, 816–822. <https://doi.org/10.1016/j.foodchem.2006.09.028>
- Tsukamoto, H., Hisada, S., Nishibe, S., 1984a. Lignans from bark of *Fraxinus mandshurica* var. *japonica* and *F. japonica*. *Chem. Pharm. Bull.* 32, 4482–4489. <https://doi.org/10.1248/cpb.32.4482>
- Tsukamoto, H., Hisada, S. Nishibe, S., 1984b. Lignans from bark of the *Olea* plants. I. *Chem. Pharm. Bull.* 32, 2730–2735. <https://doi.org/10.1248/cpb.32.2730>
- Tsukamoto, S. Hisada, S. Nishibe, S., 1985. Lignans from bark of *Olea* plants. II. *Chem. Pharm. Bull.* 33, 1232–1241. <https://doi.org/10.1248/cpb.33.1232>
- Wang, J., Liu, K., Xu, D., Wang, Q., Bi, K., Song, Y., Zhang, L., 2013. Rapid micropropagation system *in vitro* and antioxidant activity of *Scabiosa tschiliensis* Grunning. *Plant Growth Regulation* 69, 305–310. <https://doi.org/10.1007/s10725-012-9765-4>
- Wang, J.Y., Zhao, Z.L., Xue, P.F., Ma, F.X., Zhang, D.Y., Wang, N.N., Li, M.H., 2015. Chemical constituents from flowers of *Scabiosa tschiliensis*. *China Journal of Chinese Materia Medica* 40, 807–813. [https://doi.org/10.1016/s0926-2040\(97\)00018-0](https://doi.org/10.1016/s0926-2040(97)00018-0)
- Wawer, I., Zielinska, A., 2001. ¹³C CP/MAS NMR studies of flavonoids. *Magn. Reson. Chem.* 39, 374–380. <https://doi.org/10.1016/j.phytochem.2009.03.018>
- Wu, M., Wu, P., Liu, M., Xie, H., Jiang, Y., Wei, X., 2009. Iridoids from *Gentianaloureirii*. *Phytochemistry* 70, 746–750. <https://doi.org/10.1016/j.phytochem.2009.03.018>

- Xu, Z., Chang, L., 2017. Caprifoliaceae, In: Identification and control of common weeds: Volume 3, Pages 405-416. Singapore: Springer <https://doi.org/10.1007/978-981-10-5403-7>
- Yang, W., Jiansong Sun, J., Lu, W., Li, Y., Shan, L., Han, W., Zhang, W. D., Yu, B., 2010. Synthesis of kaempferol 3-O-(3'',6''-di-O-E-p-coumaroyl)- β -D-glucopyranoside, efficient glycosylation of flavonol 3-OH with glycosyl O-alkynylbenzoates as donors. *J. Org. Chem.* 75, 6879–6888. <https://doi.org/10.1021/jo1014189>
- Yu, L., Haley, S., Perret, J., Harris, M., Wilson, J., Qian, M. 2002. Free radical scavenging properties of wheat extracts. *J Agric Food Chem.* 50, 1619–1624. <http://dx.doi.org/10.1021/jf010964p>
- Zheng, Q., Koike, K., Han, L. K., Okuda, H., Nikaido, T., 2004. New biologically active triterpenoid saponins from *Scabiosa tschiliensis*. *J. Nat. Prod.* 67, 604–613. <https://doi.org/10.1021/np0304722>