



HAL
open science

Editorial: The Use of Plant Extracts and Essential Oils as Biopesticides

Rachid Lahlali, Hajar El Hamss, Jouda Mediouni-Ben Jemâa, Essaid Ait
Barka

► **To cite this version:**

Rachid Lahlali, Hajar El Hamss, Jouda Mediouni-Ben Jemâa, Essaid Ait Barka. Editorial: The Use of Plant Extracts and Essential Oils as Biopesticides. *Frontiers in Agronomy*, 2022, 4, 10.3389/fagro.2022.921965 . hal-03920695

HAL Id: hal-03920695

<https://hal.univ-reims.fr/hal-03920695>

Submitted on 3 Jan 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License



Editorial: The Use of Plant Extracts and Essential Oils as Biopesticides

Rachid Lahlali^{1*}, Hajar El Hamss¹, Jouda Mediouni-Ben Jemâa² and Essaid Ait Barka^{3*}

¹ Department of Plant Protection, Phytopathology Unit, Ecole Nationale d'Agriculture de Meknès, Meknès, Morocco, ² Institut National de la Recherche Agronomique de Tunisie (INRAT), Ariana, Tunisia, ³ Unité de Recherche Résistance Induite et Bio-Protection des Plantes-EA 4707-USC INRAE1488, Reims Champagne-Ardenne University, Reims, France

Keywords: plant pathogens, pests, plant extracts, essential oils, biopesticides

Editorial on the Research Topic

The Use of Plant Extracts and Essential Oils as Biopesticides

Essential oils (EOs) and plant extracts contain valuable natural products, many of which can be used in pest and disease control safely due to their ability to degrade in nature (Ni et al., 2021; El Khetabi et al., 2022). Despite their qualities, biopesticides represent only 5% of the overall pesticide market (Balog et al., 2017; Kumar et al., 2021; Rakshit et al., 2021). Nevertheless, biopesticides are experiencing rapid growth in recent years with an average annual growth rate of 9–20%, predicted to outpace that of chemical pesticides (Balog et al., 2017; Marrone, 2019; Kumar et al., 2021; Rakshit et al., 2021). Common management practices focus on the application of EOs and plant extract-based biopesticides without deeply understanding their mode of action (Álvarez-Martínez et al., 2021). Recent research focusing on pest management using EOs and plant extracts revealed several mechanisms involved in the insecticidal effects of EOs on targeted organisms (Ni et al., 2021). Moreover, several reports suggest specific strategies that could help in optimizing application of EOs and plant extracts as part of integrated pest management programs (de Oliveira, 2021). Apart from their role in plant development and growth, plant secondary compounds are also essential in plant resistance to biotic and abiotic stressors, and can be involved in metabolic processes that control plant tolerance (Yang et al., 2018; Karimi and Meiners, 2021; Ni et al., 2021). However, EOs and plant extracts are biologically unstable as they are easily destroyed by environmental pH, oxygen, light, and moderate temperatures. EOs exhibit poor aqueous solubility and high volatility in general. Efforts are being made to overcome these challenges. The present Research Topic gathers together studies that focus on the use of plant extracts and essential oils as biopesticides while aiming at the same time to shed light on their mode of action on different targeted key agricultural pest and plant diseases including insects, mites, nematodes, and oomycetes.

Several bioassays of different EOs and plant extracts have been used to demonstrate potential control of different key insect and mite populations. In one study, the strong ixodicidal and antifeedant agents included EOs of *Thymus zygis*, *T. vulgaris*, and *Mentha suaveolens*, which were suggested to be developed as biopesticides to effectively control ticks and insect pests (Valcárcel et al.). Interestingly, synergistic effects were also observed between these EOs (Valcárcel et al.). In a large-scale field trial, the application of aqueous extracts of *Murraya paniculata*, *Cassia tora*, *Amphineuron opulentum*, *Tithonia diversifolia*, and *C. alata* equally reduced the population of the red spider mite, a major tea pest, with a lower impact on natural enemies and increased the yield of tea plants without lethal consequence for the tea plants or consumers (Deka et al.).

OPEN ACCESS

Edited and reviewed by:

Murray B. Isman,
University of British
Columbia, Canada

*Correspondence:

Rachid Lahlali
rlahlali@enameknes.ac.ma
Essaid Ait Barka
ea.barka@univ-reims.fr

Specialty section:

This article was submitted to
Pest Management,
a section of the journal
Frontiers in Agronomy

Received: 16 April 2022

Accepted: 02 May 2022

Published: 13 June 2022

Citation:

Lahlali R, El Hamss H, Mediouni-Ben
Jemâa J and Barka EA (2022)
Editorial: The Use of Plant Extracts
and Essential Oils as Biopesticides.
Front. Agron. 4:921965.
doi: 10.3389/fagro.2022.921965

Furthermore, the larvicidal activity of EOs from five piper species including *Piper aduncum*, *P. marginatum*, *P. gaudichaudianum*, *P. crassinervium*, and *P. arboreum* against the yellow fever mosquito *Aedes aegypti* showed up to a 90% lethality at a screening concentration of 100 ppm, making these EOs potential alternatives for control of *A. aegypti* mosquito larvae (Pereira Filho et al.). Despite these effects, EO-based products are often chemically variable, requiring rigorous control of the cultivation process and an understanding of the regulatory aspects of the biosynthesis of these phenylpropanoids (Pereira Filho et al.), especially in developing countries such as sub-Saharan Africa (SSA) (Uyi et al.). The latter review summarized the existing insecticidal activity of invasive plants in SSA including Asteraceae, Solanaceae, Fabaceae, and Euphorbiaceae amongst others. These plants caused 50–100% mortality against various insect pests (Uyi et al.). However, using extracts from these invasive plants as biopesticides in African countries, especially among resource-poor smallholder farmers and locals, remains challenging (Stevenson et al., 2017; Uyi et al.).

Unlike studies on insects and mites, very few bioactivity evaluation bioassays with different EOs and plant extracts have been carried out on controlling different nematode populations. Recently the nematocidal potential of *Citrus sinensis*, *Cymbopogon nardus*, and *Melaleuca alternifolia* has been reported against the cotton root-knot nematode *Meloidogyne incognita* (Kundu et al.). On the other hand, potential botanical products against plant diseases have been extensively studied. For example, volatile organic compounds (VOCs) from soybean plants have been tested against harmful oomycetes, represented by *Phytophthora* (Ge et al.). The VOCs, mainly containing 4-ethylphenol, were simultaneously inoculated with the causal agent of soybean root rot, *P. sojae* and the black shank, *P. nicotiana*. VOCs inhibited the growth of the pathogens by destroying their cell membrane (Ge et al.). These VOCs have potent antifungal activity against other soil-borne phytopathogenic fungi including *Rhizoctonia solani*, *Fusarium graminearum*, and *Gaeumannomyces graminis* var *tritici*, and four forma specialis of *Fusarium oxysporum* (Ge et al.), making it ideal for simultaneously controlling major soilborne diseases.

The identification of active compounds of EOs and plant extracts is crucial to study their mode of action and thus develop effective biopesticide products. Nanotechnology using active biomolecules represents a potential solution to control the release of the active ingredient with less product waste. Biochemical and molecular modes of action of plant extracts and EOs have been recently investigated. *Thymus proximus* EO contains carvacrol, *p*-cymene, and γ -terpinene, representing 85.9% of the total oil, and these major constituents are responsible for both the plant suppressive effect and the insecticidal activity of the EOs (Zhou et al.). Furthermore, Azadirachtin, a tetranortriterpenoid derived from the neem seed of the Indian neem tree, have been reviewed many, many times in the past going back to the early 1980s for its insecticidal activity (Schmutterer, 2002). In this Research Topic, the literature review of Kilani-Morakchi et al. summarized the state of the art on key azadirachtin insecticidal activities and risk assessment. The effect of *T.*

zygis, *T. vulgaris*, and *M. suaveolens* was mainly due to the presence of active compounds including piperitenone oxide, carvacrol, piperitenone, and thymol (Valcárcel et al.). The major compounds of the EOs from *Piper* species were identified and included β -asarone, (E)-anethole, (E)- β -caryophyllene, γ -terpinene, *p*-cymene, limonene, α -pinene, and β -pinene and showed larvicidal activity with mortality between 90 and 100% (Pereira Filho et al.). Phytochemical analyses of EOs of *Origanum compactum* tested against *Callosobruchus maculatus* showed that the main components were carvacrol and thymol (38 and 31.5%, respectively) (Aimad et al.). A comprehensive chemoprofiling of nematocidal action of EOs was performed to understand their possible interactions with the target sites of *M. incognita*, suggesting the most prominent monoterpene was l-limonene, with a range of 32–98%. In particular, industrially important *Commiphora myrrha*, *Cymbopogon nardus*, *Artemisia absinthium*, and *Pogostemon cablin* contained a higher amount of furanoeudesm 1,3 diene, geraniol, myrcene, camphor, and patchoulol, respectively. *In silico* analysis suggested a higher binding capacity of geraniol, β -terpineol, citronellal, l-limonene, and γ -terpinene, to the selected target proteins (Kundu et al.). Terpenoids which are present in most essential oils have been reported responsible for their bioactivity (Kundu et al.). Interestingly, several studies pointed out that synergistic interactions among terpenoids in EOs can be important (Tak and Isman, 2015, 2017). Therefore, these biochemical analyses of EOs and plant extracts will open a new door to specifically devise efficient and eco-friendly biopesticides and will help in effectively targeting the plant system (Werrie et al.).

The present Research Topic provides important updates on the roles of EOs and plant extracts in pest and disease management and highlights the chemical compositions responsible for their mode of action. This Research Topic contributes to define potential EO and plant extract candidates, which can be implemented in eco-friendly and sustainable management strategies. Moreover, plants are an important source of biomolecules, which are essential for fighting against economically devastating pests and diseases, and are also listed in this Research Topic. As such, it contributes to advancing the development of sustainable strategies for pest and disease management of food crops. However, there is a big disconnect between academic studies on insecticidal activity of plant metabolites and production of commercial bioinsecticides. Many such compounds are touted as potential biopesticides, but very few meet all the necessary criteria to be produced at scale for commercial use (Isman, 2017, 2020). Finally, besides providing an update on the state of the art of biopesticide research, the current Research Topic offers a perspective on future research needs and priorities. Emerging areas of research related to biopesticides include investigating (i) biopesticides' roles and function in plant metabolism and (ii) novel biopesticide management strategies to address biopesticide waste such as that of nanotechnology. Biopesticides and related products should be evaluated in a more biological and ecological context to further enhance the penetration of biopesticides into plant tissues, thus decreasing the waste and degradation of biopesticides

and contributing to more sustainable integrated pest management systems.

AUTHOR CONTRIBUTIONS

All authors listed have made a substantial, direct, and intellectual contribution to the work and approved it for publication.

REFERENCES

- Álvarez-Martínez, F. J., Barrajón-Catalán, E., Herranz-López, M., and Micol, V. (2021). Antibacterial plant compounds, extracts and essential oils: an updated review on their effects and putative mechanisms of action. *Phytomedicine* 90, 153626. doi: 10.1016/j.phymed.2021.153626
- Balog, A., Hartel, T., Loxdale, H. D., and Wilson, K. (2017). Differences in the progress of the biopesticide revolution between the EU and other major crop-growing regions. *Pest Manag. Sci.* 73, 2203–2208. doi: 10.1002/ps.4596
- de Oliveira, J. L. (2021). “1 - nano-biopesticides: present concepts and future perspectives in integrated pest management,” in *Advances in Nano-Fertilizers and Nano-Pesticides in Agriculture*, eds S. Jogaiah, H. B. Singh, L. F. Fraceto, and R. d. Lima (Cambridge, MA: Woodhead Publishing), 1–27.
- El Khetabi, A., Lahlali, R., Ezrari, S., Radouane, N., Nadia, L., Banani, H., et al. (2022). Role of plant extracts and essential oils in fighting against postharvest fruit pathogens and extending fruit shelf life: a review. *Trends Food Sci. Technol.* 120, 402–417. doi: 10.1016/j.tifs.2022.01.009
- Isman, M. B. (2017). Bridging the gap: moving botanical insecticides from the laboratory to the farm. *Indus. Crops Products* 110, 10–14. doi: 10.1016/j.indcrop.2017.07.012
- Isman, M. B. (2020). Botanical insecticides in the twenty-first century—fulfilling their promise? *Annu. Rev. Entomol.* 65, 233–249. doi: 10.1146/annurev-ento-011019-025010
- Karimi, A., and Meiners, T. (2021). Antifungal activity of *Zataria multiflora* Boiss. essential oils and changes in volatile compound composition under abiotic stress conditions. *Indus. Crops Products* 171, 113888. doi: 10.1016/j.indcrop.2021.113888
- Kumar, J., Ramlal, A., Mallick, D., and Mishra, V. (2021). An overview of some biopesticides and their importance in plant protection for commercial acceptance. *Plants* 10, 1185. doi: 10.3390/plants10061185
- Marrone, P. G. (2019). Pesticidal natural products - status and future potential. *Pest Manag. Sci.* 75, 2325–2340. doi: 10.1002/ps.5433
- Ni, Z.-J., Wang, X., Shen, Y., Thakur, K., Han, J., Zhang, J.-G., et al. (2021). Recent updates on the chemistry, bioactivities, mode of action, and industrial applications of plant essential oils. *Trends Food Sci. Technol.* 110, 78–89. doi: 10.1016/j.tifs.2021.01.070

FUNDING

This study was financially supported by the Phytopathology Unit, Ecole Nationale d'Agriculture de Meknès and Unité de Recherche Résistance Induite et Bio-Protection des Plantes, Reims Champagne-Ardenne University.

- Rakshit, A., Meena, V. S., Abhilash, P., Sarma, B., Singh, H., Fraceto, L., et al. (2021). *Biopesticides: Volume 2: Advances in Bio-Inoculants*. Cambridge, MA: Woodhead Publishing.
- Schmutterer, H. (2002). *The Neem Tree, 2nd Edition*. Mumbai: The Neem Foundation.
- Stevenson, P. C., Isman, M. B., and Belmain, S. R. (2017). Pesticidal plants in Africa: a global vision of new biological control products from local uses. *Ind. Crops Prod.* 110, 2–9. doi: 10.1016/j.indcrop.2017.08.034
- Tak, J.-H., and Isman, M. (2015). Enhanced cuticular penetration as the mechanism for synergy of insecticidal constituents of rosemary essential oil in *Trichoplusia ni*. *Sci. Rep.* 5, 12690. doi: 10.1038/srep12690
- Tak, J.-H., and Isman, M. B. (2017). Penetration-enhancement underlies synergy of plant essential oil terpenoids as insecticides in the cabbage looper, *Trichoplusia ni*. *Sci. Rep.* 7, 42432. doi: 10.1038/srep42432
- Yang, L., Wen, K.-S., Ruan, X., Zhao, Y.-X., Wei, F., and Wang, Q. (2018). Response of plant secondary metabolites to environmental factors. *Molecules* 23, 762. doi: 10.3390/molecules23040762

Conflict of Interest: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Publisher's Note: All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article, or claim that may be made by its manufacturer, is not guaranteed or endorsed by the publisher.

Copyright © 2022 Lahlali, El Hamss, Mediouni-Ben Jemâa and Barka. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.