



Traditional Chinese medicine: saponins, critical micellar concentrations and partition coefficients

Cédric Barthel, Georges Massiot, Catherine Lavaud

► To cite this version:

Cédric Barthel, Georges Massiot, Catherine Lavaud. Traditional Chinese medicine: saponins, critical micellar concentrations and partition coefficients. *Phytochemical Analysis*, 2023, 34, pp.414. 10.1002/pca.3221 . hal-04040462

HAL Id: hal-04040462

<https://hal.univ-reims.fr/hal-04040462>

Submitted on 22 Mar 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License

RESEARCH ARTICLE

Traditional Chinese medicine: saponins, critical micellar concentrations and partition coefficients

Cédric Barthel¹ | Georges Massiot¹  | Catherine Lavaud² 

¹Institut de Chimie Moléculaire de Reims,
UMR CNRS 7312, Université
Reims-Champagne-Ardenne, UFR Sciences,
Reims, France
²Institut de Chimie Moléculaire de Reims,
UMR CNRS 7312, Université
Reims-Champagne-Ardenne, UFR Pharmacie,
Laboratoire de Pharmacognosie, Reims, France

Correspondence

Georges Massiot, Institut de Chimie
Moléculaire de Reims, UMR CNRS 7312,
Université Reims-Champagne-Ardenne, UFR
Sciences, BP 1039, 51 687 Reims Cedex 2,
France.
Email: georges.massiot@univ-reims.fr

Funding information

Agence Régionale de Santé (ARS) Grand Est

Abstract

Introduction: Traditional Chinese medicine (TCM) revolves around complex mixtures bound to specific roles within the formulation, among which saponin-containing plants with alleged properties of harmonising or detoxifying other compounds present in the preparations.

Objective: This article deals with the study of these interactions with, as a model, the interaction between saponins and selected active principles.

Methods: The measurement of the partition coefficient between water and octanol ($\log P$) was used as an indicator and determined by nuclear magnetic resonance (NMR) for these active principles in the presence of saponins. For each compound, a graph was constructed showing the evolution of $\log P$ with increasing concentrations of saponins.

Results: Four distinct patterns of interactions were distinguished. Pattern A showed a constant decrease of $\log P$, pattern B showed a decrease followed by a plateau, in pattern C the $\log P$ did not vary until the critical micellar concentration (CMC) and decreased afterwards, and pattern D exhibited an increase of $\log P$. These properties were linked to the ability of saponins to form micelles in water once the CMC is reached. The interaction of aconitine and saponins followed pattern D, thus explaining the detoxification of herbal preparations using *Aconitum* with licorice. The licorice facilitated the extraction of the notoriously water-insoluble artemisinin from *Artemisia annua*.

Conclusion: This investigation confirms that the physical properties of micelle forming saponins are intimately linked to a modification of behaviour of the other molecules in solution, as seen with the alteration of $\log P$ and the four types of interactions presented.

KEYWORDS

aconitine, artemisinin, critical micellar concentration, licorice, partition coefficient, saponins, traditional Chinese medicine

The Agence Régionale de Santé (ARS) Grand Est and the Centre Hospitalier Universitaire (CHU) of Reims (France) are acknowledged for providing a thesis grant to Cédric Barthel. Dedicated to the memory of Professor Louissette Le Men-Olivier, who initiated work on saponins in Reims

This is an open access article under the terms of the [Creative Commons Attribution](https://creativecommons.org/licenses/by/4.0/) License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

© 2023 The Authors. *Phytochemical Analysis* published by John Wiley & Sons Ltd.

1 | INTRODUCTION

Chinese herbal medicines generally consist of mixtures of a fairly large number of plants, with each of them being assigned a specific role: Sovereign (or Emperor), Minister, Assistant (or Adjuvant) and Courier (or Messenger), also known as the Jun-Chen-Zuo-Shi classification.^{1,2} Numerous attempts have been made to extract “heroic principles” from this corpus of plants to include them in the Western pharmacopeia, with little success, a noticeable exception being artemisinin (Qing Hao Su) extracted from *Artemisia annua* L.³ The main reason for this relative failure lies in a fundamental difference between the two systems of medicine: “one disease-one target-one drug” on the one hand, “target networks-multiple drugs” on the other hand.^{4,5} The occidental system revolves around curing a target disease, while traditional Chinese medicine (TCM) aims at restoring balance. Human genome sequencing and a better understanding of genetic interactions have led to a renewal of interest in combination therapy, which will hopefully help to fill the gap between the two systems.⁶ Activity against one or several molecular targets might be key to the success of these preparations and the chemical and physical properties of these complex mixtures also contribute to it. The complexity of TCM preparations makes them difficult to study at the molecular level and we simplified the problem by taking as a model, the interactions between saponins and diverse active principles.

The presence of saponin-rich plants in a large variety of preparations is a characteristic of TCM. Table 1 lists some of them (not exhaustively) as well as their role in traditional formulations.^{7,8} It should be noticed that, while they can occupy all the four positions of the Jun-Chen-Zuo-Shi classification, saponin-rich plants are most often found in the “Courier” and “Assistant” roles. For instance, the root of *Glycyrrhiza glabra* L., notoriously known to contain saponins, fills the role of Sovereign in the Zhi Gan Cao Tang and Gancao Xiaomai Dazao Tang formulations, while being the Assistant in Xiao Jian Zhong Tang or the Courier in many others (Si Ni Tang, Si Jun Zi Tang, Sang Ju Yin, Xia Qing Long Tang, Shao Yao Gan Cao Tang, Gan Cao Ganjiang Fuling Baizhu Tang, Guizhi Gan Cao Longgu Muli Tang, Zhizi Gan Cao Chi Tang, Houpo Shengjiang Banxia Gan Cao Renshen Tang preparations).

Most authors explain the ubiquitous presence of saponins in TCM by their intrinsic biological properties, but we felt that, in

addition to those, the solubilising properties of saponins must also play a key role. The purpose of this work is to investigate the effect of saponins on the bioavailability of drugs and we chose to work on a few saponins representative of plants used in TCM: glycyrrhizin (*Glycyrrhiza glabra*), α -hederin (*Hedera helix* and *Nigella sativa*), β -aescin (*Aesculus hippocastanum*), soyasaponin I (*Glycine max* and *Medicago sativa*), as pure compounds and an extract from *Quillaja saponaria*.

2 | EXPERIMENTAL

2.1 | General experimental procedures

All tested compounds were available from our laboratory or purchased from Sigma-Aldrich or Alfa-Aesar. Millipore water was used and purified 1-octanol was purchased from Prolabo.

2.2 | Sample preparation

Sample preparation required c. 2 mg of pure compound which was solubilised in a 5 ml vial with 1.5 ml of a water/deuterated water (H_2O/D_2O) (90:10) saponin solution at different concentrations. Stock solutions of saponins were prepared in H_2O/D_2O (90:10) at 2 g/L for glycyrrhizin, 0.03 g/L for α -hederin, and 0.3 g/L for soyasaponin I, β -aescin, and *Quillaja* saponin. A range of dilutions around the critical micellar concentration (CMC) was prepared from these solutions: 0.02 to 1.28 g/L for glycyrrhizin, 0.002 to 0.03 g/L for α -hederin, and 0.02 to 0.3 g/L for soyasaponin I, β -aescin, and *Quillaja* saponin. Solutions were placed in an ultrasonic bath for 5 to 10 min to optimise solubilisation and filtered on a high-performance liquid chromatography (HPLC) disposable syringe filter from Macherey-Nagel (pore size 0.45 μm). An aliquot (550 μl) of the filtered solution was placed in nuclear magnetic resonance (NMR) tube A for analysis, while 700 μl was placed in an Eppendorf tube for partition against 700 μl of 1-octanol. The Eppendorf tube was inverted at least 40 times, centrifuged at around 2000 rpm for 10 min and left to rest for 24 h to reach equilibrium. A portion (550 μl) from the aqueous phase was then placed into NMR tube B for comparative analysis.

Plant species	Role	Plant species	Role
<i>Achyranthes bidentata</i> Blume	Minister	<i>Dioscorea polystachya</i> Turcz.	Assistant
<i>Acorus tatarinowii</i> Sol.	Courier	<i>Foeniculum vulgare</i> Mill.	Courier
<i>Adenophora tetraphylla</i> A. DC.	Minister	<i>Forsythia suspensa</i> Vahl	Courier
<i>Aesculus hippocastanum</i> L.	Sovereign	<i>Gardenia jasminoides</i> J. Ellis	Courier
<i>Allium schoenoprasum</i> L.	Sovereign	<i>Glycyrrhiza glabra</i> L.	Courier
<i>Amomum villosum</i> Lour.	Assistant	<i>Indigo tinctoria</i> L.	Courier
<i>Angelica pubescens</i> Maxim.	Assistant	<i>Lonicera japonica</i> Thunb.	Courier
<i>Atractylodes lancea</i> Thunb.	Assistant	<i>Platycodon grandiflorus</i> A. DC.	Courier
<i>Cimicifuga heracleifolia</i> Kom.	Courier	<i>Polygala tenuifolia</i> Willd.	Assistant
<i>Codonopsis pilosula</i> Franch.	Assistant		

TABLE 1 A list of plant containing saponins used in traditional Chinese medicine (TCM) and their place in the Jun-Chen-Zuo-Shi classification.

2.3 | NMR procedures

NMR measurements were performed on a Bruker Avance III 600 MHz spectrometer using thin-wall NMR tubes of size 5 mm × 178 mm and equipped with a TCI (triple-resonance inverse with carbon observe) cryoprobe. Peak picking, integration and quantitative NMR were realised on the Topspin software (version 4.0.9) from Bruker. The temperature of the samples was regulated at 298 K. To suppress the solvent, the spectrometer frequency was locked onto D₂O and proton (¹H) spectra were acquired by means of the zgpg30 pulse sequence with eight scans per free induction decay (FID) and a 12 s relaxation delay, resulting in a 4 min 55 s recording time. The transmitter frequency offset (O1) was set at water frequency (around 2818 Hz) and gradient ratios (smoothed rectangular shape sequence SMSQ10.100) were fixed to gradients G1 = 31% and G2 = 11%. A 180° shaped pulse (square selective pulse Squa100.1000) was fixed at 180° pulse length p12 = 2 ms and power at 180° shaped pulse SPdB (−dB) = 33 or SPW (W) = 5.01 e^{−4}. The FIDs (64k points, spectral width = 12,000 Hz) were processed with LB = 0.3 Hz. Concentrations in sample NMR tubes A and B were measured with the ERETIC 2 method integrated in the Topspin software from Bruker. For more practical details see Barthel *et al.*⁹

2.4 | Microscopic experiments

Microscopic observations were realised on an inverted Olympus IX70 microscope with a ×100 immersion lens. Aqueous sample solutions were prepared in glass vials at 1.00 g/L (2 mg of sample in 2 ml H₂O) and a drop was deposited on the glass slide. Increasing glycyrrhizin concentrations were added to the sample solutions. Resulting pictures were analysed on the ImageJ software.

2.5 | HPLC analysis

HPLC measurements were performed with an UltiMate 3000 system equipped with Chromeleon software. The column was a C18 250 mm × 4.6 mm with 5 µm particle size from Interchim, and the mobile phase consisted of acetonitrile/H₂O (70:30, v/v) with 0.10% formic acid. The flow rate was 1 ml/min, the injection volume was 10 µl, and the column temperature was 30°C; elution duration was 20 min. Detection was made on a ultraviolet (UV) detection diode at a wavelength of 254 nm. For artemisinin, the analysis was made on a lyophilised aqueous extract of *Artemia annua* leaves (courtesy of Pierre Fabre Laboratories) which was prepared from 4 g of dried powdered leaves with a 15 min decoction in 200 ml of boiling pure water. Five similar decoctions were prepared by adding glycyrrhizin (0.04, 0.08, 0.16, 0.32, 0.64 g/L) to the solution. After filtration, the aqueous solutions were lyophilised, and 5 mg of residue were dissolved in 2 ml of HPLC mobile phase. The retention time of artemisinin was detected at 6 min.

3 | RESULTS AND DISCUSSION

3.1 | Measurement of critical micellar concentrations and partition coefficients

Saponins are tensioactive molecules containing a hydrophobic lipidic core (steroid or triterpenoid) coupled with one or several hydrophilic sugar chains. Above a certain concentration, called the CMC, saponins tend to organise themselves into micelles of various sizes and forms.¹⁰ CMCs in water, were measured on our saponin samples, alone and in the presence of a selection of guest molecules featuring the active constituents of drugs (Table 2 and Figure S39). Measurements were carried out with a du Noüy ring tensiometer¹¹ and the resulting values were in fair agreement with literature values obtained on a plate method tensiometer.^{12–15} The presence of guest molecules had little or no effect on the CMC.

According to principles of drug development in the pharmaceutical industry, the bioavailability of drugs is deemed to be linked to partition coefficients between octanol and water (or logP).¹⁶ The partition coefficients are usually measured by HPLC. However, for the sake of convenience and due to the very large number of measurements involved, we developed a quicker method based on NMR.⁹ This method is sensitive and allowed a large number of analysis since only about 2 mg of compound and a small quantity of D₂O are required (150 µl). To avoid effects due to D₂O as solvent, we chose to use a mixture of H₂O/D₂O (90:10), that allowed valid comparison with reported logP measured in H₂O in the literature. The suppression of the residual water peak was performed using water suppression by excitation sculpting (a spin-echo experiment); the dephasing of all coherences set on the water frequency, cleanly and selectively with limiting baseline distortions and phasing issues (about 0.5 ppm around water signal). The ERETIC 2 tool allows a direct quantification of NMR signals with no additional standard. The full list of bioactive molecules representing a selection of drugs from natural or chemical origin is provided in the Supporting Information section (Table S1).

3.2 | Influence of the concentration of glycyrrhizin on partition coefficients

A stock solution of glycyrrhizin (ammonium salt) was made in a mixture of H₂O/D₂O (9:1), dilutions of which were prepared to proceed to measurements below and above its CMC. These dilutions were utilised to prepare sample aqueous solutions and the concentration was determined by ¹H NMR using the ERETIC 2 tool.⁹ A part of the solution was partitioned with 1-octanol and the concentration of the aqueous phase was determined in the same manner (Figure 1). The partition coefficient of the sample was calculated from the ratio of these two values with C_w as the concentration in water and C_{wo} that after partition.

$$\log P = \log \frac{C(\text{octanol})}{C(\text{water})} = \log \frac{C_w - C_{wo}}{C_{wo}}$$

Saponin	Guest molecule	CMC (g/L)	Literature value (g/L)
Glycyrrhizin	-	0.15	0.20
	(+)-Catechin	0.20	
	Caffeine	0.30	
	Acetylsalicylic acid	0.20	
	Ascorbic acid	0.15	
	L-Phenylalanin	0.15	
	L-Tyrosin	0.15	
	Vanillin	0.25	
α -Hederin	-	0.01	0.01
	(+)-Catechin	0.01	
	L-Tyrosin	0.01	
β -Aescin	-	0.10	0.10
	(+)-Catechin	0.06	
Soyasaponin I	-	0.16	0.09
	(+)-Catechin	0.10	
Quillaja saponin	-	0.05	0.14–0.77
	(+)-Catechin	0.06	
	Salicylic acid	0.06	
	Caffeine	0.05	

TABLE 2 Critical micellar concentration (CMC) of saponin samples with or without guest molecules.




FIGURE 1 Overview of experimental procedure.

Depending on the molecules under study (Table S1), four patterns of behaviour were observed. Pattern A showed a constant decrease of $\log P$, therefore a decrease of lipophilicity, as the saponin concentration increased, with no inflexion of the curve at the CMC. Pattern B was characterised by a similar decrease of $\log P$ until the CMC was reached, followed by a plateau. In pattern C, $\log P$ did not vary, until an inflexion was observed at the CMC, followed by a continuous decrease. In contrast with the preceding observations, pattern D showed an increase of lipophilicity (a rise in $\log P$) until the CMC was reached, followed by a plateau. Examples of representative curves are presented in Figure 2. Although the curves corresponding to patterns A and B bear similarities in their descending parts, they differ in the influence of the micelle formation concentration.

The evolution of $\log P$ of salicylic acid in the presence of increasing quantities of glycyrrhizin illustrated pattern A (Figure 2a). When the glycyrrhizin concentration was increased from 0 to 3 g/L, the curve showed a decrease of $\log P$ from 2.5 to 0.5, corresponding to a 100-fold increase in water solubility. This phenomenon may be explained by a decrease of the water superficial tension, which promotes the dispersion of salicylic acid and hence its solubility. Once the CMC was reached, it was observed that the solubility still increased, suggesting a further interaction between the saponin and the molecule, with its incorporation into the micelles. Similar curves were observed for a large variety of compounds: aromatic acids (acetylsalicylic, caffeic, gallic and *p*-coumaric acids), alkaloids (atropine, colchicine and quinine), lawsone, mangiferin, stevioside and rutin (see Supporting Information Figures S1–S11).




FIGURE 2 Partition coefficient of salicylic acid (a), (+)-catechin (b), codeine (c) and caffeine (d) as a function of glycyrrhizin concentrations.

(+)-Catechin was an example of a curve characterised by a sharp decrease of logP until the CMC was reached, followed by a plateau, where no significant evolution of logP was observed (Figure 2b). Therefore, it does not seem that micelle formation interferes with the solubilisation process. Once the CMC was reached, superficial tension remained stable and no further solubilisation was observed. Microscopic examination of catechin “solutions” showed the presence of insoluble aggregates, which disappeared upon addition of glycyrrhizin, hence demonstrating its solubilising properties (Figure S30). The same behaviour was observed with the four other saponins. Cinnamaldehyde, cinchonine, cocaine, phenylalanine, tyrosine, theobromine and theophylline gave rise to the same phenomenon (Figures S12–S18).

Molecules with pattern C gave sigmoidal curves with no or little slope until the CMC was reached, followed by a decrease. Thus, logP of codeine remained virtually unchanged until the CMC (0.05) and progressively dropped towards -0.25 at three times the CMC (Figure 2c). In this case, the increase in water solubility was directly linked to micelle formation and not to diminished surface tension. Microscopic examination showed that the aggregates corresponding

to insoluble material disappeared as soon as micelles were formed (Figure S31). This reaction was observed with morphine, acetaminophen, chlorogenic acid, arbutin, vanillin and zolmitriptan (Figures S19–S25).

A totally opposite behaviour was observed for some molecules, where lipophilicity increased with saponin concentration, because of water dissociation forces becoming weaker than intermolecular forces (Figure 2d). For instance, the couple caffeine/glycyrrhizin, showed an increase in logP, from -0.5 to -0.25 , reached at the CMC, and with no significant change above it. In this case, the guest molecules become less hydrophilic and more soluble in octanol, as also seen with naringin and capecitabine (Figures S26–S29).

Several compounds such as ascorbic acid did not fit in the picture: its logP decreased until the CMC was reached and then increased to a value superior to its initial value (Table S2). This behaviour reflects the ability of ascorbic acid to destabilise micelles and to act as a solvent.^{17,18} Molecules of too high or too low lipophilicity did not lend themselves to reliable logP measurements with our method and could not be incorporated into this model. They are β -sitosterol, β -carotene,

digitoxin, 18 β -glycyrrhetic acid, docetaxel, myrcene, thymol, amygdalin, curcumin, daidzein, emodin, esculetin, hesperidin, ergotamine and podophyllotoxin.

3.3 | Generalisation with regard to TCM

Salicylic acid was chosen as a type A model compound, due to its presence in several plants from the Chinese pharmacopeia: *Illicium verum* (star anise), *Peonia lactiflora* (common garden peony), *Isatis tinctoria* (dyer's woad) and *Nelumbo nucifera* (sacred lotus). Its partition coefficient was measured against different concentrations of α -hederin, soyasaponin I, a saponin mixture from *Q. saponaria* and β -aescin (see Figures S32–S35). In the first three cases, the curves showed a profile analogous to the one observed with glycyrrhizin (pattern A), but a pattern B curve was observed with β -aescin, with a plateau after the CMC. Thus, the addition of a saponin containing plant to a plant containing salicylic acid led to a three-fold decrease in lipophilicity of the active principle, therefore to a better aqueous availability. The same four saponins led to similar lipophilicity decreases for (+)-catechin (pattern B) present in common TCM recipes based on *Camellia sinensis* (tea), *Ziziphus jujuba* (Chinese date) and *Cinnamum verum* (cinnamon tree). Chlorogenic acid, one of the active principles of *Andrographis paniculata* (green chiretta), *Salvia miltiorrhiza* (Chinese sage) and *Coptis chinensis* (goldthread) showed drops of logP at the CMCs of α -hederin and β -aescin as in pattern C combinations.

The traditional use of saponin-rich plants in TCM formulations fills a specific role in the hierarchy of ingredients. According to the Jun-Chen-Zuo-Shi classification, plants in the Courier role aim to harmonise and detoxify the formulation. Aside from the biological effects of saponins, this investigation confirms that the physical properties of micelle forming saponins are intimately linked to a modification of behaviour of the other molecules in solution, as seen with the alteration of logP and the four types of interactions presented earlier. These results agree with previous results on the effects of saponins on water solubility like those reported by Schöpke and coworkers.¹⁹

3.4 | The aconitine and artemisinin cases

LogP of aconitine in the presence of glycyrrhizin showed a sharp decrease until the CMC was reached followed by a less pronounced decrease, typical of pattern B behaviour, meaning that the compound became more water soluble with increasing saponin concentration (logP dropped from -0.60 to -1.20 , Figure S36). In TCM, plants of the genus *Aconitum* are almost systematically accompanied in the TCM by licorice root as the Courier.^{20,21} Since aconitine is a notoriously toxic compound, this suggests that lowering the partition coefficient brings more aconitine into the aqueous phase and thus enhances its urinary excretion, reducing its acute toxicity. This is illustrated by the classical herb-pair Fuzi-Gan Cao (*Aconitum carmichaelii*-*Radix* and *Rhizoma Glycyrrhizae*) where the cardiotoxicity of the former is attenuated by the later.²²

Artemisinin, the antimalarial active principle of Qing Hao Su, is so lipophilic that it could only be developed as a single-compound drug as the hemi-succinate (artesanate). It is easily identified by NMR spectroscopy when dissolved in deuterated chloroform (CDCl_3) (singlet at 6.00 ppm for the hemiketal proton), whereas it is almost impossible to detect it in an aqueous phase, rendering our method of logP measurement ineffective. However, after addition of glycyrrhizin, artemisinin's signal became observable and a logP of 1.7 was measured at a 0.28 g/L concentration of the saponin (Figure S37). This last result was also observed in HPLC: the concentration of artemisinin detected in a decoction of 20 g/L of leaves of *Artemisia annua* (retention time 6.1 min) steadily increased after adding glycyrrhizin to the mixture (Figure S38). In other words, saponins had a direct effect on the aqueous extraction of artemisinin by their action on partition coefficients and water solubility. Thus, the relative area percentage of artemisinin in the decoction increased from 3% to 21%, when glycyrrhizin concentration went from 0.08 g/L to 0.32 g/L.

These results explain why the inclusion of licorice roots in traditional formulations has a direct impact on the toxicity and elimination of bioactive molecules such as aconitine, and on the availability of lipophilic molecule such as artemisinin in aqueous formulations.

It remains to understand the nature and the structural aspects of the interactions to predict the behaviour of a given product in the presence of a saponin. Similar interactions prevail in the food domain, which also makes use of complex mixtures of ingredients, as demonstrated by a cryo electron microscopy study of *Quillaja* saponins micelles in the presence of lutein esters.²³ This work is but a small part in the understanding and modernisation of TCM. Recent reviews and reports show major research activity in the promising areas of integrated pharmacology and of multi-component and multi-target mechanisms of actions.^{24,25} Efforts are also being made to find a consensus between traditional use and drug-likeness as ruled by Lipinski and coworkers.^{26,27} Because of the complexity of the mixtures and the high number of possible interactions from pharmacological and chemical standpoints future investigations will require the use of simplified models like ours.

ACKNOWLEDGEMENTS

The authors thank Dr Pedro Lameiras (ICMR) for help with NMR Water Suppression methods and the ERETIC 2 tool. The Agence Régionale de Santé (ARS) Grand Est as well as the Centre Hospitalier Universitaire (CHU) of Reims (France) are also acknowledged for providing a thesis grant to Cédric Barthel. Numerous discussions with Professor Joseph Connolly on the content and conformity of this article are deeply appreciated.

DATA AVAILABILITY STATEMENT

Data available upon request.

ORCID

Georges Massiot  <https://orcid.org/0000-0002-3507-8559>

Catherine Lavaud  <https://orcid.org/0000-0001-7894-9388>

REFERENCES

- Wu L, Wang Y, Li Z, Zhang B, Cheng Y, Fan X. Identifying roles of «Jun-Chen-Zuo-Shi» component herbs of QiShenYiQi formula in treating acute myocardial ischemia by network pharmacology. *Chinas Med*. 2014;9(1):24. doi:10.1186/1749-8546-9-24
- Jia W, Gao WY, Yan YQ, et al. The rediscovery of ancient chinese herbal formulas. *Phytother Res*. 2004;18(8):681-686. doi:10.1002/ptr.1506
- Tu Y. The discovery of artemisinin (qinghaosu) and gifts from Chinese medicine. *Nat Med*. 2011;17(10):1217-1220. doi:10.1038/nm.2471
- Zhao J, Jiang P, Zhang W. Molecular networks for the study of TCM pharmacology. *Brief Bioinform*. 2010;11(4):417-430. doi:10.1093/bib/bbp063
- Wendy Hsiao WL, Liu L. The role of traditional chinese herbal medicines in cancer therapy – from TCM theory to mechanistic insights. *Planta Med*. 2010;76(11):1118-1131. doi:10.1055/s-0030-1250186
- Nabirotkin S, Peluffo AE, Rinaudi P, Yu J, Hajj R, Cohen D. Next-generation drug repurposing using human genetics and network biology. *Curr Op Pharmacol*. 2020;51:78-92. doi:10.1016/j.coph.2019.12.004
- TCM Wiki. List of prescriptions. <https://tcmwiki.com/wiki/prescriptions>. Last accessed July 2022.
- Liu J, Henkel T. Traditional chinese medicine (TCM): are polyphenols and saponins the key ingredients triggering biological activities? *Curr Med Chem*. 2002;9(15):1483-1485. doi:10.2174/092986702369709
- Barthel C, Massiot G, Lavaud C. An easy-to-use and general NMR method for the determination of partition coefficients of drugs and natural products. *Magn Reson Chem*. 2021;59(8):835-843. doi:10.1002/mrc.5159
- Böttcher S, Drusch S. Saponins – self-assembly and behaviour at aqueous interfaces. *Adv Colloids Interface Sc*. 2017;243:105-113. doi:10.1016/j.cis.2017.02.008
- du Noüy PL. An interfacial tensiometer for universal use. *J Gen Physiol*. 1925;7(5):625-631. doi:10.1085/jgp.7.5.625
- Schöpke T, Bartlakowski J, Hiller K. Critical micellar concentrations of a number of triterpenoid saponins of different structure type. *Pharmazie*. 1995;50:711.
- Böttger S, Hofmann K, Melzig MF. Saponins can perturb biological membranes and reduce the surface tension of aqueous solutions: a correlation? *Biorg Med Chem*. 2012;20(9):2822-2828. doi:10.1016/j.bmc.2012.03.032
- Decroos K, Vincken JP, van Koningsveld GA, Gruppen H, Verstraete W. Preparative chromatographic purification and surfactant properties of individual soyasaponins from soy hypocotyls. *Food Chem*. 2007;101:324-333. doi:10.1016/j.foodchem.2006.01.041
- Reichert CL, Salminen H, Weiss J. Quillaja saponin characteristics and functional properties. *Annu Rev Food Sci Technol*. 2019;10(1):43-73. doi:10.1146/annurev-food-032818-122010
- Lipinski CA, Lombardo F, Dominy BW, Feeney PJ. Experimental and computational approaches to estimate solubility and permeability in drug discovery and development settings. *Adv Drug Deliv Rev*. 1997;23(1-3):3-25. doi:10.1016/S0169-409X(96)00423-1
- Harutyunyan LR, Lachinyan ML, Harutyunyan RS. Effect of ascorbic acid on the colloidal and micellar properties of anionic, cationic, and nonionic surfactants: conductivity, volumetric, viscometric, and fluorescence study. *J Chem Eng Data*. 2013;58(11):2998-3008. doi:10.1021/je400667c
- Alam MS, Siddiq AM, Mandal AB. Thermodynamic and micellization studies of a cationic gemini surfactant 16-6-16: influence of ascorbic acid and temperature. *Colloid J*. 2016;78(1):9-14. doi:10.1134/S1061933X16010026
- Walther U, Dittich K, Gelbrich G, Schöpke T. Effects of Saponins on the water solubility of different model compounds. *Planta Med*. 2001;67(1):49-54. doi:10.1055/s-2001-10876
- Singhuber J, Zhu M, Prinz S, Kopp B. Aconitum in traditional Chinese medicine – a valuable drug or an unpredictable risk? *J Ethnopharmacol*. 2009;126(1):18-30. doi:10.1016/j.jep.2009.07.031
- Lu G, Dong Z, Wang Q, et al. Toxicity assessment of nine types of decoction pieces from the daughter root of *Aconitum carmichaeli* (Fuji) based on the chemical analysis of their diester diterpenoid alkaloids. *Planta Med*. 2010;76(08):825-830. doi:10.1055/s-0029-1240688
- Zhang JM, Liao W, He YX, He Y, Yan D, Fu CM. Study on intestinal absorption and pharmacokinetic characterization of diester diterpenoid alkaloids in precipitation derived from Fuji-Gancao herb-pair decoction for its potential interaction mechanism investigation. *J Ethnopharmacol*. 2013;147(1):128-135. doi:10.1016/j.jep.2013.02.019
- Tippel J, Lehmann M, von Klitzing R, Drusch S. Interfacial properties of Quillaja saponins and its use for micellization of lutein esters. *Food Chem*. 2016;212:35-42. doi:10.1016/j.foodchem.2016.05.130
- Chen M, Yang F, Yang X, Lai X, Gao Y. Systematic understanding of mechanisms of a Chinese herbal formula in treatment of metabolic syndrome by an integrated pharmacology approach. *Int J Mol Sci*. 2016;17(12):2114. doi:10.3390/ijms17122114
- Guan D, Xuan B, Wang C, et al. Improving the physicochemical and biopharmaceutical properties of active pharmaceutical ingredients derived from traditional Chinese medicine through cocrystal. *Pharmaceutics*. 2021;13(12):2160. doi:10.3390/pharmaceutics13122160
- Yang J. Application of computer-aided drug design to traditional Chinese medicine. *Int J Org Chem*. 2013;03(01):1-16. doi:10.4236/ijoc.2013.31A001
- Shen M, Tian S, Li Y, Xu X, Wang J, Hou T. Drug-likeness analysis of traditional Chinese medicines: 1. Property distribution of drug-like compounds, non-drug like compounds and natural compounds from traditional Chinese medicines. *J Chem*. 2012;4(1):31. doi:10.1186/1758-2946-4-31

SUPPORTING INFORMATION

Additional supporting information can be found online in the Supporting Information section at the end of this article.

How to cite this article: Barthel C, Massiot G, Lavaud C. Traditional Chinese medicine: saponins, critical micellar concentrations and partition coefficients. *Phytochemical Analysis*. 2023;1-7. doi:10.1002/pca.3221