



**HAL**  
open science

## High Blood Eosinophil Count at Stable State is Not Associated with Airway Microbiota Distinct Profile in COPD

Jeanne-Marie Perotin, Anaëlle Muggeo, Quentin Lecomte-Thenot, Audrey Brisebarre, Sandra Dury, Claire Launois, Julien Ancel, Valérian Dormoy, Thomas Guillard, Gaëtan Deslee

► **To cite this version:**

Jeanne-Marie Perotin, Anaëlle Muggeo, Quentin Lecomte-Thenot, Audrey Brisebarre, Sandra Dury, et al.. High Blood Eosinophil Count at Stable State is Not Associated with Airway Microbiota Distinct Profile in COPD. *International Journal of Chronic Obstructive Pulmonary Disease*, 2024, Volume 19, pp.765-771. 10.2147/COPD.S453526 . hal-04530339

**HAL Id: hal-04530339**

**<https://hal.univ-reims.fr/hal-04530339>**

Submitted on 3 Apr 2024

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

1 ORIGINAL RESEARCH

2

3 Perotin et al

4

5

6 **Title: High blood eosinophil count at stable state is not associated with airway**

7 **microbiota distinct profile in COPD**

8

9 **Authors:** Jeanne-Marie PEROTIN<sup>1,2</sup>, Anaëlle MUGGEO<sup>3</sup>, Quentin LECOMTE<sup>3</sup>, Audrey

10 BRISEBARRE<sup>1</sup>, Sandra DURY<sup>2</sup>, Claire LAUNOIS<sup>1,2</sup>, Julien ANCEL<sup>1,2</sup>, Valérian DORMOY<sup>1</sup>,

11 Thomas GUILLARD<sup>3</sup>, Gaëtan DESLEE<sup>1,2</sup>

12

13 **Affiliations :**

14 <sup>1</sup> Université de Reims Champagne-Ardenne, INSERM UMR-S 1250, P3Cell, Reims, France

15 <sup>2</sup> CHU de Reims, Service des Maladies Respiratoires, Reims, France

16 <sup>3</sup> Université de Reims Champagne-Ardenne, INSERM UMR-S 1250, P3Cell, CHU de Reims,

17 Laboratoire de bactériologie-Virologie-Hygiène hospitalière-Parasitologie-Mycologie, Reims,

18 France

19

20

21

22 **Abstract**

23 **Purpose:** The heterogeneity of clinical features in COPD at stable state has been associated  
24 with airway microbiota. Blood eosinophil count (BEC) represents a biomarker for a pejorative  
25 evolution of COPD, including exacerbations and accelerated FEV<sub>1</sub> decline. We aimed to  
26 analyse the associations between BEC and airway microbiota in COPD at stable state.

27 **Patients and methods:** Adult COPD patients at stable state (RINNOPARI cohort) were  
28 included and characterised for clinical, functional, biological and morphological features. BEC  
29 at inclusion defined 2 groups of patients with low BEC <300/mm<sup>3</sup> and high BEC ≥300/mm<sup>3</sup>.  
30 Sputa were collected and an extended microbiological culture was performed for the  
31 identification of viable airway microbiota.

32 **Results:** Fifty-nine subjects were included. When compared with the low BEC (n=40,  
33 67.8%), the high BEC group (n=19, 32.2%) had more frequent exacerbations (p<0.001) and  
34 more pronounced cough and sputum (p<0.05). The global composition, the number of  
35 bacteria per sample and the α-diversity of the microbiota did not differ between groups, as  
36 well as the predominant phyla (Firmicutes), or the gender repartition.

37 **Conclusion:** In our study, high BEC in COPD at stable state was associated with a clinical  
38 phenotype including frequent exacerbation, but no distinct profile of viable airway microbiota  
39 compared with low BEC.

40

41 **Key words:** COPD, Eosinophil, sputum, microbiota

42

43 **Correspondance :**

44 Jeanne-Marie Perotin, Service des Maladies Respiratoires, CHU Maison Blanche, Inserm  
45 UMR-S 1250, 45 rue Cognacq-Jay, 51092 Reims cedex, France.

46 Email: [jmperotin-collard@chu-reims.fr](mailto:jmperotin-collard@chu-reims.fr)

**47 Introduction**

48 Chronic obstructive pulmonary disease (COPD) is characterized by chronic respiratory  
49 symptoms including dyspnea, cough, sputum production, and exacerbations, due to  
50 abnormalities of the airways and/or alveoli that cause persistent, often progressive airflow  
51 limitation (1). COPD is a heterogeneous disease regarding clinical features, with high  
52 morbidity and mortality. The pathogenesis of COPD is still not completely elucidated,  
53 including structural and inflammatory changes (1). Blood eosinophil count (BEC) has been  
54 recently identified as a biomarker of interest in stable COPD. Greater BEC levels have been  
55 associated with greater FEV<sub>1</sub> decline in the absence of inhaled corticosteroid (ICS)  
56 treatment, and with the risk of exacerbation (1, 2). It is currently recommended to use BEC,  
57 not as a standalone biomarker but together with exacerbation history, to identify patients with  
58 the greatest likelihood of ICS treatment benefit (1, 3).

59 Airway microbiota at stable state might participate in the heterogeneity of COPD clinical  
60 presentation. Previous studies performed in a stable state identified associations between  
61 the sputum microbiome and COPD phenotypes (4-6). Associations with COPD inflammatory  
62 phenotypes including eosinophils have been recently investigated, suggesting that different  
63 profiles of bacteria phyla might be associated with different profiles of inflammation: low  
64 sputum eosinophil count in the presence of potential pathogenic microorganism (PPM) (7, 8),  
65 more diverse respiratory microbiome in subjects with BEC  $\geq 2\%$  (9). However, the  
66 relationships between BEC and airway microbiota and its clinical implications remain to be  
67 determined.

68 The most frequently used techniques for airway microbiota analyses (PCR amplification,  
69 sequencing of the bacterial 16S ribosomal RNA gene) do not involve bacterial culture. We  
70 previously described an extended conventional culture-based approach, allowing the  
71 identification of bacteria restricted to viable strains (6, 10). In this study, we applied this  
72 culture technique and compared viable airway microbiota in COPD patients at a stable state  
73 with high and low BEC.

74

75 **Methods**

## 76 Patients

77 Patients with mild to severe COPD were included prospectively in the RINNOPARI cohort  
78 (Recherche et INNOvation en PAthologie Respiratoire Inflammatoire; University Hospital of  
79 Reims, France; NCT02924818) as described previously (6) The regional ethics committee  
80 approved the study (Comité de Protection des Personnes—Dijon EST I, no. 2016-A00242-  
81 49). All patients provided written consent. Exclusion criteria were other pulmonary diseases,  
82 including asthma, bronchiectasis, cystic fibrosis, bronchopulmonary allergic aspergillosis,  
83 (CF), or pulmonary fibrosis. Stable state was defined as 4 weeks or more from the last  
84 exacerbation (defined as an acute worsening of respiratory symptoms resulting in additional  
85 therapy (6, 11). COPD was defined by postbronchodilator  $FEV_1/FVC < 70\%$ . The severity of  
86 COPD was determined by spirometric classification (GOLD 1:  $FEV_1 \geq 80\%$ ; GOLD 2:  $50\% \leq$   
87  $FEV_1 < 80\%$ ; GOLD 3:  $30\% \leq FEV_1 < 50\%$ ; GOLD 4:  $FEV_1 < 30\%$ ), and ABE classification  
88 depending on exacerbations, dyspnea and CAT score (1). Emphysema was visually  
89 assessed and quantified from CT scan images as previously described (12, 13). BEC was  
90 performed at inclusion, defining 2 groups of patients with low BEC  $< 300/mm^3$  and high BEC  
91  $\geq 300/mm^3$ , using the cut-off proposed by GOLD 2023 (1).

## 92 Sputum analysis

93 Sputa were collected and an extended microbiological culture was performed, as previously  
94 described (6). After liquefaction by N-acetylcysteine, serial dilutions (from 1/1,000 to  
95 1/100,000) of the sputum were performed and cultured at 37°C for 48 h (aerobic cultures), or  
96 5% CO<sub>2</sub> for 5 days (anaerobic cultures) on several agar media including Columbia blood,  
97 chocolate, Schaedler, and *Pseudomonas*-selective ceftrimide (Thermo Fisher Scientific, USA)  
98 We next quantified all morphologically distinct colonies as colony-forming unit (CFU) per  
99 milliliter. Colonies were then identified using MALDI-TOF mass spectrometry (MALDI

100 Biotyper<sup>®</sup>, Bruker Daltonics, Germany). We estimated the viable airway microbiota  $\alpha$ -diversity  
101 using the Shannon index.

## 102 Statistical Analysis

103 The descriptive data are expressed as numbers (percentages), median [25<sup>th</sup>-75<sup>th</sup> quartiles],  
104 or mean values  $\pm$  standard deviation, when appropriate. Qualitative variables were compared  
105 by the chi-square test or Fisher exact test. Quantitative variables were compared using the t-  
106 test or Mann–Whitney test. A p-value  $< 0.05$  was considered significant. The dissimilarities in  
107 bacterial communities between low BEC  $<300/\text{mm}^3$  and high BEC  $\geq 300/\text{mm}^3$  groups were  
108 visualised in a low-dimensional Euclidean space, using unsupervised principal component  
109 analysis (PCA), that was plotted along the first two principal components (the two explaining  
110 most of the variance).

111

## 112 Results

### 113 Patients

114 Fifty-nine subjects were included in the study, 40 in the low BEC group (BEC $<300/\text{mm}^3$ ,  
115 67.8%) and 19 in the high BEC group (BEC  $\geq 300/\text{mm}^3$ , 32.2%). Subjects' characteristics are  
116 detailed in Table 1 and Suppl Table 1. Briefly, they were predominantly men (57.6%), mean  
117 age  $61 \pm 9$  years. COPD was severe to very severe in 57.6%, 66.1% had one or more  
118 exacerbations in the last year, with a median of 2 exacerbations. Two patients received long  
119 term oral corticosteroids at inclusion, including one in low BEC group, and one in high BEC  
120 group, with no significant differences between groups (Table 1). When compared with the  
121 low BEC group, the high BEC group was characterised by a higher number of exacerbations  
122 per patient (median 2 [0-4] vs 1 [0-3],  $p<0.001$ ), more pronounced symptoms of cough and  
123 sputum ( $p<0.05$ , CASA-Q, Table 1) and a trend for lower smoking exposure ( $p=0.06$ ) and  
124 more frequent antibiotic course ( $p=0.06$ ). We did not find any differences in terms of inhaled  
125 treatment, lung function, or emphysema score.

## 126 Microbiology

127 The extended culture method was applied to 59 sputa (1 sample per patient) to determine  
128 the viable airway microbiota. We identified 386 bacteria, from 71 different species,  
129 representing a mean of 6.6 bacteria per sample (Table S2). We compared the viable  
130 microbiota of the low and high BEC groups by PCA analysis and we found no difference in  
131 the global composition of the microbiota of the 2 groups (Figure 1A). The number of bacteria  
132 per sample and the  $\alpha$ -diversity of the microbiota did not differ between low and high BEC  
133 groups (Figure 1B and C). The repartition of the bacterial phyla was similar in the 2 groups,  
134 with a predominance of Firmicutes (Figure 1D). We observed the same repartition of different  
135 genera: *Streptococcus*, *Rothia*, *Veillonella*, *Neisseria*, and *Actinomyces* were predominant  
136 and represented more than 65% of the bacteria identified (Figure 1E). The prevalence of the  
137 different species in the 2 groups was analysed and no difference was found between the low  
138 and high BEC groups (Figure 1F). The most common bacteria in both groups were  
139 *Streptococcus oralis/mitis/pneumoniae*, identified in more than 90% of samples, followed by  
140 *Veillonella parvula/dispar/atypica* found in more than 50% of samples. Bacterial  
141 quantifications of the viable microbiota ranged from  $10^2$  CFU/mL to  $10^9$  CFU/mL, with a  
142 median of  $10^6$  CFU/mL and no difference between the 2 groups (Table S2).

143 In this cohort of COPD patients at stable state, some PPM were detected: *Staphylococcus*  
144 *aureus* (n = 8, 13,6%), *Haemophilus influenzae* (n = 6, 10.2%), *Moraxella catarrhalis* (n = 5,  
145 8.5%) and *Pseudomonas aeruginosa* (n = 3, 5.1%), with the same prevalence in the 2  
146 groups (n=16; 40.0% in the low BEC group vs n= 7, 36.8% in the high BEC group) (Table  
147 S2).

148

149 **Discussion**

150 In this cross-sectional analysis focusing on COPD patients at stable state and using an  
151 original extended conventional culture-based approach that allows the identification of

152 bacteria restricted to viable strains, we identified that high BEC is not associated with a  
153 distinct profile of viable airway microbiota, despite a BEC-associated clinical phenotype.

154 In our study, the high BEC group was characterised by a clinical phenotype including more  
155 frequent exacerbations in the last year, a trend for a more frequent antibiotic course, and  
156 more pronounced cough and sputum symptoms (CASA-Q). Elevated BEC at a stable state  
157 has been previously identified as a biomarker for exacerbation risk in COPD, with an  
158 incidence rate ratio of exacerbation of 1.32 for BEC  $\geq 300/\text{mm}^3$  in the COPDgene study (2).  
159 An association between sputum inflammatory cells and symptoms at stable state has been  
160 shown, with neutrophilic inflammation being associated with cough, and eosinophilic  
161 inflammation with dyspnea (14, 15). A recent post-hoc analysis suggested that COPD  
162 patients with both high eosinophil levels in sputum ( $\geq 3\%$ ) and chronic bronchitis might  
163 present a distinct profile of gene expression, characterised by the overexpression of T2- and  
164 phosphodiesterase-4-inhibitors-related genes (16). The high BEC group in our study was  
165 further characterised by a trend for lower smoking exposure, but a similarly altered lung  
166 function. This might indirectly reflect the previously observed accelerated lung function  
167 decline in COPD subjects with high eosinophil counts (17).

168 We did not identify a distinct profile of viable airway microbiota in the high BEC group when  
169 compared with the low BEC group. Firmicutes and *Streptococcus* were the predominant  
170 bacteria phylum and species respectively, in line with previous analyses performed in COPD  
171 at a stable state (5, 18). The previously reported decrease in Proteobacteria abundance and  
172 increase in Firmicutes phyla in subjects with high BEC (19), or the more diverse microbiome  
173 in subjects with BEC  $\geq 2\%$  (9) were not found in our study. It must be pointed out that these  
174 previous studies used gene sequencing techniques (16S rRNA), which are not able to  
175 discriminate between viable and non-viable strains. Previous studies described the presence  
176 of PPM in subjects with low sputum eosinophil count (7, 8). However, these studies used  
177 qPCR detection restricted to 3 species (*H. influenzae*, *M. catarrhalis*, and *S. pneumoniae*),



178 therefore also including both viable and non-viable strains. Our study did not confirm those  
179 results.

180 Our study suffers from several limits. Although monocentric, the patient's recruitment was  
181 prospective and all benefited from an in-depth phenotypic characterisation. Exacerbation  
182 frequency was estimated retrospectively in the last year before inclusion. We performed only  
183 a one-point BEC assessment at inclusion, while BEC is known to vary over time in COPD  
184 (20). However, the clinical phenotype of the high BEC group including more frequent  
185 exacerbation matches with previous studies (2). Inhaled treatment was heterogeneous with  
186 32% of the patients using ICS. ICS treatment can alter the microbiome in the small airways  
187 of patients with COPD and might have an impact on our results (21). However, inhaled  
188 treatment strategies including ICS in our study did not significantly differ between high and  
189 low BEC groups. Finally, our extended conventional culture-based approach may have lower  
190 sensitivity than a metagenomic approach for a more in-depth characterisation of the lung  
191 microbiota.

192

## 193 **Conclusion**

194 In our study using an original extended conventional culture-based approach, high BEC in  
195 COPD at a stable state was associated with a clinical phenotype including frequent  
196 exacerbation and more cough and sputum symptoms, but no distinct profile of viable airway  
197 microbiota compared with low BEC.

198

199 **Declarations**

200 **Ethics approval and informed consent**

201 The study was approved by the regional ethics committee (Comité de Protection des  
202 Personnes—Dijon EST I, no. 2016-A00242-49). Informed consent was obtained from all the  
203 patients.

204 **Consent for publication**

205 Not applicable

206 **Data availability**

207 The data that support the findings of this study are available from the corresponding author  
208 upon reasonable request.

209 **Funding**

210 No funding

211 **Competing interests**

212 J.M. Perotin reports lecture honoraria from AstraZeneca, and support for attending meetings  
213 from AstraZeneca and Chiesi; outside the submitted work. C. Launois reports support for  
214 attending meeting from Chiesi; outside the submitted work. V. Dormoy reports lecture  
215 honoraria from Chiesi and AstraZeneca; outside the submitted work. G. Deslée reports  
216 lecture honoraria from Chiesi, AstraZeneca and GlaxoSmithKline; outside the submitted  
217 work. There are no further conflicting interests to disclose.

218 **Authors' contributions**

219 Study concept: J.M. Perotin, A. Muggeo, T. Guillard and G. Deslee; study design: J.M.  
220 Perotin and A. Muggeo; acquisition data: J.M. Perotin, A. Muggeo, Q. Lecomte, S. Dury, C.  
221 Launois, J. Ancel and V. Dormoy; analysis and data interpretation: J.M. Perotin, A. Muggeo,  
222 Q. Lecomte, A. Brisebarre, S. Dury, C. Launois, J. Ancel, V. Dormoy, T. Guillard and G.  
223 Deslee; revision of manuscript: J.M. Perotin, A. Muggeo, Q. Lecomte, A. Brisebarre, S. Dury,  
224 C. Launois, J. Ancel, V. Dormoy, T. Guillard and G. Deslee; manuscript writing: J.M. Perotin,  
225 A. Muggeo, T. Guillard and G. Deslee.

226

227 **References**

- 228 1. Global strategy for the diagnosis, management, and prevention of chronic obstructive  
229 pulmonary disease (2023 report). Available from <https://goldcopd.org>.
- 230 2. Yun JH, Lamb A, Chase R, Singh D, Parker MM, Saferali A, et al. Blood eosinophil  
231 count thresholds and exacerbations in patients with chronic obstructive pulmonary disease. *J*  
232 *Allergy Clin Immunol*. 2018;141(6):2037-47.e10.
- 233 3. Singh D, Agusti A, Martinez FJ, Papi A, Pavord ID, Wedzicha JA, et al. Blood  
234 Eosinophils and Chronic Obstructive Pulmonary Disease: A Global Initiative for Chronic  
235 Obstructive Lung Disease Science Committee 2022 Review. *Am J Respir Crit Care Med*.  
236 2022;206(1):17-24.
- 237 4. Galiana A, Aguirre E, Rodriguez JC, Mira A, Santibañez M, Candela I, et al. Sputum  
238 microbiota in moderate versus severe patients with COPD. *Eur Respir J*. 2014;43(6):1787-  
239 90.
- 240 5. Garcia-Nuñez M, Millares L, Pomares X, Ferrari R, Pérez-Brocal V, Gallego M, et al.  
241 Severity-related changes of bronchial microbiome in chronic obstructive pulmonary disease.  
242 *J Clin Microbiol*. 2014;52(12):4217-23.
- 243 6. Muggeo A, Perotin JM, Brisebarre A, Dury S, Dormoy V, Launois C, et al. Extended  
244 Bacteria Culture-Based Clustering Identifies a Phenotype Associating Increased Cough and.  
245 *Front Microbiol*. 2021;12:781797.7. Beech AS, Lea S, Kolsum U, Wang Z, Miller BE,  
246 Donaldson GC, et al. Bacteria and sputum inflammatory cell counts; a COPD cohort  
247 analysis. *Respir Res*. 2020;21(1):289.
- 248 8. Kolsum U, Donaldson GC, Singh R, Barker BL, Gupta V, George L, et al. Blood and  
249 sputum eosinophils in COPD; relationship with bacterial load. *Respir Res*. 2017;18(1):88.
- 250 9. Millares L, Pascual S, Montón C, García-Núñez M, Lalmolda C, Faner R, et al.  
251 Relationship between the respiratory microbiome and the severity of airflow limitation, history  
252 of exacerbations and circulating eosinophils in COPD patients. *BMC Pulm Med*.  
253 2019;19(1):112.
- 254 10. Dingle TC, Butler-Wu SM. Maldi-tof mass spectrometry for microorganism  
255 identification. *Clin Lab Med*. 2013;33(3):589-609.
- 256 11. Global strategy for the diagnosis, management, and prevention of chronic obstructive  
257 pulmonary disease (2022 report). Available from <https://goldcopd.org>
- 258 12. Perotin JM, Adam D, Vella-Boucaud J, Delepine G, Sandu S, Jonvel AC, et al. Delay  
259 of airway epithelial wound repair in COPD is associated with airflow obstruction severity.  
260 *Respir Res*. 2014;15:151.
- 261 13. Washko GR, Criner GJ, Mohsenifar Z, Scirba FC, Sharafkhaneh A, Make BJ, et al.  
262 Computed tomographic-based quantification of emphysema and correlation to pulmonary  
263 function and mechanics. *COPD*. 2008;5(3):177-86.
- 264 14. Zysman M, Deslee G, Caillaud D, Chanez P, Escamilla R, Court-Fortune I, et al.  
265 Relationship between blood eosinophils, clinical characteristics, and mortality in patients with  
266 COPD. *Int J Chron Obstruct Pulmon Dis*. 2017;12:1819-24.
- 267 15. Contoli M, Baraldo S, Conti V, Gnesini G, Marku B, Casolari P, et al. Airway  
268 inflammatory profile is correlated with symptoms in stable COPD: A longitudinal proof-of-  
269 concept cohort study. *Respirology*. 2020;25(1):80-8.
- 270 16. Singh D, Bassi M, Balzano D, Lucci G, Emirova A, Anna Nandeuil M, et al. COPD  
271 patients with chronic bronchitis and higher sputum eosinophil counts show increased type-2  
272 and PDE4 gene expression in sputum. *J Cell Mol Med*. 2021;25(2):905-18.
- 273 17. Tan WC, Bourbeau J, Nadeau G, Wang W, Barnes N, Landis SH, et al. High  
274 eosinophil counts predict decline in FEV. *Eur Respir J*. 2021;57(5).
- 275 18. Tangedal S, Nielsen R, Aanerud M, Persson LJ, Wiker HG, Bakke PS, et al. Sputum  
276 microbiota and inflammation at stable state and during exacerbations in a cohort of chronic  
277 obstructive pulmonary disease (COPD) patients. *PLoS One*. 2019;14(9):e0222449.

- 278 19. Dicker AJ, Huang JTJ, Lonergan M, Keir HR, Fong CJ, Tan B, et al. The sputum  
279 microbiome, airway inflammation, and mortality in chronic obstructive pulmonary disease. *J*  
280 *Allergy Clin Immunol.* 2021;147(1):158-67.
- 281 20. Schumann DM, Tamm M, Kostikas K, Stolz D. Stability of the Blood Eosinophilic  
282 Phenotype in Stable and Exacerbated COPD. *Chest.* 2019;156(3):456-65.
- 283 21. Yip W, Li X, Koelwyn GJ, Milne S, Leitao Filho FS, Yang CX, et al. Inhaled  
284 Corticosteroids Selectively Alter the Microbiome and Host Transcriptome in the Small  
285 Airways of Patients with Chronic Obstructive Pulmonary Disease. *Biomedicines.* 2022;10(5).

286

287

288 **Tables**

289 Table 1. Patients' characteristics

	Total	Low BEC	High BEC	p-value
n	59	40	19	
Age, yrs	61.0 ± 9.3	61.3 ± 9.6	60.4 ± 8.9	0.365
Men	34 (57.6)	22 (55.0)	12 (63.2)	0.380
BMI, kg/m <sup>2</sup>	25.8 ± 5.7	25.7 ± 6.1	25.9 ± 4.8	0.440
Current smoker	21 (35.6)	15 (37.5)	6 (31.6)	0.443
Smoking history, pack-year	43.4 ± 19.5	46.1 ± 18.6	37.6 ± 20.5	0.059
Inhaled treatment strategy				
No inhaled treatment	11 (18.6)	5 (12.5)	6 (31.6)	
LABA or LAMA	5 (8.5)	4 (10.0)	1 (5.3)	
LABA + LAMA	24 (40.7)	18 (45.0)	6 (31.6)	0.204
ICS + LABA or ICS +LAMA	5 (8.5)	2 (5.0)	3 (15.8)	
ICS + LAMA + LABA	14 (23.7)	11 (27.5)	3 (15.8)	
Long-term oral corticosteroids	2 (3.4)	1 (2.5)	1 (5.3)	0.643
Long-term oxygen	10 (17.0)	7 (17.5)	3 (15.8)	0.561
Exacerbation in the previous year, n	39 (66.1)	27 (67.5)	12 (63.2)	0.481
Number per patient	1 [0-3]	1 [0-3]	2 [0-4]	0.001
Antibiotics (6 Mo, nb/patient)	1 [0-1]	1 [0-1]	1 [0-3]	0.06
Dyspnea mMRC ≥ 2	44 (77.2)	31 (81.6)	13 (68.4)	0.323
CAT score	18.4 ± 7.4	17.7 ± 7.4	19.8 ± 7.7	0.165
CASA-Q scores				
Symptoms: cough	65.5 ± 24.0	70.2 ± 23.2	56.1 ± 23.2	0.018
Symptoms: sputum	64.9 ± 25.9	69.5 ± 26.8	55.7 ± 21.9	0.029
Impact: cough	72.4 ± 24.4	74.6 ± 26.4	67.9 ± 19.5	0.168
Impact: sputum	75.2 ± 22.3	78.2 ± 22.8	69.3 ± 20.6	0.079
Blood eosinophils, G/L	0.2 [0.1-0.3]	0.1 [0.1-0.2]	0.3 [0.3-0.4]	< 0.001
Total IgE, IU/mL	76 [20-256]	54 [18-339]	102 [45-418]	0.267
Positive fungal sputum analysis	20 (34.5)	16 (40.0)	4 (21.0)	0.239

290 Values are n (%), mean ± SD and median [25<sup>th</sup>-75<sup>th</sup>]. BMI: body mass index; LABA: long-  
291 acting beta agonist; LAMA: long-acting muscarinic antagonist; ICS: inhaled corticosteroid;  
292 CAT: COPD assessment test

293

294 **Figure**

295 Figure 1: Bacterial analysis of airway microbiota in low and high BEC groups. A: Principal  
296 component analysis (PCA) of the airway microbiota. The axes are the first eigenvalues, the  
297 ones explaining the most variance of the dataset. Individual patients are represented in blue  
298 triangles (high BEC group, n=19) and green dots (low BEC group, n=40). The largest blue  
299 triangle and red dot at the center of the ellipses represent the 95% confidence interval. B:  
300 Number of species per sample. C: Alpha diversity of viable microbiota (Shannon index). D:  
301 Phyla distribution. E: Genus distribution. F: Bacteria prevalence (note: bacteria with less than  
302 5% frequency are not listed)